

NOTE ON THE FUNDAMENTAL THEOREM ON IRREDUCIBLE NON-NEGATIVE MATRICES

by HANS SCHNEIDER
 (Received 12th February 1958)

1. Let $A = [a_{ij}]$ be an n -th order irreducible non-negative matrix. As is very well-known, the matrix A has a positive characteristic root ρ (provided that $n > 1$), which is simple and maximal in the sense that every characteristic root λ satisfies $|\lambda| \leq \rho$, and the characteristic vector x belonging to ρ may be chosen positive. These results, originally due to Frobenius, have been proved by Wielandt (4) by means of a strikingly simple basic idea. Recently, a variant of Wielandt's proof has been given by Householder (2).

We shall sketch part of the proof. For each non-negative column vector y we set

$$\rho_*(y) = \sup r : ry \leq Ay, \dots\dots\dots(1)$$

$$\rho^*(y) = \inf r : ry \geq Ay. \dots\dots\dots(2)$$

For strictly positive y , we may replace (1) and (2) by

$$\rho_*(y) = \min_i \frac{(Ay)_i}{y_i} = \min_i \frac{\sum_{j=1}^n a_{ij}y_j}{y_i}, \dots\dots\dots(1')$$

$$\rho^*(y) = \max_i \frac{(Ay)_i}{y_i} = \max_i \frac{\sum_{j=1}^n a_{ij}y_j}{y_i} \dots\dots\dots(2')$$

Let P be the section of the non-negative cone (all $y_i \geq 0$) by the plane $\sum_{i=1}^n y_i = 1$. Since $\rho_*(\lambda y) = \rho_*(y)$, for all positive λ , the supremum of $\rho_*(y)$ over P equals the supremum of $\rho_*(y)$ over all $y \geq 0$. On P , $\rho_*(y)$ attains this supremum, say $\rho = \sup \rho_*(y) = \rho_*(x)$, where x is on P . It is then shown that $Ax = \rho x$, that ρ is simple and maximal, and that $x > 0$. Similarly the infimum of $\rho^*(y)$ over all $y \geq 0$ is attained on P .

2. In this argument there arises a dilemma :

(i) *Either* the whole of P is considered, in which case $\rho_*(y)$ (or $\rho^*(y)$) may have singularities and discontinuities at vectors y which have zero elements ;

(ii) *Or*, the subset P_1 of P , consisting of all positive y on P , is considered, in which case $\rho_*(y)$ (and $\rho^*(y)$) are everywhere continuous on P_1 , but P_1 is not closed.

In either case, some justification is required for the assertion that $\rho_*(y)$ attains its supremum (or $\rho^*(y)$ its infimum).

For an example of a discontinuity in $\rho_*(y)$, examine the case of

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 2 \\ 1 & 0 & 2 \end{bmatrix}$$

At $z = (0, 0, 1)$, we have $\rho_*(z) = 2$. But for all y of form $(\alpha, \alpha, 1 - 2\alpha)$ with $0 < \alpha \leq \frac{1}{4}$, $\rho_*(y) = 1$.

It may however be shown that $\rho_*(y)$ is upper semi-continuous on P , and hence $\rho_*(y)$ attains its supremum on P . For $\rho^*(y)$ slightly more complex reasoning is required. If R_1 denotes the real line with infinity adjoined, topologised in the normal way, then $\rho^*(y)$ is a lower semi-continuous function into R_1 , for all non-negative A , and is even continuous when A is irreducible. Again it follows that $\rho^*(y)$ attains its infimum, on P .

3. In this note we shall demonstrate an alternative method of resolving the dilemma. We begin by proving an inequality, which is of some intrinsic interest when applied to the characteristic vector x .

An inequality: Let A be an n -th order non-negative irreducible matrix, κ a least diagonal element of A , and λ a least non-vanishing non-diagonal element. Let y be a positive column vector, and suppose that $y_1 \geq y_2 \geq \dots \geq y_n$. If $\rho^*(y) \leq M$, then

$$\frac{y_n}{y_1} \geq \left(\frac{\lambda}{M - \kappa} \right)^{n-1} \dots \dots \dots (3)$$

Proof: Let $1 \leq k < n$. Then for all $i > k$,

$$My_i \geq \sum_{j=1}^n a_{ij} y_j \geq \sum_{j=1}^k a_{ij} y_j + a_{ii} y_i$$

whence

$$(M - \kappa)y_{k+1} \geq (M - \kappa)y_i \geq (\sum_{j=1}^k a_{ij})y_k \dots \dots \dots (4)$$

Since A is irreducible, here is at least one $i > k$ for which $\sum_{j=1}^k a_{ij} \geq \lambda > 0$. Hence it follows from (4) that

$$\frac{y_{k+1}}{y_k} \geq \frac{\lambda}{M - \kappa} \dots \dots \dots (5)$$

The inequality (3) is obtained from (5) by setting $k = 1, \dots, n - 1$ and multiplying.

Corollary: If, in addition, $\sum_{i=1}^n y_i = 1$, then for all i

$$y_i \geq \frac{1}{n} \left(\frac{\lambda}{M - \kappa} \right)^{n-1} \dots \dots \dots (6)$$

For

$$1 = \sum_{i=1}^n y_i \leq ny_1 \leq n \left(\frac{M - \kappa}{\lambda} \right)^{n-1} y_n$$

4. We now turn to the proof of the fundamental theorem. We shall consider $\rho^*(y)$ only. Set

$$R = \max_i \sum_{j=1}^n a_{ij}$$

and

$$\delta = \frac{1}{n} \left(\frac{\lambda}{R - \kappa} \right)^{n-1}$$

Clearly $R \geq k + \lambda$, whence

$$\delta \leq \frac{1}{n} \dots \dots \dots (7)$$

Let P_2 be that part of the plane section of the non-negative cone defined by $\sum_{i=1}^n y_i = 1$, and $y_i \geq \delta$, for all i . Evidently P_2 is bounded and closed, $\rho^*(y)$ is continuous everywhere on P_2 , and hence $\rho^*(y)$ attains its infimum over P_2 , say at the vector x on P_2 . Set $\rho = \rho^*(x) = \inf \rho^*(y)$ over P_2 .

We note that ρ is also the infimum of $\rho^*(y)$ over all positive y . It is sufficient to consider the infimum on the plane section P_1 , and to show that there exists a z on P_2 for which $\rho^*(z) < \rho^*(y)$ for all y on P_1 which do not lie on P_2 . By (7), the vector $z = \frac{1}{n}(1, 1, \dots, 1)$ belongs to P_2 , while by (6),

$$\rho^*(y) > R = \rho^*(z) \geq \rho, \dots\dots\dots(8)$$

if y belongs to P_1 but not to P_2 .

5. In this section we shall show that ρ is a characteristic root of A , and that the positive vector x is a characteristic vector belonging to ρ , viz. that $Ax = \rho x$. We shall prove the equivalent proposition: *If $z > 0$ and $\rho_*(z) < \rho^*(z)$, then $\rho < \rho^*(z)$.* Our proof is entirely due to Householder (2). Suppose that

$$\frac{(Az)_i}{z_i} = \rho^*(z), \quad i = 1, \dots, k < n \dots\dots\dots(9)$$

$$\frac{(Az)_i}{z_i} < \rho^*(z), \quad i = k + 1, \dots, n. \dots\dots\dots(10)$$

Since A is irreducible there exists a positive element a_{pq} with $1 \leq p \leq k$ and $k + 1 \leq q \leq n$. Define the vector z' by setting $z'_i = z_i$ if $i \neq q$ and $0 < z'_q < z_q$, where z'_q is chosen sufficiently close to z_q to ensure that

$$\frac{(Az')_q}{z'_q} < \rho^*(z). \dots\dots\dots(11)$$

But for all $i \neq q$

$$\frac{(Az')_i}{z'_i} \leq \frac{(Az)_i}{z_i} \dots\dots\dots(12)$$

It follows from (9), (10), (11) and (12) that

$$\rho^*(z') \leq \rho^*(z).$$

Since

$$\frac{(Az')_p}{z'_p} < \frac{(Az)_p}{z_p}$$

the equality

$$\frac{(Az')_i}{z'_i} = \rho^*(z)$$

can hold for at most $k - 1$ indices i . Thus, by repetition of this process we may construct a vector z'' satisfying $\rho^*(z'') < \rho^*(z)$. We deduce that $\rho < \rho^*(z)$.

6. If ρ is not a simple characteristic root of A , suppose first that ρ has the linearly independent characteristic vectors $x > 0$ and z belonging to it. By choosing the real numbers α and β suitably, we may obtain a positive characteristic vector $\alpha x + \beta z$ on P_i belonging to ρ , which has some element less than δ . But this is impossible, by (6) and (8). Next, suppose that ρ is not simple, but has only one linearly independent characteristic vector belonging to it.

Then there exists a column vector y satisfying

$$Ay = \rho y - x, \dots\dots\dots(13)$$

and replacing y by $y + \gamma x$, it follows there exists a positive y satisfying (13). For this y , $\rho^*(y) < \rho$, which is impossible. We have proved that ρ is a simple characteristic root.

7. For the sake of completeness, we add a standard proof that ρ is a maximal characteristic root. Let σ be any characteristic root of A and u a row vector satisfying $uA = \sigma u$. Then, for all j ,

$$|\sigma| |u_j| = \left| \sum_{i=1}^n u_i a_{ij} \right| \leq \sum_{i=1}^n |u_i| |a_{ij}|$$

whence $|\sigma| \sum_{j=1}^n |u_j| x_j \leq \sum_{i,j=1}^n |u_i| |a_{ij}| x_j = \rho \sum_{i=1}^n |u_i| x_i$,

and so $|\sigma| \leq \rho$,

since $\sum_{i=1}^n |u_i| x_i > 0$.

8. The inequality (3) leads to a positive lower bound for the ratios of the elements of the characteristic vector x . In view of $\rho^*(x) = \rho$ and (8) it follows that

$$\frac{\min_i x_i}{\max_i x_i} \geq \left(\frac{\lambda}{\rho - \kappa} \right)^{n-1} \geq \left(\frac{\lambda}{R - \kappa} \right)^{n-1}.$$

For the case $A > 0$, lower bounds for the ratios x_j/x_i have already been found by Ostrowski (3) and A. Brauer (1). These bounds, however, reduce to 0 when A has zero elements.

REFERENCES

(1) A. BRAUER, The theorems of Ledermann and Ostrowski on positive matrices, *Duke Math. J.*, **24** (1957), 265-274.
 (2) A. S. HOUSEHOLDER, *On the convergence of matrix iterations*, Oak Ridge National Laboratory, Physics No. 1883 (1955).
 (3) A. OSTROWSKI, Bounds for the greatest latent root of a positive matrix, *J. London Math. Soc.*, **27** (1952), 253-256.
 (4) H. WIELANDT, Unzerlegbare, nicht negative Matrizen, *Math. Zeitschrift*, **52** (1950), 642-8.

THE QUEEN'S UNIVERSITY
 BELFAST