

# HAAR FRAME CHARACTERIZATIONS OF BESOV-SOBOLEV SPACES AND OPTIMAL EMBEDDINGS INTO THEIR DYADIC COUNTERPARTS

GUSTAVO GARRIGÓS    ANDREAS SEEGER    TINO ULLRICH

ABSTRACT. We study the behavior of Haar coefficients in Besov and Triebel-Lizorkin spaces on  $\mathbb{R}$ , for a parameter range in which the Haar system is not an unconditional basis. First, we obtain a range of parameters, extending up to smoothness  $s < 1$ , in which the spaces  $F_{p,q}^s$  and  $B_{p,q}^s$  are characterized in terms of doubly oversampled Haar coefficients (Haar frames). Secondly, in the case that  $1/p < s < 1$  and  $f \in B_{p,q}^s$ , we actually prove that the usual Haar coefficient norm,  $\|\{2^j \langle f, h_{j,\mu} \rangle\}_{j,\mu}\|_{b_{p,q}^s}$  remains equivalent to  $\|f\|_{B_{p,q}^s}$ , i.e., the classical Besov space is a closed subset of its dyadic counterpart. At the endpoint case  $s = 1$  and  $q = \infty$ , we show that such an expression gives an equivalent norm for the Sobolev space  $W_p^1(\mathbb{R})$ ,  $1 < p < \infty$ , which is related to a classical result by Bočkarëv. Finally, in several endpoint cases we clarify the relation between dyadic and standard Besov and Triebel-Lizorkin spaces.

## 1. INTRODUCTION AND STATEMENT OF MAIN RESULTS

In this paper we investigate the validity of norm characterizations for elements  $f$  in Besov and Triebel-Lizorkin spaces,  $B_{p,q}^s(\mathbb{R})$  and  $F_{p,q}^s(\mathbb{R})$ , in terms of expressions involving its Haar coefficients or suitable variations thereof. The novelty in the current paper is that we obtain results for a range of the parameters  $(s, p, q)$  in which the Haar system is not an unconditional basis of the above spaces (see Figures 1 and 2 below); this complements earlier work of the authors [10, 11, 12, 13, 21] where a complete description was given for the parameter range in which the unconditional or Schauder basis property holds in each such space.

---

*Date:* September 7, 2022.

*2010 Mathematics Subject Classification.* 46E35, 46B15, 42C40, 42B35.

*Key words and phrases.* Haar system, Haar frames, Sobolev spaces, Besov and Triebel-Lizorkin spaces, dyadic versions of function spaces, wavelets, splines.

G.G. was supported in part by grants MTM2017-83262-C2-2-P and PID2019-105599GB-I00 from Ministerio de Ciencia e Innovación (Spain), and grant 20906/PI/18 from Fundación Séneca (Región de Murcia, Spain). A.S. was supported in part by National Science Foundation grant DMS 2054220. T.U. was supported in part by Deutsche Forschungsgemeinschaft (DFG), grant 403/2-1.

We denote the (inhomogeneous) Haar system in  $\mathbb{R}$  by

$$(1.1) \quad \mathcal{H} = \{h_{j,\mu} : j \geq -1, \mu \in \mathbb{Z}\},$$

where we let  $h(x) = \mathbb{1}_{[0, \frac{1}{2})}(x) - \mathbb{1}_{[\frac{1}{2}, 1)}(x)$  and

$$(1.2) \quad h_{j,\mu}(x) := h(2^j x - \mu), \quad \text{if } \mu \in \mathbb{Z}, j = 0, 1, 2, \dots$$

Note that  $h_{j,\mu}$  is supported in the closure of the dyadic interval

$$I_{j,\mu} = [2^{-j}\mu, 2^{-j}(\mu + 1)).$$

In the case  $j = -1$ , we just let

$$h_{-1,\mu} := \mathbb{1}_{I_{-1,\mu}} = \mathbb{1}_{[\mu, \mu+1)}, \quad \mu \in \mathbb{Z}.$$

Let  $F_{p,q}^s(\mathbb{R})$  and  $B_{p,q}^s(\mathbb{R})$  denote the usual Triebel-Lizorkin and Besov spaces [25]. It has been shown in [28, 21, 22] that  $\mathcal{H}$  is an unconditional basis of  $F_{p,q}^s(\mathbb{R})$  if and only if  $s$  belongs to the range

$$(1.3) \quad \max \left\{ 1/p - 1, 1/q - 1 \right\} < s < \min \left\{ 1/p, 1/q, 1 \right\};$$

moreover in the range (1.3) we also have the Haar characterization

$$(1.4) \quad \|f\|_{F_{p,q}^s} \approx \left\| \left( \sum_{j=-1}^{\infty} 2^{jsq} \left[ \sum_{\mu \in \mathbb{Z}} 2^j |\langle f, h_{j,\mu} \rangle| \mathbb{1}_{I_{j,\mu}} \right]^q \right)^{1/q} \right\|_p$$

for all  $f \in \mathcal{B}$ .




FIGURE 1. Parameter domain for  $\mathcal{H}$  to be an unconditional basis (left figure) or a Schauder basis (right figure) in  $F_{p,q}^s(\mathbb{R})$ .

It was also shown in [11] that  $\mathcal{H}$  is a Schauder basis of  $F_{p,q}^s(\mathbb{R})$  (with respect to natural enumerations) in the larger range

$$(1.5) \quad 1/p - 1 < s < \min \left\{ 1/p, 1 \right\}, \quad (\text{for all } 0 < q < \infty).$$

At the endpoints, the Schauder basis property holds for  $F_{p,q}^s$  if and only if

$$(1.6) \quad s = 1/p - 1 \quad \text{and} \quad 1/2 < p \leq 1$$

also for all  $0 < q < \infty$ ; see [12]. These regions are depicted in Figure 1.

For the spaces  $B_{p,q}^s(\mathbb{R})$  there is no such distinction, and  $\mathcal{H}$  is an unconditional basis under (1.5) (and also a Schauder basis under (1.6), if  $p = q$ ); see [13]. Moreover in the range (1.5) we have the Haar characterization

$$(1.7) \quad \|f\|_{B_{p,q}^s} \approx \left( \sum_{j=-1}^{\infty} 2^{j(s-\frac{1}{p})q} \left( \sum_{\mu \in \mathbb{Z}} |2^j \langle f, h_{j,\mu} \rangle|^p \right)^{q/p} \right)^{1/q}$$

for all  $f \in \mathcal{B}$ .

1.1. *The oversampled Haar systems - Haar frames.* A main feature of this paper is to show that the above characterizations in terms of Haar coefficients can be extended to larger regions depicted in Figure 2 below if we doubly oversample with Haar type coefficients obtained by a shift.

More concretely, we now define

$$(1.8) \quad \tilde{h}_{j,\nu}(x) := h(2^j x - \frac{\nu}{2}) \quad \text{if } j = 0, 1, 2, \dots \text{ and } \nu \in \mathbb{Z}.$$

Observe that for even  $\nu = 2\mu$  we recover the original Haar functions,  $\tilde{h}_{j,2\mu} = h_{j,\mu}$  supported in  $I_{j,\mu}$  but for odd  $\nu$  we obtain a *shifted Haar function*

$$\tilde{h}_{j,2\mu+1} = h_{j,\mu}(\cdot - 2^{-j-1}),$$

which is supported in the interval  $[2^{-j}(\mu+1/2), 2^{-j}(\mu+3/2)) = I_{j,\mu} + 2^{-j-1}$ . As before, for  $j = -1$  we just let

$$\tilde{h}_{-1,\nu} := h_{-1,\nu} = \mathbb{1}_{[\nu,\nu+1)}.$$

Then the *extended Haar system* is defined by

$$(1.9) \quad \mathcal{H}^{\text{ext}} = \left\{ \tilde{h}_{j,\nu} : j \geq -1, \nu \in \mathbb{Z} \right\}.$$

In what follows we will need to work with appropriate spaces of functions and distributions on which the (generalized) Haar coefficients are well defined. For a bounded interval define the linear functional (distribution)  $\lambda_I$  by

$$(1.10) \quad \lambda_I(f) = \int_I f(x) dx.$$

Obviously  $\lambda_I$  extends to  $L_1^{\text{loc}}$  and we have the trivial inequality

$$(1.11) \quad |\lambda_I(f)| \leq \int_I |f(x)| dx.$$

To deal with distributions associated with certain negative smoothness parameters we choose as a reference space  $\mathcal{B}$  of distributions the space

$$(1.12) \quad \mathcal{B} := B_{\infty,1}^{-1}(\mathbb{R}).$$

By standard embeddings, see e.g. [25, 2.7.1], we have

$$(1.13) \quad B_{p,q}^s(\mathbb{R}) \hookrightarrow \mathcal{B}, \quad \text{if } s > \frac{1}{p} - 1, \text{ or } s = \frac{1}{p} - 1 \text{ and } 0 < q \leq 1,$$

and

$$(1.14) \quad F_{p,q}^s(\mathbb{R}) \hookrightarrow \mathcal{B}, \quad \text{if } s > \frac{1}{p} - 1, \text{ or } s = \frac{1}{p} - 1 \text{ and } 0 < p \leq 1.$$

In particular, all the spaces that are used in Theorems 1.2-1.10 below are embedded into  $\mathcal{B}$ .

**Proposition 1.1.** *For a bounded interval  $I$  consider the distribution  $\lambda_I$  in (1.10). Then*

(i)  $\lambda_I$  extends to a bounded linear functional on  $\mathcal{B}$ , with operator norm  $O(1 + |I|)$ .

(ii) For every  $h \in \mathcal{H}^{\text{ext}}$  the linear functional  $f \mapsto \langle f, h \rangle$  is bounded on  $\mathcal{B}$  with uniformly bounded operator norm.

(iii) If  $f \in \mathcal{B}$  and  $\langle f, h \rangle = 0$ , for all  $h \in \mathcal{H}$ , then  $f = 0$ .

*Remark.* Clearly, using (1.11) one can also replace in (i) the space  $\mathcal{B}$  with  $\mathcal{B} + L_1^{\text{loc}}$ . We note that  $L_1$  is not embedded in  $\mathcal{B}$ , see the proof of Proposition 11.3 below.

In the rest of the paper, when  $f \in \mathcal{B}$ , we use the following notation, combining the standard Haar coefficients with the coefficients obtained from the shifted Haar functions:

$$(1.15) \quad \mathbf{c}_{j,\mu}(f) = 2^j |\langle f, \tilde{h}_{j,2\mu} \rangle| + 2^j |\langle f, \tilde{h}_{j,2\mu+1} \rangle|,$$

when  $j = 0, 1, \dots$ , and

$$\mathbf{c}_{-1,\mu}(f) = \langle f, h_{-1,\mu} \rangle = \langle f, \mathbb{1}_{[\mu,\mu+1)} \rangle.$$

Our first main result provides a characterization where in (1.7) the Haar coefficients  $2^j \langle f, h_{j,\mu} \rangle$  are replaced with the  $\mathbf{c}_{j,\mu}(f)$ . This covers as well the quasi-Banach range of parameters; see Figure 2 below.

**Theorem 1.2.** *Let  $1/2 < p < \infty$ ,  $1/2 < q \leq \infty$  and*

$$(1.16) \quad \max\{1/p - 1, 1/q - 1\} < s < 1.$$

*Then  $F_{p,q}^s(\mathbb{R})$  is the collection of all  $f \in \mathcal{B}$  such that*

$$(1.17) \quad \left\| \left( \sum_{j=-1}^{\infty} 2^{jsq} \left| \sum_{\mu \in \mathbb{Z}} \mathbf{c}_{j,\mu}(f) \mathbb{1}_{I_{j,\mu}} \right|^q \right)^{1/q} \right\|_p < \infty.$$

*Moreover, the latter quantity represents an equivalent quasi-norm in  $F_{p,q}^s(\mathbb{R})$ .*

Using terminology introduced by Gröchenig [15], one may say that  $\mathcal{H}^{\text{ext}}$  is a (quasi-)Banach frame<sup>1</sup> for  $F_{p,q}^s(\mathbb{R})$ . In signal processing language, this can be interpreted by saying that one may stably recover  $f$  from the sampled information  $\{\langle f, h \rangle : h \in \mathcal{H}^{\text{ext}}\}$ .

<sup>1</sup>In a Hilbert space  $H$ , a frame is a system of vectors  $\{e_j\} \subset H$ , which for some constants  $A, B > 0$  satisfies  $A\|f\|_H^2 \leq \sum_j |\langle f, e_j \rangle|^2 \leq B\|f\|_H^2$ ,  $\forall f \in H$ .

We remark that the condition  $s > \frac{1}{q} - 1$  in (1.16) is necessary, in view of the examples in [21]; see Remark 4.3 below. The analogous characterization for Besov spaces is valid in a larger range:

**Theorem 1.3.** *Let  $1/2 < p \leq \infty$ ,  $0 < q \leq \infty$  and*

$$1/p - 1 < s < 1.$$

*Then  $B_{p,q}^s(\mathbb{R})$  is the collection of all functions  $f \in \mathcal{B}$  such that*

$$\left( \sum_{j=-1}^{\infty} 2^{j(s-1/p)q} \left( \sum_{\mu \in \mathbb{Z}} |\mathfrak{c}_{j,\mu}(f)|^p \right)^{q/p} \right)^{1/q} < \infty.$$

*Moreover, the latter quantity represents an equivalent quasi-norm in  $B_{p,q}^s(\mathbb{R})$ .*




FIGURE 2. Parameter domain for  $\mathcal{H}^{\text{ext}}$  to be a characterizing frame for  $F_{p,q}^s(\mathbb{R})$  (left figure, Theorem 1.2) and  $B_{p,q}^s(\mathbb{R})$  (right figure, Theorem 1.3).

Figure 2 shows the regions of parameters where  $\mathcal{H}^{\text{ext}}$  is a characterizing frame for each of the spaces  $F_{p,q}^s(\mathbb{R})$  and  $B_{p,q}^s(\mathbb{R})$ .

We remark that a related result, in the special case of the Hölder spaces  $C^\alpha = B_{\infty,\infty}^\alpha(\mathbb{R})$ ,  $\alpha \in (0, 1)$ , and using a  $1/3$ -shifted Haar frame, has been recently obtained by Jaffard and Krim; see [16, Thm.1].

**1.2. Characterization of  $W_p^1(\mathbb{R})$  via Haar frames.** We now let  $s = 1$ , and consider in the Banach range  $1 \leq p \leq \infty$  the Sobolev space  $W_p^1(\mathbb{R})$ , endowed with the usual norm

$$\|f\|_{W_p^1(\mathbb{R})} = \|f\|_p + \|f'\|_p.$$

We also let  $BV(\mathbb{R})$  be the subspace of  $L_1(\mathbb{R})$  for which the distributional derivative belongs to the space  $\mathcal{M}$  of bounded Borel measures (with the norm given by the total variation of the measure) and define

$$\|f\|_{BV(\mathbb{R})} = \|f\|_1 + \|f'\|_{\mathcal{M}}.$$

Note that by our definition  $BV \subset L_1$  which deviates from the definition in some other places in the literature. We have the following result, that provides characterizations in terms of the oversampled Haar system  $\mathcal{H}^{\text{ext}}$ .

**Theorem 1.4.** *For all  $f \in \mathcal{B}$  the following hold.*

(i) *If  $1 < p \leq \infty$  then*

$$\|f\|_{W_p^1} \approx \sup_{j \geq -1} 2^{j(1-1/p)} \left( \sum_{\mu \in \mathbb{Z}} |\mathbf{c}_{j,\mu}(f)|^p \right)^{1/p}.$$

(ii) *In the case  $p = 1$  we have instead*

$$\|f\|_{BV} \approx \sup_{j \geq -1} \sum_{\mu \in \mathbb{Z}} |\mathbf{c}_{j,\mu}(f)|.$$

Clearly part (ii) implies the inequality

$$(1.18) \quad \sup_{j \geq -1} \sum_{\mu \in \mathbb{Z}} |\mathbf{c}_{j,\mu}(f)| \lesssim \|f\|_{W_1^1}$$

for all  $f \in \mathcal{B}$ . However the converse of this inequality fails as one checks by testing it with  $f = \mathbb{1}_{[0,1]} \in BV \setminus W_1^1$ ; we have  $\sup_j \sum_{\mu} |\mathbf{c}_{j,\mu}(\mathbb{1}_{[0,1]})| < \infty$ .

The fact that the Sobolev  $W_p^1(\mathbb{R})$  norm can be expressed in terms of a discrete norm of  $b_{p,\infty}^1$  type may seem surprising at first, but actually results of this sort can be found in the literature since the 60s, see [1]. The theorem is also reminiscent of characterizations via the uniform bounds for difference quotients  $h^{-1}(f(\cdot + h) - f)$ , see [24, Prop V.3] and more recently [5, 4].

**1.3. Dyadic Besov spaces.** In this section we present stronger results involving the standard Haar system  $\mathcal{H}$ , and suitable dyadic variants  $B_{p,q}^{s,\text{dyad}}$  of the Besov spaces.

We first recall the definition of the sequence spaces  $b_{p,q}^s$  and  $f_{p,q}^s$ ; see [9]. If  $s \in \mathbb{R}$  and  $0 < p, q \leq \infty$ , we define, for  $\beta = \{\beta_{j,\mu}\}_{j \geq -1, \mu \in \mathbb{Z}}$ ,

$$(1.19) \quad \|\beta\|_{b_{p,q}^s} := \left( \sum_{j=-1}^{\infty} \left[ 2^{j(s-1/p)} \left( \sum_{\mu \in \mathbb{Z}} |\beta_{j,\mu}|^p \right)^{1/p} \right]^q \right)^{1/q},$$

and if  $p < \infty$  we let

$$(1.20) \quad \|\beta\|_{f_{p,q}^s} := \left\| \left( \sum_{j=-1}^{\infty} \left| 2^{js} \sum_{\mu \in \mathbb{Z}} \beta_{j,\mu} \mathbb{1}_{I_{j,\mu}}(\cdot) \right|^q \right)^{1/q} \right\|_p.$$

These expressions have the obvious interpretations if  $\max\{p, q\} = \infty$ .

We additionally define for every  $f \in \mathcal{B}$  the quantity

$$\|f\|_{B_{p,q}^{s,\text{dyad}}} := \left\| \{2^j \langle f, h_{j,\mu} \rangle\}_{j,\mu} \right\|_{b_{p,q}^s}$$

and the vector spaces

$$B_{p,q}^{s,\text{dyad}}(\mathbb{R}) = \left\{ f \in \mathcal{B} : \|f\|_{B_{p,q}^{s,\text{dyad}}} < \infty \right\}.$$

Observe that  $\text{span } \mathcal{H} \subset B_{p,q}^{s,\text{dyad}}$ , so the spaces are not null. Also, the quantity  $\|f\|_{B_{p,q}^{s,\text{dyad}}}$  is a quasi-norm (not just a semi-norm), by Proposition 1.1. Since  $2^j |\langle f, h_{j,\mu} \rangle| \leq \mathfrak{c}_{j,\mu}(f)$  we note the following immediate consequence of Theorem 1.4.

**Corollary 1.5.** *If  $1 \leq p \leq \infty$  then*

$$(1.21) \quad W_p^1 \hookrightarrow B_{p,\infty}^{1,\text{dyad}}.$$

To avoid pathological cases, below we shall typically consider the range

$$(1.22) \quad \frac{1}{p} - 1 < s < 1,$$

and some end-point cases of these. Recall that in the smaller range (1.5) we have  $B_{p,q}^s = B_{p,q}^{s,\text{dyad}}$ , cf. (1.7). We remark that when  $s > 1$  (or  $s = 1$  and  $0 < q < \infty$ ), the spaces  $B_{p,q}^{1,\text{dyad}}$  contain no nontrivial  $C^1$  functions (see Proposition 11.1) while for  $s < 1/p - 1$  the spaces are not complete (see Proposition 11.2.)

Assume now that (1.22) holds. By Theorem 1.3 we have  $B_{p,q}^s \hookrightarrow B_{p,q}^{s,\text{dyad}}$ , and the inclusion is proper provided that

$$(1.23) \quad 1/p < s < 1, \quad \text{or } s = 1/p \quad \text{and} \quad q < \infty$$

(since in that range Haar functions do not belong to  $B_{p,q}^s$ ). Our goal is to prove converse inequalities of the form

$$(1.24) \quad \|f\|_{B_{p,q}^s} \lesssim \|f\|_{B_{p,q}^{s,\text{dyad}}}, \quad \text{provided that } f \in B_{p,q}^s(\mathbb{R}).$$

Such inequalities will imply that  $\|\cdot\|_{B_{p,q}^{s,\text{dyad}}}$  is an equivalent norm in  $B_{p,q}^s$ , a result which may seem surprising outside the usual unconditionality region. Our first result in this direction is the following.

**Theorem 1.6.** *Let  $1 < p \leq \infty$ ,  $0 < q \leq \infty$ , and  $1/p < s < 1$ . Then (1.24) holds. In particular,  $B_{p,q}^s$  is a proper closed subspace of  $B_{p,q}^{s,\text{dyad}}$ , and we have*

$$(1.25) \quad \|f\|_{B_{p,q}^s} \approx \|f\|_{B_{p,q}^{s,\text{dyad}}}, \quad \text{for all } f \in B_{p,q}^s(\mathbb{R}).$$

There are also some precedent results of this nature in the literature. When  $p = q = \infty$ , a norm equivalence as in (1.25) (for continuous functions in the interval  $[0,1]$ ) was proved by Golubov [14, Corollary 6]; see also [18, Corollary 3.2], [19, Theorem 7.c.3] and references therein.

1.4. *Inclusions for the limiting case  $s = 1$ .* In what follows the notation  $\mathfrak{X}_1 \hookrightarrow \mathfrak{X}_2$  will indicate a continuous embedding of the space  $\mathfrak{X}_1$  in the space  $\mathfrak{X}_2$ . As already remarked above we may focus on the cases  $s < 1$  or  $s = 1$ ,  $q = \infty$ , cf. Proposition 11.1.

We now state inclusions into the spaces  $B_{p,q}^{s,\text{dyad}}$ , in the case that  $s = 1$  and  $q = \infty$ . Note that in one of the inclusions we use the smaller space

$$F_{p,\infty}^{1,\text{dyad}} := \left\{ f \in \mathcal{B} : \left\{ 2^j \langle f, h_{j,\mu} \rangle \right\}_{\substack{j \geq -1 \\ \mu \in \mathbb{Z}}} \in f_{p,\infty}^1 \right\}.$$

**Theorem 1.7.** *Let  $1/2 \leq p \leq \infty$ . Then the following hold.*

(i) *If  $1/2 \leq p < \infty$  then*

$$(1.26) \quad B_{p,q}^1 \hookrightarrow B_{p,\infty}^{1,\text{dyad}} \iff q \leq \min\{p, 2\},$$

$$(1.27) \quad F_{p,q}^1 \hookrightarrow B_{p,\infty}^{1,\text{dyad}} \iff q \leq 2.$$

*For  $p = \infty$  we have*

$$(1.28) \quad B_{\infty,q}^1 \hookrightarrow B_{\infty,\infty}^{1,\text{dyad}} \iff q \leq 1.$$

(ii) *For  $1/2 < p < \infty$*

$$(1.29) \quad F_{p,2}^1 \hookrightarrow F_{p,\infty}^{1,\text{dyad}}.$$

The next result is a converse inequality to (1.21), which in particular implies that

$$\|f\|_{B_{p,\infty}^{1,\text{dyad}}} = \sup_{j \geq -1} 2^{j(1-1/p)} \left( \sum_{\mu \in \mathbb{Z}} |2^j \langle f, h_{j,\mu} \rangle|^p \right)^{1/p}$$

is an equivalent norm in  $W_p^1(\mathbb{R})$ . Lower bounds of this type, for absolutely continuous functions in the interval  $[0, 1]$ , can be found in the work of Bočkarev [1, Theorem 7], [2, Theorem I.3.4], or [19, Corollary 7.b.2] and references therein. Below we establish, by different methods, the following result, which is complementary to Theorem 1.4.

**Theorem 1.8.** *Let  $1 < p \leq \infty$ . Then*

$$(1.30) \quad \|f\|_{W_p^1} \lesssim \|f\|_{B_{p,\infty}^{1,\text{dyad}}}, \quad \text{provided } f \in W_p^1(\mathbb{R}).$$

*In particular,  $W_p^1(\mathbb{R})$  is a proper closed subspace of  $B_{p,\infty}^{1,\text{dyad}}$ , and it holds*

$$(1.31) \quad \|f\|_{W_p^1} \approx \|f\|_{B_{p,\infty}^{1,\text{dyad}}} \approx \|f\|_{F_{p,\infty}^{1,\text{dyad}}}, \quad \text{for } f \in W_p^1(\mathbb{R}), \quad 1 < p < \infty.$$

*Remark.* The inequality in (1.30) (and hence, the first equivalence in (1.31)) is also true when  $p = 1$ , due to the result of Bočkarev [1]. The proof we give here, however, is only valid for  $p > 1$ .

1.5. *Inclusions for the limiting case  $s = 1/p - 1$ .* We state inclusions into the spaces  $B_{p,q}^{s,\text{dyad}}$ , in the case that  $s = 1/p - 1$  and  $q = \infty$ .

**Theorem 1.9.** (i) *For  $0 < p, u \leq \infty$  the embedding  $B_{p,u}^{1/p-1} \hookrightarrow B_{p,q}^{1/p-1,\text{dyad}}$  can only hold when  $q = \infty$ .*

(ii) *If  $p \geq 1/2$  then*

$$(1.32) \quad B_{p,q}^{1/p-1} \hookrightarrow B_{p,\infty}^{1/p-1,\text{dyad}} \iff q \leq \min\{1, p\}$$

(iii) *If  $1/2 < p \leq 1$  then*

$$(1.33) \quad F_{p,\infty}^{1/p-1} \hookrightarrow B_{p,\infty}^{1/p-1,\text{dyad}}.$$



*Remark.* In case  $p = 1$  we have for all  $f \in L_1$  the straightforward inequality

$$\sup_{j \geq -1} \sum_{\mu \in \mathbb{Z}} |\langle f, h_{j,\mu} \rangle| \lesssim \|f\|_1;$$

in particular  $L_1 \cap \mathcal{B} \subset B_{1,\infty}^{0,\text{dyad}}$ .

1.6. *The case  $s = 1/p$ .* When  $1 < p < \infty$ , the unconditionality of the Haar basis implies that

$$B_{p,q}^s(\mathbb{R}) = B_{p,q}^{s,\text{dyad}}(\mathbb{R}), \quad \frac{1}{p} - 1 < s < \frac{1}{p}.$$

On the other hand, Theorem 1.6 implies the norm equivalence

$$\|f\|_{B_{p,q}^s} \approx \|f\|_{B_{p,q}^{s,\text{dyad}}}, \quad f \in B_{p,q}^s, \quad \frac{1}{p} < s < 1.$$

These two results might suggest that the norm equivalence could hold also at the dividing line  $s = 1/p$ . Here we show that this is not the case, at least when  $q = \infty$ .

**Theorem 1.10.** *Let  $1 \leq p < \infty$ . Then*

(i) *there exists a sequence  $\{f_N\}_{N=1}^\infty$  of functions in  $B_{p,\infty}^{1/p}$  such that*

$$\|f_N\|_{B_{p,\infty}^{1/p,\text{dyad}}} = 1 \text{ and } \|f_N\|_{B_{p,\infty}^{1/p}} \gtrsim N,$$

(ii)  $B_{p,\infty}^{1/p,\text{dyad}} \setminus B_{p,\infty}^{1/p} \neq \emptyset$ .

*Remarks.* (i) The following comments on the case  $s = 1/p$  as a dividing line between Theorems 1.3 and 1.6 on which both theorems fail. If  $1 < p < \infty$ , the embedding  $B_{p,\infty}^{1/p}(\mathbb{R})$  into  $B_{p,\infty}^{1/p,\text{dyad}}(\mathbb{R})$  which is established by Theorem 1.3 is proper, i.e.,  $B_{p,\infty}^{1/p}(\mathbb{R}) \subsetneq B_{p,\infty}^{1/p,\text{dyad}}(\mathbb{R})$ . Moreover, on the smaller space  $B_{p,\infty}^{1/p}(\mathbb{R})$ , the norms are not equivalent, that is,

$$(1.34) \quad \sup \{ \|f\|_{B_{p,\infty}^{1/p}} : f \in B_{p,\infty}^{1/p} \text{ and } \|f\|_{B_{p,\infty}^{1/p,\text{dyad}}} = 1 \} = \infty.$$

Both statements are an immediate consequence of Theorem 1.10.

(ii) Observe that  $BV(\mathbb{R}) \hookrightarrow B_{1,\infty}^1(\mathbb{R})$  and that for  $p = 1$  we have the embedding  $BV(\mathbb{R}) \hookrightarrow B_{1,\infty}^{1,\text{dyad}}(\mathbb{R})$  as an immediate consequence of Theorem 1.4. Theorem 1.10 shows that this embedding is also proper, i.e.  $B_{1,\infty}^{1,\text{dyad}}(\mathbb{R}) \setminus BV(\mathbb{R}) \neq \emptyset$ .

1.7. *Further directions.* We mention a few problems left open in this paper.

1.7.1. *Besov-type spaces.* Concerning (1.34) in Theorem 1.10, we do not know whether the inequality

$$(1.35) \quad \|f\|_{B_{p,\infty}^{1/p}} \leq C \|f\|_{B_{p,\infty}^{1/p,\text{dyad}}}$$

could be true for  $1 < p < \infty$  when restricted to  $f \in \mathcal{S}(\mathbb{R})$ . It is also open to determine whether such inequality could hold if the  $B_{p,\infty}^{1/p}$  norm is replaced by  $B_{p,q}^{1/p}$  with  $q < \infty$ .

1.7.2.  $F_{p,q}^s$  versus  $F_{p,q}^{s,\text{dyad}}$ . It would be interesting to establish an optimal analogue of Theorems 1.6/1.8 for Triebel-Lizorkin spaces.

1.7.3. *Wavelet frames.* The sharp results on the failure of unconditional convergence of Haar expansions in [21] (described above) have been extended by R. Srivastava [23] to classes of spline wavelets with more restrictive smoothness assumptions. It is then natural to investigate extensions of our results on Haar frames to suitable classes of oversampled systems of spline wavelets.

1.8. *Structure of the paper.* In §2 we introduce notation and review some preliminary known results on maximal characterizations of function spaces, and Chui-Wang wavelet bases that we need in the proofs of our results.

In §3 we clarify the role of the space  $\mathcal{B}$  and proof Proposition 1.1.

In §4 we consider the characterization of function spaces via Haar frames and give the proofs of Theorems 1.2 and 1.3 (as a combination of four Propositions 4.1, 4.4, 4.5 and 4.9).

In §5 we establish Haar frame characterizations of Sobolev and bounded variation spaces and give a proof of Theorem 1.4.

In §6 we prove the sufficiency of the conditions for the embeddings into  $B_{p,\infty}^{1,\text{dyad}}$  or  $F_{p,\infty}^{1,\text{dyad}}$  in Theorem 1.7 and the sufficiency for the conditions of embedding into  $B_{p,\infty}^{1/p-1,\text{dyad}}$  in Theorem 1.9.

In §7 we prove Theorems 1.6 and 1.8.

In §8 we prove necessary conditions for the embeddings into  $B_{p,\infty}^{1,\text{dyad}}$ . Specifically, in Theorem 1.7 the necessary conditions  $q \leq p$  in (1.26),  $q \leq 2$  in (1.27), (1.26), and  $q \leq 1$  in (1.28) correspond to Lemma 8.1, Lemma 8.2, and Lemma 8.3, respectively.

In §9 we obtain necessary conditions in part (ii) of Theorem 1.9 for the embeddings into  $B_{p,\infty}^{1/p-1,\text{dyad}}$ .

In §10 we prove Theorem 1.10.

In §11 we give a simple proof for the fact that  $C^1$  functions in  $B_{p,q}^{1,\text{dyad}}$  are constant (Proposition 11.1); moreover show that  $B_{p,q}^{s,\text{dyad}}$  is not complete when  $s < 1/p - 1$  (Proposition 11.2) and finally prove part (i) of Theorem 1.9 (see Proposition 11.3).

## 2. PRELIMINARIES ON FUNCTION SPACES AND WAVELET BASES

2.1. *Definition of spaces.* Let  $s \in \mathbb{R}$  and  $0 < p, q \leq \infty$  be given. We shall use both definitions and characterizations of  $B_{p,q}^s$  and  $F_{p,q}^s$  in terms of dyadic frequency decompositions and in terms of sequences of compactly supported kernels with cancellation (see e.g. [26, 2.5.3, 2.4.6] or [27, §1.3,1.4] where the terminology local means is used).

Consider two functions  $\beta_0, \beta \in \mathcal{S}(\mathbb{R})$  such that  $|\widehat{\beta}_0(\xi)| > 0$  when  $|\xi| \leq 1$  and  $|\widehat{\beta}(\xi)| > 0$  when  $1/4 \leq |\xi| \leq 1$ . Assume further that  $\beta(\cdot)$  has vanishing moments up to a sufficiently large order  $M \in \mathbb{N}$ , that is,

$$(2.1) \quad \int_{\mathbb{R}} \beta(x) x^m dx = 0 \quad \text{when } m < M.$$

The precise value of  $M$  is not relevant, but for the properties used in the paper it will suffice with

$$(2.2) \quad M > 1/p + |s| + 2.$$

We let  $\beta_k(x) := 2^{kd}\beta(2^kx)$ ,  $k \geq 1$ , and define for  $k \in \mathbb{N}_0$  the convolution operators

$$L_k f := \beta_k * f,$$

acting on distributions  $f \in \mathcal{S}'(\mathbb{R})$ .

The Besov space  $B_{p,q}^s(\mathbb{R})$  is the set of all distributions  $f \in \mathcal{S}'(\mathbb{R})$  such that

$$(2.3) \quad \|f\|_{B_{p,q}^s} := \left( \sum_{k=0}^{\infty} \left( 2^{ks} \|L_k f\|_p \right)^q \right)^{1/q} < \infty.$$

If  $p < \infty$ , the Triebel-Lizorkin space  $F_{p,q}^s(\mathbb{R})$  is the set of all  $f \in \mathcal{S}'(\mathbb{R})$  such that

$$(2.4) \quad \|f\|_{F_{p,q}^s} := \left\| \left( \sum_{k=0}^{\infty} 2^{ksq} |L_k f(x)|^q \right)^{1/q} \right\|_p < \infty.$$

Different choices of  $\beta_0, \beta$  give rise to the same spaces and equivalent quasi-norms; see e.g. [27, Theorem 1.7]. From now on we will assume that

$$\text{supp } \beta_0 \subset (-1/2, 1/2) \quad \text{and} \quad \text{supp } \beta \subset (-1/2, 1/2).$$

We shall often use the following decomposition of distributions in  $\mathcal{S}'(\mathbb{R})$ . Let  $\eta_0 \in C_0^\infty(\mathbb{R})$  be supported on  $\{|\xi| < 3/4\}$  and such that  $\eta_0(\xi) = 1$  when  $|\xi| \leq 1/2$ . Define the convolution operators  $\Lambda_0$ , and  $\Lambda_k$  for  $k \geq 1$ , by

$$\begin{aligned} \widehat{\Lambda_0 f}(\xi) &= \frac{\eta_0(\xi)}{\widehat{\beta}_0(\xi)} \widehat{f}(\xi) \\ \widehat{\Lambda_k f}(\xi) &= \frac{\eta_0(2^{-k}\xi) - \eta_0(2^{-k+1}\xi)}{\widehat{\beta}(2^{-k}\xi)} \widehat{f}(\xi), \quad k \geq 1. \end{aligned}$$

Then, for all  $f \in \mathcal{S}'(\mathbb{R})$  we have

$$(2.5) \quad f = \sum_{k=0}^{\infty} L_k \Lambda_k f$$

with convergence in  $\mathcal{S}'(\mathbb{R})$ . Also, it holds

$$\left( \sum_{k=0}^{\infty} \left( 2^{ks} \|\Lambda_k f\|_p \right)^q \right)^{1/q} \lesssim \|f\|_{B_{p,q}^s},$$

and likewise for the  $F$ -norms.

**2.2. Maximal functions.** We follow Triebel [25, 26]. Given  $f \in L_1^{\text{loc}}(\mathbb{R})$ , consider the Hardy-Littlewood maximal function, defined by

$$(2.6) \quad Mf(x) := \sup_{x \in I} \frac{1}{|I|} \int_I |f(x)| dx,$$

where the sup is taken over all intervals  $I$  that contain  $x$ . A classical result of Fefferman and Stein asserts that, if  $1 < p < \infty$  and  $1 < q \leq \infty$  then

$$(2.7) \quad \left\| \left( \sum_j |Mf_j|^q \right)^{1/q} \right\|_p \lesssim \left\| \left( \sum_j |f_j|^q \right)^{1/q} \right\|_p.$$

for all sequences of measurable functions  $\{f_j\}$  with finite right hand side.

Let us further define the Peetre maximal functions [20]. Given  $j \in \mathbb{N}$  and  $A > 0$  we let

$$\mathfrak{M}_{j,A}^{**} f(x) = \sup_{h \in \mathbb{R}} \frac{|f(x+h)|}{(1+2^j|h|)^A}.$$

Let  $\mathcal{E}_j$  be the set of distributions  $f \in \mathcal{S}'(\mathbb{R})$  such that  $\text{supp } \hat{f}$  is supported in an interval of diameter  $\leq 2^{j+2}$ . Then for all  $f \in \mathcal{E}_j$  it holds

$$(2.8) \quad \mathfrak{M}_{j,A}^{**} f(x) \lesssim_{s,A} [M(|f|^s)(x)]^{1/s},$$

provided that  $s \geq 1/A$ ; see [20] or [25, Theorem 1.3.1]. In particular, if  $0 < p \leq \infty$  and  $A > 1/p$  then

$$(2.9) \quad \|\mathfrak{M}_{j,A}^{**} f\|_p \leq C_{p,A} \|f\|_p, \quad f \in \mathcal{E}_j.$$

Also, from (2.7) and (2.8), if  $0 < p < \infty$ ,  $0 < q \leq \infty$  and  $A > \max\{1/p, 1/q\}$ , then

$$(2.10) \quad \left\| \left( \sum_j |\mathfrak{M}_{j,A}^{**} f_j|^q \right)^{1/q} \right\|_p \leq C_{p,q,A} \left\| \left( \sum_j |f_j|^q \right)^{1/q} \right\|_p$$

for all sequences of functions  $(f_j)$  such that  $f_j \in \mathcal{E}_j$ .

Below we shall also use the (smaller) maximal functions

$$(2.11) \quad \mathfrak{M}_j f(x) = \sup_{|h| \leq 2^{-j}} |f(x+h)| \quad \text{and} \quad \mathfrak{M}_j^* f(x) = \sup_{|h| \leq 2^{-j+2}} |f(x+h)|.$$

Note that for all  $A > 0$  it holds

$$\mathfrak{M}_j f(x) \leq \mathfrak{M}_j^* f(x) \lesssim \mathfrak{M}_{j,A}^{**} f(x),$$

so in particular, for all  $0 < p \leq \infty$  we have

$$(2.12) \quad \|\mathfrak{M}_j^* f\|_p \lesssim \|f\|_p, \quad f \in \mathcal{E}_j.$$

We shall also make use of the following elementary inequality: if  $0 < p \leq \infty$  then

$$(2.13) \quad \|\mathfrak{M}_j^* f\|_p \lesssim 2^{\ell/p} \|\mathfrak{M}_{j+\ell} f\|_p, \quad \ell \geq 0.$$

To prove this assertion, if we let  $x_\nu = \nu 2^{-(j+\ell)}$ , then we have

$$\mathfrak{M}_j^* f(x) \leq \sup_{|\nu| \leq 2^{\ell+2}} |\mathfrak{M}_{j+\ell} f(x + x_\nu)| \leq \left( \sum_{|\nu| \leq 2^{\ell+2}} |\mathfrak{M}_{j+\ell} f(x + x_\nu)|^p \right)^{\frac{1}{p}}.$$

Then, taking  $L_p$  quasi-norms one easily obtains (2.13).

**2.3. Chui-Wang wavelets.** The proofs of Theorems 1.2 and 1.3 will require a characterization in terms of a wavelet basis generated by the *Chui-Wang polygon* and its dual.

Define the  $m$ -fold convolution  $\mathcal{N}_m$  of characteristic functions of  $[0, 1)$ , i.e.  $\mathcal{N}_1 = \mathbb{1}_{[0,1)}$ , and, for  $m \geq 2$ ,  $\mathcal{N}_m = \mathcal{N}_{m-1} * \mathbb{1}_{[0,1)}$ . In particular we get for  $m = 2$  the *hat function*

$$(2.14) \quad \mathcal{N}_2(x) = \begin{cases} x, & x \in [0, 1], \\ 2 - x, & x \in [1, 2], \\ 0, & x \in \mathbb{R} \setminus [0, 2]. \end{cases}$$

Let

$$\mathcal{N}_{2;j,\nu}(x) := \mathcal{N}_2(2^j x - \nu), \quad j \geq 0, \nu \in \mathbb{Z},$$

which is a hat function adapted to  $\text{supp } \mathcal{N}_{2;j,\nu} = [2^{-j}\nu, 2^{-j}(\nu+2)]$ .

The next elementary observation will be crucial in what follows.

**Lemma 2.1.** *If  $f$  is locally absolutely continuous in  $\mathbb{R}$ , then for all  $j \geq 1$  and  $\nu \in \mathbb{Z}$  it holds*

$$(2.15) \quad \langle f', \mathcal{N}_{2;j,\nu} \rangle = -2^j \langle f, \tilde{h}_{j-1,\nu} \rangle,$$

while for  $j = 0$  it holds

$$\langle f', \mathcal{N}_{2;0,\nu} \rangle = -\langle f, h_{-1,\nu} \rangle + \langle f, h_{-1,\nu+1} \rangle, \quad \nu \in \mathbb{Z}.$$

*Proof.* Integrating by parts one has

$$\langle f', \mathcal{N}_{2;j,\nu} \rangle = -\langle f, (\mathcal{N}_{2;j,\nu})' \rangle.$$

Now, a simple computation gives

$$\begin{aligned} (\mathcal{N}_{2;j,\nu})' &= 2^j \mathbb{1}_{(2^{-j}\nu, 2^{-j}(\nu+1))} - 2^j \mathbb{1}_{(2^{-j}(\nu+1), 2^{-j}(\nu+2))} \\ &= 2^j \mathbb{1}_{I_{j,\nu}} - 2^j \mathbb{1}_{I_{j,\nu+1}} = 2^j \tilde{h}_{j-1,\nu}, \end{aligned}$$

where the last equality follows from the definition of the shifted Haar system in (1.8) (if  $j \geq 1$ ). Combining the two expressions one obtains (2.15). The case  $j = 0$  is similar.  $\square$

The Chui-Wang polygon [7, Theorem 1], [6, 6.2.5, 6.2.6] is the compactly supported wavelet given by

$$\begin{aligned}
 \psi(x) &= \frac{1}{2} \sum_{\ell=0}^2 (-1)^\ell \mathcal{N}_4(\ell+1) \mathcal{N}_4''(2x-\ell), \\
 (2.16) \quad &= \frac{1}{2} \sum_{\ell=0}^2 (-1)^\ell \mathcal{N}_4(\ell+1) \sum_{j=0}^2 (-1)^j \binom{2}{j} \mathcal{N}_2(2x-j-\ell) \\
 &= \sum_{k \in \mathbb{Z}} b_k \cdot \mathcal{N}_2(2x-k)
 \end{aligned}$$

with a finite sequence  $(b_k)$ . The wavelet  $\psi$  is compactly supported and has two vanishing moments, i.e.,  $\int \psi(x) dx = \int x \psi(x) dx = 0$ . For  $j \in \mathbb{N}_0$  and  $\mu \in \mathbb{Z}$  let

$$\psi_{j,\mu}(x) = \psi(2^j x - \mu),$$

while for  $j = -1$  we let

$$\psi_{-1,\mu}(x) = \mathcal{N}_{2;0,\mu}(x) = \mathcal{N}_2(x - \mu).$$

Then we have the orthogonality relations with respect to different scales

$$\langle \psi_{j,\mu}, \psi_{j',\mu'} \rangle = 0 \quad , \quad j \neq j'.$$

In contrast to that it only forms a Riesz basis within one and the same scale with respect to different translations. The dual basis can be computed precisely [8] and does not provide compact support. However, the coefficients




FIGURE 3. The Chui-Wang wavelet  $\psi_2$  of order 2 and its dual  $\psi_2^*$

in

$$(2.17) \quad \psi^*(x) = \sum_{k \in \mathbb{Z}} a_k \psi(x - k)$$

are exponentially decaying; see the paper [8] for explicit formulas for  $a_k$ . Observe from (2.17) that also  $\psi^*$  has two vanishing moments.

Using this construction, Derevianko and Ullrich provided the following characterization for the  $F_{p,q}^s$  and  $B_{p,q}^s$  spaces; see [8, Theorem 5.1].

**Theorem 2.2.** [8] *Let  $0 < p \leq \infty$ ,  $0 < q \leq \infty$ .*

(i) *If  $p < \infty$  and*

$$(2.18) \quad \max\{1/p, 1/q\} - 2 < r < 1 + \min\{1/p, 1/q, 1\}$$

*then we have for all  $f \in B_{\infty,1}^{-2}(\mathbb{R})$*

$$(2.19) \quad \|f\|_{F_{p,q}^r} \approx \left\| \left( \sum_{j=-1}^{\infty} 2^{jr q} \left| \sum_{\mu \in \mathbb{Z}} 2^j \langle f, \psi_{j,\mu} \rangle \mathbb{1}_{I_{j,\mu}} \right|^q \right)^{1/q} \right\|_p.$$

(ii) *If*

$$(2.20) \quad 1/p - 2 < r < \max\{1 + 1/p, 2\}$$

*then we have for all  $f \in B_{\infty,1}^{-2}(\mathbb{R})$*

$$(2.21) \quad \|f\|_{B_{p,q}^r} \approx \left( \sum_{j=-1}^{\infty} 2^{j(r-1/p)q} \left[ \sum_{\mu \in \mathbb{Z}} |2^j \langle f, \psi_{j,\mu} \rangle|^p \right]^{q/p} \right)^{1/q}.$$

*Remark 2.3.* Concerning part (i), we remark that the result stated in [8, Theorem 5.1], requires the additional restriction  $r < 1$ , which comes from a similar restriction in [8, Proposition 5.4]. This restriction, however, can be lifted and replaced by  $r < 1 + \max\{1/p, 1/q\}$ , using a complex interpolation argument which involves part (ii) (case  $p = q$ ), as we discuss in Step 3 of Proposition 4.5 below.

### 3. HAAR FUNCTIONS AS LINEAR FUNCTIONALS ON $\mathcal{B}$ : PROOF OF PROPOSITION 1.1

3.1. *Proof of (i) and (ii).* Since every  $h \in \mathcal{H}$  is a difference of two characteristic functions of intervals of length  $\leq 1$  part (ii) is an immediate consequence of part (i). It suffices to analyze  $\lambda_I$  on  $\mathcal{B}$ , for each bounded interval  $I$ . Let  $f \in \mathcal{B} = B_{\infty,1}^{-1}(\mathbb{R})$ . Using the decomposition in (2.5) we can write  $f = \sum_{k=0}^{\infty} L_k f_k$  where the Fourier transform  $\widehat{f_k}$  is supported in  $\{\xi : |\xi| \leq 2^k\}$  and the  $f_k$  satisfy

$$(3.1) \quad \sum_{k \geq 0} 2^{-k} \|f_k\|_{\infty} \lesssim \|f\|_{\mathcal{B}};$$

here,  $L_k f = \beta_k * f$  where  $\beta_0, \beta$  are even functions in  $C_c^{\infty}(-1/2, 1/2)$ , and  $\beta_k = 2^k \beta(2^k \cdot)$ , for  $k \geq 1$ . Also,  $\int \beta(x) dx = 0$ .

Note that

$$(3.2) \quad \|L_0 \mathbb{1}_I\|_1 \lesssim |I|.$$

By (3.1) one needs to show that  $\|L_k \mathbb{1}_I\|_1 \lesssim 2^{-k}$  for  $k \geq 1$  but one actually gets the better estimate

$$(3.3) \quad \|L_k \mathbb{1}_I\|_1 \lesssim \min\{|I|, 2^{-k}\}, \quad k \geq 1.$$

Hence  $|\langle f, \mathbb{1}_I \rangle| \lesssim \max\{|I|, 1\} \sum_{k \geq 0} 2^{-k} \|f_k\|_{\infty}$  and we deduce that  $\lambda_I$  extends to a bounded linear functional on  $\mathcal{B}$ , with  $\|\lambda_I\|_{\mathcal{B}^*} \lesssim \max\{1, |I|\}$ .

It remains to verify (3.3). Fix  $I$ , with center  $y_I$ , and assume first that  $2^{-k} > |I|$ . Then the function  $L_k \mathbb{1}_I = \beta_k * \mathbb{1}_I$  is supported in an interval centered at  $y_I$  with length  $O(2^{-k})$  and satisfies  $|\beta_k * \mathbb{1}_I(x)| \leq \|\beta_k\|_\infty \|\mathbb{1}_I\|_1 \lesssim 2^k |I|$ . Thus we obtain  $\|\beta_k * \mathbb{1}_I\|_1 \lesssim |I|$  which is in (3.3) in this case.

Now assume  $2^{-k} \leq |I|$ . Let  $y_+$  and  $y_-$  be the endpoints of  $I$ . Let  $\mathcal{U}_k$  be the union of the two closed intervals of length  $2^{-k+2}$  centered at  $y_+$  and  $y_-$ . Then  $\beta_k * \mathbb{1}_I$  is supported in  $\mathcal{U}_k$ , which has size  $|\mathcal{U}_k| = O(2^{-k})$ . This assertion, combined with  $\|\beta_k * \mathbb{1}_I\|_\infty \leq \|\beta_k\|_1 = O(1)$ , also implies (3.3) in this case.

3.2. *Proof of (iii).* For the argument below we shall use the dyadic averaging operators, defined for  $N \geq 0$  by

$$(3.4) \quad \mathbb{E}_N f(x) := \sum_{\mu \in \mathbb{Z}} 2^N \langle f, \mathbb{1}_{I_{N,\mu}} \rangle \mathbb{1}_{I_{N,\mu}}(x).$$

In view of (i), these operators can be defined acting on distributions  $f \in \mathcal{B}$  such that  $E_N : \mathcal{B} \rightarrow L_\infty$  has operator norm  $O(2^N)$ .

Let now  $f \in \mathcal{B}$  such that  $\langle f, h \rangle = 0$ , for all  $h \in \mathcal{H}$ . Since each  $I_{N,\mu}$  belongs to  $\text{span } \mathcal{H}$ , this implies that  $\mathbb{E}_N f = 0$ , for all  $N \geq 0$ . We then must show that  $f = 0$ , which is a direct consequence of part (b) in the following lemma.

**Lemma 3.1.** (a) *The operators  $\mathbb{E}_N$  satisfy the uniform bound*

$$\sup_{N \geq 0} \|\mathbb{E}_N\|_{\mathcal{B} \rightarrow B_{\infty,\infty}^{-1}} < \infty.$$

(b) *If  $f \in \mathcal{B}$  then  $\|\mathbb{E}_N f - f\|_{B_{\infty,\infty}^{-1}} \rightarrow 0$  as  $N \rightarrow \infty$ .*

*Proof.* Part (a) is implicit in [13]. Indeed, it follows by combining the estimates stated in the four propositions in [13, §4], for the cases  $s = -1$  and  $p = \infty$ .

We now show part (b). Let  $f \in \mathcal{B}$  and write  $f = \sum_{k=0}^{\infty} L_k f_k$  as at the beginning of §3.1. The above series converges in  $\mathcal{S}'$  and also in the  $\mathcal{B}$ -norm. Then, given  $\varepsilon > 0$  one can find  $g = \sum_{k=0}^J L_k f_k$  such that

$$\|f - g\|_{\mathcal{B}} < \varepsilon.$$

Observe that  $g$  is bounded, since

$$\|g\|_\infty \leq \sum_{k=0}^J \|L_k f_k\|_\infty \lesssim \sum_{k=0}^J \|f_k\|_\infty \lesssim 2^J \|f\|_{\mathcal{B}}.$$

A similar reasoning shows that  $\|g'\|_\infty < \infty$ , so in particular  $g$  is uniformly continuous. Thus exists an integer  $N_0 \in \mathbb{N}$  such that

$$\|g - \mathbb{E}_N g\|_\infty < \varepsilon, \text{ for all } N \geq N_0.$$



Combining these assertions, and using the trivial embeddings

$$L_\infty \hookrightarrow \mathcal{B} \hookrightarrow B_{\infty,\infty}^{-1},$$

we obtain, for all  $N \geq N_0$ ,

$$\begin{aligned} \|f - \mathbb{E}_N f\|_{B_{\infty,\infty}^{-1}} &\leq \|f - g\|_{\mathcal{B}} + \|g - \mathbb{E}_N g\|_\infty + \|\mathbb{E}_N(g - f)\|_{B_{\infty,\infty}^{-1}} \\ &\lesssim 2\varepsilon + C \|g - f\|_{\mathcal{B}} \lesssim \varepsilon, \end{aligned}$$

where in the second inequality we have used part (a).  $\square$

#### 4. CHARACTERIZATIONS BY HAAR FRAMES: PROOFS OF THEOREMS 1.2 AND 1.3

The proofs of Theorems 1.2 and 1.3 will follow from the four Propositions 4.1, 4.4, 4.5 and 4.9 stated below.

The first proposition is a strengthening of [28, Proposition 2.8]. It gives one of the inclusions asserted in Theorems 1.2 and 1.3. The region of indices is the same as in Figure 2. We set  $\mathbf{c}(f) = \{\mathbf{c}_{j,\mu}(f)\}_{j \geq -1, \mu \in \mathbb{Z}}$  with  $\mathbf{c}_{j,\mu}(f)$  as in (1.15).

**Proposition 4.1.** *Let  $0 < p, q \leq \infty$  and  $s \in \mathbb{R}$ .*

(i) *If  $p < \infty$  and  $\max\{1/p - 1, 1/q - 1\} < s < 1$ , then for all  $f \in F_{p,q}^s$*

$$(4.1) \quad \|\mathbf{c}(f)\|_{f_{p,q}^s} \lesssim \|f\|_{F_{p,q}^s}.$$

(ii) *If  $1/p - 1 < s < 1$ , then for all  $f \in B_{p,q}^s$*

$$\|\mathbf{c}(f)\|_{b_{p,q}^s} \lesssim \|f\|_{B_{p,q}^s}.$$

*Proof.* To avoid dealing separately with  $\tilde{h}_{j,\nu}$  with  $\nu$  even or odd, we prove a slightly more general result. For a fixed  $\delta \in [0, 1]$  and for  $j \geq 0$  and  $\mu \in \mathbb{Z}$ , consider the shifted Haar function

$$h_{j,\mu}^\delta(x) := h_{j,\mu}(x - \delta 2^{-j}) = h(2^j x - (\mu + \delta)),$$

whose support is the interval  $I_{j,\mu} + \delta 2^{-j}$ . When  $j = -1$ , we just let  $h_{-1,\mu}^\delta = h_{-1,\mu} = \mathbb{1}_{[\mu, \mu+1)}$ . Part (i) will then be a consequence of the following estimate

$$(4.2) \quad \left\| \left( \sum_{j=-1}^{\infty} 2^{jsq} \left| \sum_{\mu \in \mathbb{Z}} 2^j \langle f, h_{j,\mu}^\delta \rangle \mathbb{1}_{I_{j,\mu}} \right|^q \right)^{1/q} \right\|_p \lesssim \|f\|_{F_{p,q}^s},$$

where the constants are independent of  $\delta \in [0, 1]$ . Indeed, (4.1) follows from (4.2) applied to  $\delta = 0$  and  $\delta = 1/2$ .

We now prove (4.2) for a fixed  $\delta \in [0, 1]$ . In the proof below we denote by  $\Delta_{j,\mu}$  the set of discontinuity points of  $h_{j,\mu}^\delta$ , that is

$$\Delta_{j,\mu} = \left\{ (\mu + \delta + i)2^{-j} : i = 0, \frac{1}{2}, 1 \right\}.$$

**Lemma 4.2.** *Let  $g \in L_1^{\text{loc}}(\mathbb{R})$ ,  $k \in \mathbb{N}_0$  and  $j \geq -1$ . Then*

a) *If  $k \geq j$  then*

$$(4.3) \quad |2^j \langle g, L_k h_{j,\mu}^\delta \rangle| \lesssim 2^{-(k-j)} \sum_{z \in \Delta_{j,\mu}} \mathfrak{M}_k(g)(z);$$

moreover, for every  $A > 0$ ,

$$(4.4) \quad \begin{aligned} |2^j \langle g, L_k h_{j,\mu}^\delta \rangle| \mathbb{1}_{I_{j,\mu}}(x) &\lesssim 2^{-(k-j)} \mathfrak{M}_j^*(g)(x) \\ &\lesssim 2^{-(k-j)} 2^{(k-j)A} \mathfrak{M}_{k,A}^{**}(g)(x). \end{aligned}$$

b) *If  $j > k$  then, for every  $A > 0$ ,*

$$(4.5) \quad \begin{aligned} |2^j \langle g, L_k h_{j,\mu}^\delta \rangle| \mathbb{1}_{I_{j,\mu}}(x) &\lesssim 2^{-(j-k)} \mathfrak{M}_k(g)(x) \\ &\lesssim 2^{-(j-k)} \mathfrak{M}_{k,A}^{**}(g)(x). \end{aligned}$$

*Proof.* a) If  $k \geq j$  then the function  $L_k h_{j,\mu}^\delta = \beta_k * h_{j,\mu}^\delta$  is supported in  $\Delta_{j,\mu} + O(2^{-k})$  and has size

$$\|\beta_k * h_{j,\mu}^\delta\|_\infty \leq \|\beta_k\|_1 = O(1).$$

This immediately gives (4.3). Now, if  $z \in \Delta_{j,\mu}$  and  $x \in I_{j,\mu}$  we have

$$\begin{aligned} \mathfrak{M}_k g(z) &= \sup_{|h| \leq 2^{-k}} |g(z+h)| \leq \sup_{|u| \leq 2^{-j+2}} |g(x+u)| = \mathfrak{M}_j^*(g)(x) \\ &\lesssim 2^{(k-j)A} \mathfrak{M}_{k,A}^{**}(g)(x), \end{aligned}$$

which together with (4.3) proves (4.4).

b) If  $k < j$  and  $x \in I_{j,\mu}$ , then the function  $L_k h_{j,\mu}^\delta = \beta_k * h_{j,\mu}^\delta$  is supported in  $x + O(2^{-k})$ , and we can bound its size by

$$(4.6) \quad |L_k h_{j,\mu}^\delta(u)| \lesssim 2^{2(k-j)}.$$

This last assertion follows from the property  $\int h_{j,\mu}^\delta = 0$ , by writing

$$\begin{aligned} |L_k h_{j,\mu}^\delta(u)| &= \left| \int (\beta_k(u-y) - \beta_k(u-x)) h_{j,\mu}^\delta(y) dy \right| \\ &= \left| \int_0^1 \int_0^1 \beta_k'(u - (1-t)x - ty) dt (x-y) h_{j,\mu}^\delta(y) dy \right| \\ &\lesssim 2^{2k-2j}, \end{aligned}$$

using in the last step that  $|x-y| \leq 2^{-j+1}$  when  $x \in I_{j,\mu}$  and  $y \in \text{supp } h_{j,\mu}^\delta$ . Combining the above support and size estimates, one easily obtains (4.5).  $\square$

We continue the proof of Proposition 4.1.i. Let  $f \in F_{p,q}^s$ , and write it as  $f = \sum_{k=0}^{\infty} L_k f_k$  with  $f_k = \Lambda_k f$  as in (2.5). Note that, since  $F_{p,q}^s \subset \mathcal{B}$ , we have

$$\langle f, h_{j,\mu}^\delta \rangle = \sum_{k=0}^{\infty} \langle f_k, L_k h_{j,\mu}^\delta \rangle.$$

Now, the estimates in Lemma 4.2, suitably applied to each  $f_k$ , can be grouped into

$$(4.7) \quad \begin{aligned} \sum_{\mu} |2^j \langle f, h_{j,\mu}^\delta \rangle| \mathbb{1}_{I_{j,\mu}}(x) &\lesssim \sum_{k \geq 0} 2^{-|k-j|} 2^{A(k-j)_+} \mathfrak{M}_{k,A}^{**}(f_k)(x) \\ (\ell = k-j) &\lesssim \sum_{\ell \in \mathbb{Z}} a(\ell, A) \mathfrak{M}_{j+\ell,A}^{**}[f_{j+\ell}](x) =: G_j(x), \end{aligned}$$

where we set  $f_m \equiv 0$  for  $m < 0$  and

$$(4.8) \quad a(\ell, A) = \begin{cases} 2^\ell & : \ell < 0, \\ 2^{(A-1)\ell} & : \ell \geq 0. \end{cases}$$

At this point one takes  $L_p(\ell_q)$  quasi-norms of the above expressions. Letting  $u := \min\{p, q, 1\}$ , and using the  $u$ -triangle inequality we obtain

$$(4.9) \quad \begin{aligned} &\left\| \left( \sum_{j \geq -1} |2^{js} G_j|^q \right)^{1/q} \right\|_p \\ &\lesssim \left( \sum_{\ell \in \mathbb{Z}} \left[ a(\ell, A) 2^{-s\ell} \left\| \left( \sum_{j \in \mathbb{Z}} |2^{(j+\ell)s} \mathfrak{M}_{j+\ell,A}^{**}[f_{j+\ell}](x)|^q \right)^{1/q} \right\|_p \right]^u \right)^{1/u} \\ &\lesssim \|f\|_{F_{p,q}^s} \left( \sum_{\ell \in \mathbb{Z}} (a(\ell, A) 2^{-s\ell})^u \right)^{1/u} \lesssim \|f\|_{F_{p,q}^s}, \end{aligned}$$

where in the last line we use Peetre's maximal inequality (2.10) and  $A > \max\{1/p, 1/q\}$ , and in the last step we additionally need that  $A-1 < s < 1$ . This can always be achieved for an appropriate choice of  $A$  because of our assumption  $\max\{\frac{1}{p}, \frac{1}{q}\} - 1 < s < 1$ .

As before, part (ii) in Proposition 4.1 will be a consequence of the more general estimate

$$(4.10) \quad \left( \sum_{j=-1}^{\infty} 2^{j(s-1/p)q} \left( \sum_{\mu \in \mathbb{Z}} |2^j \langle f, h_{j,\mu}^\delta \rangle|^p \right)^{q/p} \right)^{1/q} \lesssim \|f\|_{B_{p,q}^s},$$

for  $\delta \in [0, 1]$ . Notice that

$$(4.11) \quad B_j := 2^{-j/p} \left( \sum_{\mu} |2^j \langle f, h_{j,\mu}^\delta \rangle|^p \right)^{\frac{1}{p}} = \left\| \sum_{\mu} |2^j \langle f, h_{j,\mu}^\delta \rangle| \mathbb{1}_{I_{j,\mu}} \right\|_p.$$

We shall argue a bit differently to refine the pointwise estimate in (4.7). Observe from Lemma 4.2 that we can also write

$$\begin{aligned} \left\| \sum_{\mu} 2^j \langle f_k, h_{j,\mu}^{\delta} \rangle \mathbb{1}_{I_{j,\mu}} \right\|_p &\lesssim 2^{-|k-j|} \|\mathfrak{M}_{\min\{j,k\}}^*(f_k)\|_p \\ &\stackrel{\text{by (2.13)}}{\lesssim} 2^{-|k-j|} 2^{\frac{(k-j)_+}{p}} \|\mathfrak{M}_k(f_k)\|_p \\ &= a(k-j, \frac{1}{p}) \|\mathfrak{M}_k(f_k)\|_p, \end{aligned}$$

using in the last step the definition of  $a(\ell, A)$  in (4.8). So, letting as before  $u := \min\{p, q, 1\}$ , and using the  $u$ -triangle inequality we obtain

$$\begin{aligned} &\left( \sum_{j \geq -1} (2^{js} B_j)^q \right)^{\frac{1}{q}} \lesssim \\ (4.12) \quad &\lesssim \left( \sum_{\ell \in \mathbb{Z}} \left[ a(\ell, \frac{1}{p}) 2^{-s\ell} \left( \sum_{j \in \mathbb{Z}} |2^{(j+\ell)s}| \|\mathfrak{M}_{j+\ell}[f_{j+\ell}]\|_p^q \right)^{1/q} \right]^u \right)^{1/u} \\ &\lesssim \|f\|_{B_{p,q}^s} \left( \sum_{\ell \in \mathbb{Z}} (a(\ell, \frac{1}{p}) 2^{-s\ell})^u \right)^{1/u} \lesssim \|f\|_{B_{p,q}^s}. \end{aligned}$$

Observe that this time we apply the simpler estimate  $\|\mathfrak{M}_k[f_k]\|_p \lesssim \|f_k\|_p$ , see (2.12), while the very last step requires  $1/p - 1 < s < 1$ .  $\square$

*Remark 4.3.* In view of the examples in [21], the condition  $s > 1/q - 1$  is necessary in Proposition 4.1.i, even for the validity of the weaker inequality

$$(4.13) \quad \|f\|_{F_{p,q}^{s,\text{dyad}}} := \left\| \left( \sum_{j=-1}^{\infty} 2^{jsq} \left| \sum_{\mu \in \mathbb{Z}} 2^j \langle f, h_{j,\mu} \rangle \mathbb{1}_{I_{j,\mu}} \right|^q \right)^{1/q} \right\|_p \lesssim \|f\|_{F_{p,q}^s}.$$

Indeed, arguing as in [21, §6], for each  $N \geq 2$  one constructs a (smooth) function  $f = f_N \in F_{p,q}^{1/q-1}(\mathbb{R})$  and a finite set<sup>2</sup>  $E = E_N \subset \mathcal{H}$  such that if  $0 < q \leq p < \infty$  then

$$(4.14) \quad \|f\|_{F_{p,q}^{1/q-1}(\mathbb{R})} \lesssim N^{1/q} \quad \text{and} \quad \|P_E(f)\|_{F_{p,q}^{1/q-1}(\mathbb{R})} \geq N^{1+\frac{1}{q}},$$

where  $P_E(f)$  denotes the projection onto  $\text{span}\{h\}_{h \in E}$ . Now observe that

$$\|f\|_{F_{p,q}^{s,\text{dyad}}} \geq \|P_E(f)\|_{F_{p,q}^{s,\text{dyad}}} \gtrsim \|P_E(f)\|_{F_{p,q}^s},$$

using in the last step the inequality in (4.20) (which holds for  $s = 1/q - 1$ ). Thus, we conclude from (4.14) that

$$\|f\|_{F_{p,q}^{1/q-1,\text{dyad}}} \gtrsim N \|f\|_{F_{p,q}^{1/q-1}}.$$

We turn to the converse inequalities in Theorems 1.2 and 1.3, which will be proved in Proposition 4.9 below. Before doing so we shall need some

<sup>2</sup>In the notation of [21, §6], one should consider sets  $A$  of *consecutive* Haar frequencies, so that the associated “density” number in [21, (43)] takes the value  $Z = N$ .

results concerning the inclusions  $F_{p,q}^{s,\text{dyad}} \hookrightarrow F_{p,q}^s$  (and likewise for  $B$ -spaces) for the usual Haar system. Results of this nature can be found in [28, Proposition 2.6], but we give here direct proofs which are valid in a larger range of indices. These may have an independent interest.

The first result is obtained by duality from Proposition 4.1, but the range of indices is restricted to  $1 < p, q < \infty$ . Recall the definitions of the sequence spaces  $b_{p,q}^s, f_{p,q}^s$  in (1.19) and (1.20).

**Proposition 4.4.** *Let  $1 < p, q < \infty$ .*

(i) *If  $-1 < s < \min\{1/p, 1/q\}$  and  $\beta = \{\beta_{j,\mu}\} \in f_{p,q}^s$  then*

$$(4.15) \quad f := \sum_{j=-1}^{\infty} \sum_{\mu \in \mathbb{Z}} \beta_{j,\mu} h_{j,\mu}$$

*converges in  $F_{p,q}^s(\mathbb{R})$  and*

$$(4.16) \quad \|f\|_{F_{p,q}^s} \lesssim \|\beta\|_{f_{p,q}^s}$$

(ii) *If  $-1 < s < 1/p$  and  $\beta = \{\beta_{j,\mu}\} \in b_{p,q}^s$  then (4.15) converges in  $B_{p,q}^s(\mathbb{R})$  and*

$$(4.17) \quad \|f\|_{B_{p,q}^s} \lesssim \|\beta\|_{b_{p,q}^s}.$$

*Proof.* Consider the following duality pairing for sequences

$$\begin{aligned} \langle \alpha, \beta \rangle &:= \sum_j 2^{-j} \sum_{\mu} \alpha_{j,\mu} \beta_{j,\mu} \\ &= \sum_j \int_{-\infty}^{\infty} \left( \sum_{\mu} \alpha_{j,\mu} \mathbb{1}_{I_{j,\mu}}(x) \right) \left( \sum_{\mu'} \beta_{j,\mu'} \mathbb{1}_{I_{j,\mu'}}(x) \right) dx \leq \|\alpha\|_{f_{p,q}^s} \|\beta\|_{f_{p',q'}^{-s}}. \end{aligned}$$

We define the so-called *analysis* (or *sampling*) *operator*  $\mathcal{A}$  by

$$(4.18) \quad \begin{aligned} \mathcal{A} : F_{p,q}^s(\mathbb{R}) &\longrightarrow f_{p,q}^s \\ f &\longmapsto \mathcal{A}f(j, \mu) = 2^j \langle f, h_{j,\mu} \rangle. \end{aligned}$$

Its dual operator  $\mathcal{A}'$  (the *synthesis operator*) is given by

$$(4.19) \quad \mathcal{A}'\beta(x) = \sum_{j,\mu} \beta_{j,\mu} h_{j,\mu}(x).$$

Indeed, if  $f \in F_{p,q}^s$  and  $\beta = \{\beta_{j,\mu}\}$  is a finite sequence, then

$$\begin{aligned} \langle \mathcal{A}f, g' \rangle &= \sum_j 2^{-j} \sum_{\mu} 2^j \langle f, h_{j,\mu} \rangle \cdot \beta_{j,\mu} \\ &= \int f(x) \sum_j \sum_{\mu} \beta_{j,\mu} h_{j,\mu}(x) dx = \int f(x) \mathcal{A}'\beta(x) dx. \end{aligned}$$

By Proposition 4.1, the operator  $\mathcal{A} : F_{p,q}^s(\mathbb{R}) \rightarrow f_{p,q}^s$  is bounded when  $\max\{1/p - 1, 1/q - 1\} < s < 1$ . Hence, if we assume that  $1 \leq p, q < \infty$ , then

$\mathcal{A}'$  will be bounded from  $f_{p',q'}^{-s}$  to  $F_{p',q'}^{-s}(\mathbb{R})$ , where

$$-1 < -s < \min\{1/p', 1/q'\}.$$

In other words, if  $-1 < s < \min\{1/p, 1/q\}$  and  $1 < p, q \leq \infty$  then

$$\left\| \sum_{j,\mu} \beta_{j,\mu} h_{j,\mu} \right\|_{F_{p,q}^s} \lesssim \|\beta\|_{f_{p,q}^s}.$$

The proof of part (ii) is completely analogous.  $\square$

In the following proposition we extend the previous estimate to the quasi-Banach range of parameters. The proof, which is now more involved, uses a non-trivial interpolation argument from [28, Proposition 2.6]. The range of indices we obtain is larger than in [28]; see Figure 4 below.




FIGURE 4. Parameter domain for the validity of Proposition 4.5, for  $F_{p,q}^s(\mathbb{R})$  (left figure) and  $B_{p,q}^s(\mathbb{R})$  (right figure).

**Proposition 4.5.** *Let  $0 < p, q \leq \infty$ .*

(i) *If  $p < \infty$  and  $\max\{1/p, 1/q, 1\} - 2 < s < \min\{1/p, 1/q\}$ , then for every  $\beta \in f_{p,q}^s$  the series in (4.15) converges to a distribution  $f$  in  $F_{p,q}^s(\mathbb{R})$ , and it holds*

$$(4.20) \quad \|f\|_{F_{p,q}^s} \lesssim \|\beta\|_{f_{p,q}^s}.$$

*Moreover if  $q < \infty$  the series in (4.15) converges unconditionally in  $F_{p,q}^s$ , and otherwise in  $F_{p,\infty}^{s-\varepsilon}(\mathbb{R})$ , for all  $\varepsilon > 0$ .*

(ii) *If  $\max\{1/p, 1\} - 2 < s < 1/p$ , then for every  $\beta \in b_{p,q}^s$  the series in (4.15) converges to a distribution  $f$  in  $B_{p,q}^s(\mathbb{R})$ , and it holds*

$$(4.21) \quad \|f\|_{B_{p,q}^s} \lesssim \|\beta\|_{b_{p,q}^s}.$$

*Moreover, if  $q < \infty$  the series in (4.15) converges unconditionally in the norm of  $B_{p,q}^s$ , and otherwise in the quasi-norm of  $B_{p,\infty}^{s-\varepsilon}(\mathbb{R})$ , for all  $\varepsilon > 0$ .*

*Proof.* It suffices to prove the results when  $f = \sum_{j \geq -1} \sum_{\mu \in \mathbb{Z}} \beta_{j,\mu} h_{j,\mu}$ , with  $(\beta_{j,\mu})$  a finite sequence of scalars. The other assertions will then follow by completeness of the spaces.

*Step 1.* Let  $L_k$ ,  $k \geq 0$ , be the local convolution operators from Section 2.1. If  $j \geq -1$  is fixed, then we have

$$(4.22) \quad \left| L_k \left( \sum_{\mu \in \mathbb{Z}} \beta_{j,\mu} h_{j,\mu} \right) (x) \right| \leq \sum_{\mu \in \mathbb{Z}} |\beta_{j,\mu}| \cdot |(L_k h_{j,\mu})(x)| =: G_{j,k}(x).$$

Arguing as in Lemma 4.2, one sees that, in the case  $\ell = j - k > 0$ , then

$$(4.23) \quad \text{supp } L_k h_{j,\mu} \subset \mu 2^{-j} + O(2^{-k}) \quad \text{and} \quad |(L_k h_{j,\mu})(x)| \lesssim 2^{-2\ell},$$

while in the case  $\ell = j - k \leq 0$  we have

$$(4.24) \quad \text{supp } L_k h_{j,\mu} \subset \mu 2^{-j} + O(2^{-j}) \quad \text{and} \quad |(L_k h_{j,\mu})(x)| \lesssim 1.$$

The following lemma is a variation of a result by Kyriazis; see [17, Lem. 7.1]. For  $r > 0$  we denote

$$M_r(g)(x) = [M(|g|^r)(x)]^{1/r},$$

where  $M$  is the usual Hardy-Littlewood maximal operator.

**Lemma 4.6.** *Let  $0 < r \leq 1$ . Then, it holds*

$$(4.25) \quad G_{j,k}(x) \lesssim a(j-k) M_r \left( \sum_{\mu} |\beta_{j,\mu}| \mathbb{1}_{I_{j,\mu}} \right) (x).$$

where  $a(\ell) = 2^{-2\ell} 2^{\ell/r}$  if  $\ell > 0$ , and  $a(\ell) = 1$  if  $\ell \leq 0$ .

*Proof.* Assume that  $\ell = j - k > 0$ . If  $x \in I_{j,\nu}$ , for some fixed  $\nu \in \mathbb{Z}$ , then the size and support estimates in (4.23) give

$$G_{j,k}(x)^r \lesssim \sum_{\mu: |\mu-\nu| \leq 2^{j-k}} |\beta_{j,\mu}|^r 2^{-2\ell r}.$$

On the other hand, if  $x \in I_{j,\nu}$ ,

$$\begin{aligned} M \left( \sum_{\mu} |\beta_{j,\mu}|^r \mathbb{1}_{I_{j,\mu}} \right) (x) &\gtrsim 2^k \int_{|y-2^{-j}\nu| \leq 2 \cdot 2^{-k}} \left( \sum_{\mu} |\beta_{j,\mu}|^r \mathbb{1}_{I_{j,\mu}} \right) (y) dy \\ &\geq 2^{k-j} \sum_{\mu: |\mu-\nu| \leq 2^{j-k}} |\beta_{j,\mu}|^r. \end{aligned}$$

These two estimates clearly give (4.25) in the case  $\ell > 0$ . The case  $\ell \leq 0$  is proved similarly using (4.24).  $\square$

We continue with the proof of Proposition 4.5.i. Below we shall agree that  $\beta_{m,\mu} = 0$  and  $G_{m,k} \equiv 0$  whenever  $m < -1$ . Then letting  $u = \min\{p, q, 1\}$ ,

we can apply the  $u$ -triangle inequality and the above results to obtain

$$\begin{aligned}
\|f\|_{F_{p,q}^s} &= \left\| \left( \sum_{k \geq 0} |2^{ks} L_k f|^q \right)^{1/q} \right\|_p \leq \left\| \left( \sum_{k \geq 0} (2^{ks} \sum_{\ell \in \mathbb{Z}} G_{k+\ell,k})^q \right)^{1/q} \right\|_p \\
&\lesssim \left[ \sum_{\ell \in \mathbb{Z}} \left( a(\ell) 2^{-\ell s} \right)^u \left\| \left( \sum_{k \geq 0} \left[ 2^{(k+\ell)s} M_r \left( \sum_{\mu \in \mathbb{Z}} |\beta_{k+\ell,\mu}| \mathbb{1}_{I_{k+\ell,\mu}} \right) \right]^q \right)^{\frac{1}{q}} \right\|_p^u \right]^{\frac{1}{u}} \\
&\lesssim \left\| \left( \sum_{m \in \mathbb{Z}} \left( 2^{ms} \sum_{\mu \in \mathbb{Z}} |\beta_{m,\mu}| \mathbb{1}_{I_{m,\mu}} \right)^q \right)^{\frac{1}{q}} \right\|_p,
\end{aligned}$$

where the last line is justified by the Fefferman-Stein inequality (2.7), provided  $r < \min\{p, q, 1\}$ , and the finite summation in  $\ell \in \mathbb{Z}$  holds whenever  $1/r - 2 < s < 0$ . Such an  $r$  can always be chosen under the assumption

$$\max\{1/p, 1/q, 1\} - 2 < s < 0$$

(which in particular implies  $p, q > 1/2$ ). We shall see in Step 3 below how to enlarge this range to cover as well the cases  $s \geq 0$ .

*Step 2.* We now prove (4.21). The same notation as above gives

$$\|f\|_{B_{p,q}^s} = \left( \sum_{k \geq 0} (2^{ks} \|L_k f\|_p)^q \right)^{1/q} \leq \left( \sum_{k \geq 0} (2^{ks} \left\| \sum_{\ell \in \mathbb{Z}} G_{k+\ell,k} \right\|_p)^q \right)^{1/q}.$$

At this point we distinguish two cases,  $\ell > 0$  and  $\ell \leq 0$ . In the first case we use literally the same arguments as above; since for the  $\ell_q(L_p)$  quasi-norm we just use the *scalar* Hardy-Littlewood maximal inequality we only need to impose  $r < \min\{1, p\}$ , together with  $s > 1/r - 2$ . Such an  $r$  can always be chosen under the assumption

$$\max\{1/p, 1\} - 2 < s.$$

To control the sum over  $\ell \leq 0$  we must replace the crude bound in (4.24) by the sharper estimate

$$(4.26) \quad \text{supp } L_k h_{j,\mu} \subset \Delta_{j,k} + O(2^{-j}) \quad \text{and} \quad |(L_k h_{j,\mu})(x)| \lesssim 1,$$

where  $\Delta_{j,\mu}$  are the discontinuity points of  $h_{j,\mu}$ ; see the proof of Lemma 4.2.a. So, if  $\ell = j - k \leq 0$  we have

$$\|G_{k+\ell,k}\|_p \lesssim 2^{-(k+\ell)/p} \left( \sum_{\mu \in \mathbb{Z}} |\beta_{k+\ell,\mu}|^p \right)^{1/p}.$$



Using as before the  $u$ -triangle inequality, this yields

$$\begin{aligned} & \left( \sum_{k \geq 0} \left( 2^{ks} \left\| \sum_{\ell \leq 0} G_{k+\ell, k} \right\|_p \right)^q \right)^{1/q} \\ & \lesssim \left[ \sum_{\ell \leq 0} 2^{u(s-\frac{1}{p})\ell} \left( \sum_{k \geq 0} \left( 2^{(k+\ell)(s-\frac{1}{p})} \left( \sum_{\mu \in \mathbb{Z}} |\beta_{k+\ell, \mu}|^p \right)^{1/p} \right)^{\frac{u}{q}} \right)^{\frac{1}{u}} \right] \\ & \lesssim \left( \sum_{\ell \leq 0} 2^{u(s-\frac{1}{p})\ell} \right)^{\frac{1}{u}} \left( \sum_{m \geq \mathbb{Z}} \left( 2^{m(s-\frac{1}{p})} \left( \sum_{\mu \in \mathbb{Z}} |\beta_{m, \mu}|^p \right)^{1/p} \right)^{\frac{1}{q}} \right), \end{aligned}$$

where the sum in  $\ell \leq 0$  is a finite constant due to the assumption  $s < 1/p$ . This completes the proof of part (ii) in Proposition 4.5.

*Step 3.* In the Triebel-Lizorkin case, the direct argument in Step 1 only allows for  $s < 0$  (and  $p, q > 1/2$ ), which is the desired region only when  $q = \infty$  or  $p \rightarrow \infty$ . By Step 2, the range of parameters can be extended to  $s < 1/p$  when  $p = q$ . Then, a complex interpolation argument in the three indices  $(s, 1/p, 1/q)$ , as proposed by Triebel in [28, Prop. 2.6], gives the validity of the result for all  $\max\{1/p, 1/q, 1\} - 2 < s < \min\{1/p, 1/q\}$ ; see Figure 5.  $\square$




FIGURE 5. Parameter domain for  $F$ -spaces in Steps 1 and 2 (left figure), and after the interpolation argument in Step 3 (right figure) of the proof of Proposition 4.5.

*Remark 4.7.* We remark that the decomposition of a distribution  $f \in \mathcal{S}'$  as an infinite series

$$(4.27) \quad f = \sum_{j \geq -1} \sum_{\mu \in \mathbb{Z}} \beta_{j, \mu} h_{j, \mu}$$

may not necessarily be unique. For instance, the Dirac delta satisfies

$$\delta = \mathbb{1}_{[0,1)} + \sum_{j=0}^{\infty} 2^j h_{j,0} = \mathbb{1}_{[-1,0)} - \sum_{j=0}^{\infty} 2^j h_{j,-1} \quad \text{in } \mathcal{S}'(\mathbb{R}).$$

In this example, the coefficient sequences belong to  $b_{p,q}^s$  if  $s < \frac{1}{p} - 1$  (or  $s = \frac{1}{p} - 1$  and  $q = \infty$ ), and the same happens for the property  $\delta \in B_{p,q}^s(\mathbb{R})$ .

For such cases of non-uniqueness, Proposition 4.5 should be interpreted as

$$\|f\|_{B_{p,q}^s} \lesssim \inf \left\{ \|(\beta_{j,\mu})\|_{b_{p,q}^s} : (4.27) \text{ holds} \right\},$$

and likewise for the  $F_{p,q}^s$ -quasinorms.

The next result shows that uniqueness holds when  $s > 1/p - 1$ .

**Corollary 4.8.** *Let  $0 < p, q \leq \infty$  and  $s \in \mathbb{R}$ .*

(i) *If  $p < \infty$  and  $\max\{1/p - 1, 1/q - 2\} < s < \min\{1/p, 1/q\}$  then for all  $f \in \mathcal{B}$  it holds*

$$(4.28) \quad \|f\|_{F_{p,q}^s} \lesssim \left\| \left( \sum_{j=-1}^{\infty} \left| 2^{js} \sum_{\mu \in \mathbb{Z}} 2^j \langle f, h_{j,\mu} \rangle \mathbb{1}_{I_{j,\mu}} \right|^q \right)^{1/q} \right\|_p.$$

(ii) *If  $1/p - 1 < s < 1/p$  then for all  $f \in \mathcal{B}$  it holds*

$$(4.29) \quad \|f\|_{B_{p,q}^s} \lesssim \left( \sum_{j=-1}^{\infty} 2^{j(s-1/p)q} \left( \sum_{\mu \in \mathbb{Z}} |2^j \langle f, h_{j,\mu} \rangle|^p \right)^{q/p} \right)^{1/q}.$$

*Proof.* Let  $f \in \mathcal{B}$  be such that the right hand side of (4.28) is finite. By Proposition 4.5 this implies the convergence of the series

$$g := \sum_{j=-1}^{\infty} \sum_{\mu \in \mathbb{Z}} 2^j \langle f, h_{j,\mu} \rangle h_{j,\mu},$$

to some distribution  $g \in F_{p,q}^s \hookrightarrow \mathcal{B}$ . Due to the range of parameters, and the convergence in  $F_{p,q}^{s-\varepsilon}(\mathbb{R})$  for  $\varepsilon$  small enough, we also have convergence in  $\mathcal{B}$ . We deduce that  $\langle g, h \rangle = \langle f, h \rangle$  for all  $h \in \mathcal{H}$ , and therefore, by Proposition 1.1, that  $f = g$ . Finally, Proposition 4.5 gives (4.28). The proof for (4.29) works analogously.  $\square$

We finally turn to the remaining implications in Theorems 1.2 and 1.3, which are also valid in a larger range.

**Proposition 4.9.** *Let  $0 < p, q \leq \infty$  and  $s \in \mathbb{R}$  be such that*

$$\frac{1}{p} - 1 < s < 1 + \frac{1}{p}.$$

(i) *If  $p < \infty$  and additionally  $1/q - 2 < s < 2 + 1/q$ , then for all  $f \in \mathcal{B}$*

$$(4.30) \quad \|f\|_{F_{p,q}^s} \lesssim \|\mathbf{c}(f)\|_{f_{p,q}^s}$$

(ii) *For all  $f \in \mathcal{B}$  it holds*

$$(4.31) \quad \|f\|_{B_{p,q}^s} \lesssim \|\mathbf{c}(f)\|_{b_{p,q}^s}$$

*Proof.* Note that, for all  $f \in \mathcal{S}'(\mathbb{R})$ , it holds

$$(4.32) \quad \|f\|_{F_{p,q}^s} \approx \|f\|_{F_{p,q}^{s-1}} + \|f'\|_{F_{p,q}^{s-1}},$$

see e.g. [25, 2.3.8]. We shall bound each of the summands in (4.32) by the right hand side of (4.30).

Clearly  $\|f\|_{F_{p,q}^{s-1}} \lesssim \|f\|_{B_{p,p}^r}$  for any  $s-1 < r$ . We distinguish two cases. In case  $s > 1/p$  we choose  $r := 1/p - \varepsilon$ , for a sufficiently small  $\varepsilon > 0$  so that

$$(4.33) \quad 1/p - 1 < r < 1/p \quad \text{and} \quad s - 1 < r < s;$$

this is possible by the assumption  $s < 1 + 1/p$ . In case  $s \leq 1/p$  we put  $r := s - \varepsilon$ , for some  $\varepsilon > 0$  so that (4.33) also holds (this time using the assumption  $s > 1/p - 1$ ). Hence, from the embeddings  $B_{pp}^r \hookrightarrow F_{pq}^{s-1}$  and  $f_{pq}^s \hookrightarrow b_{pp}^r$ , together with Corollary 4.8.ii, we obtain

$$(4.34) \quad \|f\|_{F_{p,q}^{s-1}} \lesssim \|f\|_{B_{p,p}^r} \lesssim \|\{2^j \langle f, h_{j,\mu} \rangle\}_{j,\mu}\|_{b_{pp}^r} \lesssim \|\{\mathfrak{c}_{j,\mu}(f)\}_{j,\mu}\|_{f_{pq}^s}.$$

We now take care of the second term in (4.32). Here we quote the analog of Corollary 4.8.i for the Chui-Wang system  $\{\psi_{j,\mu}\}$ , which can be obtained from [8, Proposition 5.4] and Remark 2.3 above. Letting  $r = s - 1$ , this gives

$$(4.35) \quad \|f'\|_{F_{p,q}^r} \lesssim \left\| \left( \sum_{j \geq -1} 2^{jrq} \left| \sum_{\mu \in \mathbb{Z}} 2^j \langle f', \psi_{j,\mu} \rangle \mathbb{1}_{I_{j,\mu}}(x) \right|^q \right)^{1/q} \right\|_p,$$

provided

$$\max\{1/p - 1, 1/q - 2\} - 1 < r < 1 + \min\{1/p, 1/q\},$$

which holds when  $1/p - 1 < s < 1/p + 1$  and  $1/q - 2 < s < 2 + 1/q$ .

Now, recall that

$$(4.36) \quad \psi_{j,\mu}(x) = \sum_{k \in \mathbb{Z}} b_k \mathcal{N}_2(2^{j+1}x - (2\mu + k)),$$

for a finite sequence of coefficients  $b_k$ ; see (2.16). In addition, we know from Lemma 2.1 that

$$(4.37) \quad \langle f', \mathcal{N}_2(2^{j+1} \cdot -\nu) \rangle = -2^{j+1} \langle f, \tilde{h}_{j,\nu} \rangle, \quad j \geq 0, \nu \in \mathbb{Z}$$

(and a similar expression for  $j = -1$ ). So, combining (4.36) and (4.37), we obtain an estimate for  $|2^j \langle f', \psi_{j,\mu} \rangle|$  in terms of  $\mathfrak{c}_{j,\nu}(f)$  coefficients, which inserted into (4.35) gives

$$\|f'\|_{F_{p,q}^r} \lesssim \|(\mathfrak{c}_{j,\mu}(f))_{j,\mu}\|_{f_{pq}^s}.$$

This, together with (4.34) concludes the proof of part (i).

The result for  $B_{p,q}^s$  in part (ii) goes similarly, using instead Corollary 4.8.ii, and the corresponding version for the Chui-Wang system  $\{\psi_{j,\mu}\}$  which can be obtained from [8, Proposition 5.3].  $\square$

We are finally ready to give the

*Proof of Theorems 1.2 and 1.3.* Just combine Proposition 4.1 and Proposition 4.9. Note that the smallest range of parameters corresponds to that in Proposition 4.1.  $\square$

5.  $W_p^1$  AND  $BV$ : PROOF OF THEOREM 1.4

The proof has three steps. We use the classical norm definition for  $W_p^1(\mathbb{R})$ , when  $1 \leq p \leq \infty$ , namely

$$\|f\|_{W_p^1} := \|f\|_p + \|f'\|_p.$$

*Step 1.* We show that, for  $1 \leq p \leq \infty$ , it holds

$$(5.1) \quad \sup_{j \geq -1} 2^{j(1-1/p)} \left( \sum_{\nu \in \mathbb{Z}} |2^j \langle f, \tilde{h}_{j,\nu} \rangle|^p \right)^{1/p} \lesssim \|f\|_{W_p^1}, \quad f \in W_p^1(\mathbb{R}),$$

moreover

$$(5.2) \quad \sup_{j \geq -1} \sum_{\nu \in \mathbb{Z}} |2^j \langle f, \tilde{h}_{j,\nu} \rangle| \lesssim \|f\|_{BV}, \quad f \in BV(\mathbb{R}),$$

In view of Lemma 2.1, for every  $j \geq 0$  we have

$$(5.3) \quad |2^{j+1} \langle f, \tilde{h}_{j,\nu} \rangle| \leq \int |f'(x) \mathcal{N}_{2;j+1,\nu}(x)| dx \lesssim 2^{-j/p'} \left[ \int_{\tilde{I}_{j,\nu}} |f'|^p dx \right]^{1/p},$$

where  $\tilde{I}_{j,\nu} = \text{supp } \mathcal{N}_{2;j+1,\nu} = [\nu/2^{j+1}, (\nu+2)/2^{j+1}]$ . Hence,

$$\sup_{j \geq 0} 2^{j(1-1/p)} \left( \sum_{\nu \in \mathbb{Z}} |2^j \langle f, \tilde{h}_{j,\nu} \rangle|^p \right)^{1/p} \lesssim \|f'\|_p.$$

Likewise, if  $j = -1$  we simply have

$$|\langle f, \tilde{h}_{-1,\nu} \rangle| = \left| \int_{I_{0,\nu}} f(x) dx \right| \leq \left( \int_{I_{0,\nu}} |f(x)|^p dx \right)^{1/p}$$

and hence

$$\left( \sum_{\nu \in \mathbb{Z}} |\langle f, \tilde{h}_{-1,\nu} \rangle|^p \right)^{1/p} \lesssim \|f\|_p.$$

We have thus established (5.1). To handle (5.2) we work with an approximation of the identity,  $\{\Phi_\ell\}$  where  $\Phi_\ell = 2^\ell \Phi(2^\ell \cdot)$  with  $\Phi \in C_c^\infty$  and  $\int \Phi = 1$ . Let  $f \in BV$  (which implies  $f \in L_\infty$ ). Then  $\Phi_\ell * f \in W_1^1$  with  $\|\Phi_\ell * f\|_{W_1^1} \lesssim \|f\|_{BV}$  and  $\Phi_\ell * f(x) \rightarrow f(x)$  almost everywhere. By dominated convergence  $\langle \Phi_\ell * f, \tilde{h}_{j,\nu} \rangle \rightarrow \langle f, \tilde{h}_{j,\nu} \rangle$  and by a further application of Fatou's lemma and (5.1)

$$\sum_{\nu \in \mathbb{Z}} |2^j \langle f, \tilde{h}_{j,\nu} \rangle| \leq \liminf_{\ell \rightarrow \infty} \sum_{\nu \in \mathbb{Z}} |2^j \langle f * \Phi_\ell, \tilde{h}_{j,\nu} \rangle| \lesssim \liminf_{\ell \rightarrow \infty} \|\Phi_\ell * f\|_{W_1^1} \lesssim \|f\|_{BV},$$

where the implicit constants are independent of  $j$ . This yields (5.2).

*Step 2.* We show that, for  $1 \leq p \leq \infty$ , we have

$$\|f\|_p \lesssim \sup_{j \geq -1} 2^{j(1-1/p)} \left( \sum_{\nu \in \mathbb{Z}} |2^j \langle f, \tilde{h}_{j,\nu} \rangle|^p \right)^{1/p} =: A(p) = A.$$

Recall from Lemma 3.1.b that  $\mathbb{E}_N f \rightarrow f$  in  $\mathcal{S}'$ . Assuming (as we may) that  $f \in \mathcal{B}$  has finite right hand side it suffices to show that  $\mathbb{E}_N f$  converges in  $L_p$  and  $\sup_{N \geq 0} \|\mathbb{E}_N f\|_p \lesssim A$ . To see this, one expands

$$\mathbb{E}_N f = \mathbb{E}_0 f + \sum_{0 \leq j < N} \sum_{\mu \in \mathbb{Z}} 2^j \langle f, h_{j,\mu} \rangle h_{j,\mu},$$

and notes that

$$\left\| \sum_{\mu \in \mathbb{Z}} 2^j \langle f, h_{j,\mu} \rangle h_{j,\mu} \right\|_p = 2^{-j/p} \left( \sum_{\mu \in \mathbb{Z}} |2^j \langle f, h_{j,\mu} \rangle|^p \right)^{\frac{1}{p}} \leq 2^{-j} A,$$

hence  $\|\mathbb{E}_{N_1} f - \mathbb{E}_{N_2} f\|_p \lesssim 2^{-N_1} A$ , for all  $N_2 > N_1$ , and thus

$$\|f\|_p \leq \sup_N \|\mathbb{E}_N f\|_p \leq A.$$

*Step 3.* We finally show, for  $1 < p \leq \infty$ , that

$$\|f'\|_p \lesssim \sup_{j \geq -1} 2^{j(1-1/p)} \left( \sum_{\nu \in \mathbb{Z}} |2^j \langle f, \tilde{h}_{j,\nu} \rangle|^p \right)^{1/p} =: A,$$

and when  $p = 1$  then  $f'$  is a finite Borel measure and  $\|f'\|_{\mathcal{M}} \lesssim A$ . Consider the multiresolution analysis in  $L_2(\mathbb{R})$  generated by the subspaces

$$V_N = \overline{\text{span}} \left\{ \mathcal{N}_{N,\mu} := \mathcal{N}_2(2^{N+1} \cdot -\mu) : \mu \in \mathbb{Z} \right\}, \quad N = -1, 0, 1, 2, \dots$$

That is,  $V_N$  consists of continuous piecewise linear functions with nodes in  $2^{-N-1}\mathbb{Z}$ . Let  $\mathcal{N}^*(\cdot)$  be the (polygonal) function which generates the dual Riesz basis to  $\{\mathcal{N}_{N,\mu} : \mu \in \mathbb{Z}\}$ ; see e.g. [7, §3]. Then, the operator

$$h \in L_2 \mapsto P_N(h) := \sum_{\mu \in \mathbb{Z}} 2^N \langle h, \mathcal{N}_{N,\mu}^* \rangle \mathcal{N}_{N,\mu}$$

is the orthogonal projection onto  $V_N$ . Let

$$g_N(x) := \sum_{\mu \in \mathbb{Z}} 2^N \langle f', \mathcal{N}_{N,\mu} \rangle \mathcal{N}_{N,\mu}^*(x).$$

Using Lemma 2.1,<sup>3</sup> we have the uniform bound

$$2^{-\frac{N}{p}} \left( \sum_{\mu \in \mathbb{Z}} |2^N \langle f', \mathcal{N}_{N,\mu} \rangle|^p \right)^{\frac{1}{p}} \lesssim 2^{N(1-\frac{1}{p})} \left( \sum_{\mu \in \mathbb{Z}} |2^N \langle f, \tilde{h}_{N,\mu} \rangle|^p \right)^{\frac{1}{p}} \leq A < \infty.$$

So, the exponential decay of  $\mathcal{N}^*(\cdot)$  guarantees that the series defining  $g_N(x)$  converges, and moreover

$$(5.4) \quad \|g_N\|_p \lesssim 2^{-N/p} \left( \sum_{\mu \in \mathbb{Z}} |2^N \langle f', \mathcal{N}_{N,\mu} \rangle|^p \right)^{1/p} \leq A.$$

<sup>3</sup>Note that, in view of Proposition 1.1, one can give a meaning to the identity  $\langle f', \mathcal{N}_{N,\mu} \rangle = -2^{N+1} \langle f, \tilde{h}_{N,\mu} \rangle$  in Lemma 2.1 also for distributions  $f \in \mathcal{B}$ .

Then, if  $p > 1$  there exists  $g \in L_p$  which is the weak \*-limit of a subsequence of  $\{g_N\}$ . Now, if  $j, \nu$  are fixed, for all  $N \geq j$  we have

$$(5.5) \quad \langle g_N, \psi_{j,\nu} \rangle = \left\langle f', \sum_{\mu \in \mathbb{Z}} 2^N \langle \psi_{j,\nu}, \mathcal{N}_{N,\mu}^* \mathcal{N}_{N,\mu} \rangle \right\rangle = \langle f', \psi_{j,\nu} \rangle,$$

because  $P_N(\psi_{j,\nu}) = \psi_{j,\nu}$ . Thus, taking limits as  $N \rightarrow \infty$  in (5.5) we obtain

$$\langle g, \psi_{j,\nu} \rangle = \langle f', \psi_{j,\nu} \rangle, \quad \text{for all } j, \nu.$$

This implies that  $g = f' \in L_p$  and  $\|f'\|_p \lesssim A$ .

When  $p = 1$  the weak\* sequential-compactness argument only provides that  $\|g\|_{\mathcal{M}} \lesssim A$  and then  $\|f'\|_{\mathcal{M}} \lesssim A$ .  $\square$

## 6. EMBEDDINGS INTO $B_{p,\infty}^{s,\text{dyad}}$ : THE CASES $s = 1$ AND $s = 1/p - 1$

In this section we prove the sufficiency of the conditions for the embeddings into  $B_{p,\infty}^{1,\text{dyad}}$  or  $F_{p,\infty}^{1,\text{dyad}}$  in Theorem 1.7 and the sufficiency for the conditions of embedding into  $B_{p,\infty}^{1/p-1,\text{dyad}}$  in Theorem 1.9.

**Lemma 6.1.** *Let  $1/2 \leq p \leq \infty$ ,  $\frac{1}{p} - 1 \leq s \leq 1$ . Then  $B_{p,\min\{p,1\}}^s \hookrightarrow B_{p,\infty}^{s,\text{dyad}}$  is a continuous embedding.*

*Proof.* Using the notation in the proof of part (ii) of Proposition 4.1 we can write

$$\|f\|_{B_{p,\infty}^{s,\text{dyad}}} = \sup_{j \geq -1} 2^{js} B_j$$

where  $B_j$  is defined as in (4.11) with  $\delta = 0$ . Letting  $u = \min\{p, 1\}$ , we obtain for each  $j$ , arguing as in relation (4.12)

$$\begin{aligned} 2^{js} B_j &\lesssim \left( \sum_{\ell \in \mathbb{Z}} \left[ a(\ell, \frac{1}{p}) 2^{j\ell} \|\mathfrak{M}_{j+\ell}(f_{j+\ell})\|_p \right]^u \right)^{1/u} \\ &\lesssim \left( \sup_{\ell \in \mathbb{Z}} a(\ell, \frac{1}{p}) 2^{-\ell s} \right) \|f\|_{B_{p,u}^s}, \end{aligned}$$

where  $a(\ell, 1/p) = 2^{(1/p-1)\ell}$  for  $\ell > 0$  and  $a(\ell, 1/p) = 2^\ell$  for  $\ell < 0$ . Since  $\sup_{\ell \in \mathbb{Z}} a(\ell, 1/p) 2^{-s\ell} < \infty$  whenever  $1/p - 1 \leq s \leq 1$ , we obtain the desired inequality  $\|f\|_{B_{p,\infty}^{s,\text{dyad}}} \lesssim \|f\|_{B_{p,\min\{p,1\}}^s}$ .  $\square$

**Proposition 6.2.** *Let  $1/2 < p < \infty$ . Then  $F_{p,2}^1 \hookrightarrow F_{p,\infty}^{1,\text{dyad}}$  is a continuous embedding.*

*Proof.* Let  $f \in F_{p,2}^1$ . we must show that

$$\|f\|_{F_{p,\infty}^{1,\text{dyad}}} = \left\| \sup_{j \geq -1} 2^j \sum_{\nu \in \mathbb{Z}} |2^j \langle f, h_{j,\nu} \rangle| \mathbb{1}_{I_{j,\nu}} \right\|_p \lesssim \|f\|_{F_{p,2}^1}.$$

With the notation from Section 2.1, we write  $f = \sum_{k \geq 0} L_k f_k$ , and for each  $j \geq -1$  we split  $f = \Pi_j f + \Pi_j^\perp f$  where

$$\Pi_j f := \sum_{k=0}^j L_k f_k, \quad \text{and} \quad \Pi_j^\perp f := \sum_{k>j} L_k f_k.$$

If  $j = -1$ , we understand that  $\Pi_{-1} f = 0$  and  $\Pi_{-1}^\perp f = f$ . We shall first bound

$$A_0 := \left\| \sup_{j \geq 0} 2^j \sum_{\nu \in \mathbb{Z}} |2^j \langle \Pi_j f, h_{j,\nu} \rangle| \mathbb{1}_{I_{j,\nu}} \right\|_p.$$

The same argument in the proof of [12, Lemma 3.3] gives

$$|2^j \langle \Pi_j f, h_{j,\nu} \rangle| \lesssim \mathfrak{M}_{j,A}^{**} (2^{-j} (\Pi_j f)')(x), \quad x \in I_{j,\nu}.$$

Thus,

$$2^j \sum_{\nu \in \mathbb{Z}} |2^j \langle \Pi_j f, h_{j,\nu} \rangle| \mathbb{1}_{I_{j,\nu}}(x) \lesssim \mathfrak{M}_{j,A}^{**} ((\Pi_j f)')(x), \quad x \in \mathbb{R},$$

and taking a supremum over all  $j \geq 0$ , and then  $L_p$ -norms, we obtain

$$(6.1) \quad A_0 \lesssim \left\| \sup_{j \geq 0} \mathfrak{M}_{j,A}^{**} ((\Pi_j f)') \right\|_p \lesssim \left\| \sup_{j \geq 0} |(\Pi_j f)'| \right\|_p,$$

using (2.10) in the last step with  $A > 1/p$ . Now, the maximal function characterization of the  $h^p = F_{p,2}^0$  norms yields

$$(6.2) \quad \left\| \sup_{j \geq 0} |(\Pi_j f)'| \right\|_p = \left\| \sup_{j \geq 0} |\Pi_j(f')| \right\|_p \lesssim \|f'\|_{F_{p,2}^0} \lesssim \|f\|_{F_{p,2}^1}.$$

To estimate the remaining part involving  $\Pi_j^\perp f$ , we may quote the standard proof in Proposition 4.1<sup>4</sup> above, which gives

$$(6.3) \quad A_1 := \left\| \sup_{j \geq -1} 2^j \sum_{\nu \in \mathbb{Z}} |2^j \langle \Pi_j^\perp f, h_{j,\nu} \rangle| \mathbb{1}_{I_{j,\nu}}(x) \right\|_p \lesssim \|f\|_{F_{p,\infty}^1},$$

provided that  $s > \max\{1/p, 1/q\} - 1$ , with  $s = 1$  and  $q = \infty$ . So, we obtain

$$A_1 \lesssim \|f\|_{F_{p,\infty}^1} \lesssim \|f\|_{F_{p,2}^1},$$

under the assumption that  $p > 1/2$ . Finally, (6.1), (6.2), (6.3) yield the desired estimate  $\|f\|_{F_{p,\infty}^{1,\text{dyad}}} \lesssim \|f\|_{F_{p,2}^1}$ .  $\square$

**Corollary 6.3.** *Let  $1/2 \leq p < \infty$  and  $q \leq \min\{p, 2\}$ . Then  $B_{p,q}^1 \hookrightarrow B_{p,\infty}^{1,\text{dyad}}$  is a continuous embedding.*

*Proof.* For  $q \leq p \leq 1$  this follows from Lemma 6.1. For  $1/2 < p \leq 1$  it follows from Proposition 6.2 together with the inequality

$$(6.4) \quad \|f\|_{B_{p,\infty}^{1,\text{dyad}}} \lesssim \|f\|_{F_{p,\infty}^{1,\text{dyad}}}$$

and the inequality

$$\|f\|_{F_{p,2}^1} \lesssim \|f\|_{B_{p,\min\{p,2\}}^1};$$

<sup>4</sup>That is, the part of the proof of Proposition 4.1 involving the indices  $\ell = k - j \geq 0$ .

the latter being a consequence of Minkowski's inequalities.

Inequality (6.4) in turn follows by definition from the sequence space inequality  $\|\beta\|_{b_{p,\infty}^s} \leq \|\beta\|_{f_{p,\infty}^s}$  (in the case  $s = 1$ ), i.e. from the elementary inequality

$$(6.5) \quad \sup_j \left( \sum_{\mu \in \mathbb{Z}} \int_{I_{j,\mu}} |2^{js} \beta_{j,\mu} \mathbb{1}_{I_{j,\mu}}(x)|^p dx \right)^{1/p} \\ \leq \left( \int \left[ \sup_j 2^{js} \left| \sum_{\mu \in \mathbb{Z}} \beta_{j,\mu} \mathbb{1}_{I_{j,\mu}}(x) \right|^p dx \right]^{1/p} \right)^{1/p}. \quad \square$$

We now consider the other limiting case, where  $s = 1/p - 1$ .

**Proposition 6.4.** *Let  $1/2 < p \leq 1$ ,  $q \leq \infty$ . Then  $F_{p,q}^{\frac{1}{p}-1} \hookrightarrow B_{p,\infty}^{\frac{1}{p}-1, \text{dyad}}$  is a continuous embedding.*

*Proof.* Since  $F_{p,q}^{\frac{1}{p}-1} \hookrightarrow F_{p,\infty}^{\frac{1}{p}-1}$  it suffices to prove this for  $q = \infty$ .

Let  $f \in F_{p,\infty}^{1/p-1}$ , which as before we shall split as

$$f = \Pi_j f + \Pi_j^\perp f.$$

This time, the standard proof in Proposition 4.1 (that is, the part of the proof involving the indices  $\ell = k - j \leq 0$ ) gives

$$\left\| \left( \sum_{j \geq 0} 2^{jsq} \sum_{\nu \in \mathbb{Z}} |2^j \langle \Pi_j f, h_{j,\nu} \rangle|^q \mathbb{1}_{I_{j,\nu}} \right)^{1/q} \right\|_{L_p(\mathbb{R})} \lesssim \|f\|_{F_{p,q}^s}$$

provided that  $s < 1$ . So in particular, letting  $s = 1/p - 1$  and  $q = \infty$  we obtain (after trivial embeddings)

$$(6.6) \quad \sup_{j \geq 0} 2^{j(1/p-1)} \left\| \sum_{\nu \in \mathbb{Z}} 2^j \langle \Pi_j f, h_{j,\nu} \rangle \mathbb{1}_{I_{j,\nu}} \right\|_{L_p(\mathbb{R})} \lesssim \|f\|_{F_{p,\infty}^{\frac{1}{p}-1}},$$

whenever  $p > 1/2$ .

So, it remains to establish a similar estimate with  $\Pi_j^\perp f$  instead of  $\Pi_j f$ . We borrow some notation from [12]. Let  $\mathcal{D}_j$  denote the dyadic intervals of length  $2^{-j}$ . If  $I \in \mathcal{D}_j$  is fixed and  $k \geq j$  we let

- (a)  $\mathcal{D}_k(\partial I) = \{J \in \mathcal{D}_k : \bar{J} \cap \partial I \neq \emptyset\}$
- (b)  $\omega(J) = \{x \in J : \text{dist}(x, \partial I) \geq 2^{-k-1}\}$ , when  $J \in \mathcal{D}_k(\partial I)$ .

We will use the maximal function  $\mathfrak{M}_k^*$  (cf. (2.11)) and note that, as in [12, (42)], when  $J \in \mathcal{D}_k(\partial I)$ ,  $k > j$ , it holds

$$(6.7) \quad \sup_{x \in J} |g(x)| \leq \left[ \int_{\omega(J)} |\mathfrak{M}_k^* g|^p \right]^{\frac{1}{p}}, \quad 0 < p < \infty.$$

Now, let  $I = I_{j,\nu} \in \mathcal{D}_j$  be fixed, and let  $I^\pm$  be its dyadic sons. For each  $k > j$ , the function  $L_k(h_{j,\nu})$  has support contained in the union of the intervals  $J$



belonging to  $\mathcal{D}_k(\partial I^\pm)$ , and  $|L_k(h_{j,\nu})| \lesssim 1$ . Thus, since we are assuming  $p \leq 1$ , we have

$$\begin{aligned} |\langle 2^j h_{j,\nu}, \Pi_j^\perp f \rangle|^p &\leq \sum_{k>j} |\langle L_k(2^j h_{j,\nu}), f_k \rangle|^p \\ &\lesssim \sum_{k>j} 2^{-(k-j)p} \sum_{J \in \mathcal{D}_k(\partial I^\pm)} \|f_k\|_{L^\infty(J)}^p \end{aligned}$$

which, by (6.7), is bounded by

$$\sum_{k>j} 2^{-(k-j)p} \sum_{J \in \mathcal{D}_k(\partial I^\pm)} \int_{\omega(J)} |\mathfrak{M}_k^* f_k(x)|^p \lesssim 2^{jp} \int_{I^*} \sup_{k>j} |2^{(\frac{1}{p}-1)k} \mathfrak{M}_k^* f_k|^p dx,$$

where  $I^*$  is the 2-fold dilation of the interval  $I$ . Summing up in all intervals  $I = I_{j,\nu} \in \mathcal{D}_j$  we obtain

$$\sum_{\nu \in \mathbb{Z}} |\langle 2^j h_{j,\nu}, \Pi_j^\perp f \rangle|^p \lesssim 2^{jp} \int_{\mathbb{R}} \sup_{k>j} |2^{(\frac{1}{p}-1)k} \mathfrak{M}_k^* f_k|^p.$$

This implies

$$(6.8) \quad \sup_j 2^{-j} \left( \sum_{\nu \in \mathbb{Z}} |\langle 2^j h_{j,\nu}, \Pi_j^\perp f \rangle|^p \right)^{1/p} \lesssim \left( \int_{\mathbb{R}} \sup_{k>j} |2^{(\frac{1}{p}-1)k} \mathfrak{M}_k^* f_k|^p \right)^{1/p}.$$

Now,  $\mathfrak{M}_k^* g \lesssim \mathfrak{M}_{k,A}^{**} g$  for any  $A > 0$ , so choosing  $A > 1/p$  and using (2.10) we have

$$(6.9) \quad \left\| \sup_{k \geq 0} 2^{(\frac{1}{p}-1)k} \mathfrak{M}_k^* f_k \right\|_p \lesssim \left\| \sup_{k \geq 0} 2^{(\frac{1}{p}-1)k} |f_k| \right\|_p \lesssim \|f\|_{F_{p,\infty}^{\frac{1}{p}-1}}.$$

Finally, the inequality

$$\|f\|_{B_{p,\infty}^{\frac{1}{p}-1, \text{dyad}}} \lesssim \|f\|_{F_{p,\infty}^{\frac{1}{p}-1}}$$

follows by combining (6.6), (6.8) and (6.9).  $\square$

We now consider the remaining endpoint case where the two limiting cases coincide, that is, we have both  $s = 1$  and  $s = 1/p - 1$ , and thus  $p = 1/2$  (the corresponding Besov embedding is already covered in Lemma 6.1).

**Proposition 6.5.** *For  $p = 1/2$  we have the continuous embedding*

$$F_{1/2,2}^1 \hookrightarrow B_{1/2,\infty}^{1, \text{dyad}}.$$

*Proof.* We examine the proof of Propositions 6.4 and 6.2 and note that (6.8) and (6.9) remain valid for  $p = 1/2$ , that is

$$\sup_{j \geq -1} 2^{-j} \left( \sum_{\nu \in \mathbb{Z}} |\langle 2^j h_{j,\nu}, \Pi_j^\perp f \rangle|^{\frac{1}{2}} \right)^2 \lesssim \|f\|_{F_{1/2,\infty}^1}.$$

Similarly, the arguments in (6.1) and (6.2) do not require a restriction on  $p$ , so we also have

$$\left\| \sup_{j \geq 0} 2^j \sum_{\nu \in \mathbb{Z}} |2^j \langle \Pi_j f, h_{j,\nu} \rangle| |h_{j,\nu}| \right\|_{1/2} \lesssim \|f\|_{F_{1/2,2}^1}.$$

Now the proposition follows from the trivial embeddings  $F_{1/2,2}^1 \hookrightarrow F_{1/2,\infty}^1$  and  $f_{1/2,\infty}^1 \hookrightarrow b_{1/2,\infty}^1$  (cf. (6.5)).  $\square$

## 7. NORM EQUIVALENCES ON SUITABLE SUBSPACES: THE PROOFS OF THEOREMS 1.6 AND 1.8

7.1. *A bootstrapping lemma.* Consider a sequence  $\mathbf{a} = \{a_{j,\mu}\}$  indexed by  $j \in \mathbb{N} \cup \{0\}$  and  $\mu \in \mathbb{Z}$ . As in (1.19) let  $b_{p,q}^s$  be the set of all  $\mathbf{a}$  for which

$$(7.1) \quad \|\mathbf{a}\|_{b_{p,q}^s} = \left( \sum_{j=0}^{\infty} 2^{j(s-\frac{1}{p})q} \left[ \sum_{\mu \in \mathbb{Z}} |a_{j,\mu}|^p \right]^{\frac{q}{p}} \right)^{1/q} < \infty.$$

We split each sequence as  $\mathbf{a} = \mathbf{a}^{\text{even}} + \mathbf{a}^{\text{odd}}$ , where

$$\mathbf{a}_{j,\mu}^{\text{even}} = \begin{cases} a_{j,\mu} & \text{if } \mu \text{ is even} \\ 0 & \text{if } \mu \text{ is odd,} \end{cases} \quad \text{and} \quad \mathbf{a}_{j,\mu}^{\text{odd}} = \begin{cases} 0 & \text{if } \mu \text{ is even} \\ a_{j,\mu} & \text{if } \mu \text{ is odd.} \end{cases}$$

Then

$$\begin{aligned} \|\mathbf{a}^{\text{even}}\|_{b_{p,q}^s} &= \left( \sum_{j=0}^{\infty} 2^{j(s-\frac{1}{p})q} \left[ \sum_{\nu \in \mathbb{Z}} |a_{j,2\nu}|^p \right]^{\frac{q}{p}} \right)^{1/q} \\ \|\mathbf{a}^{\text{odd}}\|_{b_{p,q}^s} &= \left( \sum_{j=0}^{\infty} 2^{j(s-\frac{1}{p})q} \left[ \sum_{\nu \in \mathbb{Z}} |a_{j,2\nu+1}|^p \right]^{\frac{q}{p}} \right)^{1/q} \end{aligned}$$

The key result is the following lemma, which, under suitable conditions, allows us to control  $\|\mathbf{a}\|_{b_{p,q}^s}$  in terms of  $\|\mathbf{a}^{\text{even}}\|_{b_{p,q}^s}$ . The general hypothesis in (7.3) will be linked later to a refinement condition which will appear in (7.6).

**Lemma 7.1.** *Let  $0 < p, q \leq \infty$ ,  $s \in \mathbb{R}$  and  $\vec{\lambda} = (\lambda_0, \lambda_1, \lambda_2) \in \mathbb{C}^3$  such that*

$$(7.2) \quad |\lambda_1| < 2^{s-\frac{1}{p}}.$$

*Then, there exists  $C = C(p, q, s, \vec{\lambda}) > 0$  such that for every sequence  $\mathbf{a} \in b_{p,q}^s$  satisfying the condition*

$$(7.3) \quad |a_{j,2\nu+1}| \leq \sum_{\ell=0}^2 |\lambda_\ell| |a_{j+1,4\nu+2+\ell}|, \text{ for all } j \geq 0, \nu \in \mathbb{Z}.$$

*we have*

$$(7.4) \quad \|\mathbf{a}\|_{b_{p,q}^s} \leq C \|\mathbf{a}^{\text{even}}\|_{b_{p,q}^s}.$$

*Proof.* Let  $\rho = \min\{1, p, q\}$ , and for simplicity write  $\sigma = s - 1/p$ . Condition (7.3) together with the  $\rho$ -triangle inequality gives

$$\begin{aligned} \|\mathbf{a}^{\text{odd}}\|_{b_{p,q}^s}^\rho &\leq \sum_{\ell=0,1,2} |\lambda_\ell|^\rho \left( \sum_{j=0}^{\infty} 2^{j\sigma q} \left[ \sum_{\nu \in \mathbb{Z}} |a_{j+1,4\nu+2+\ell}|^p \right]^{q/p} \right)^{\rho/q} \\ &= (|\lambda_1|2^{-\sigma})^\rho \left( \sum_{j=0}^{\infty} 2^{(j+1)\sigma q} \left[ \sum_{\nu \in \mathbb{Z}} |a_{j+1,4\nu+3}|^p \right]^{q/p} \right)^{\rho/q} \\ &\quad + \sum_{\ell=0,2} (|\lambda_\ell|2^{-\sigma})^\rho \left( \sum_{j=0}^{\infty} 2^{(j+1)\sigma q} \left[ \sum_{\nu \in \mathbb{Z}} |a_{j+1,4\nu+2+\ell}|^p \right]^{q/p} \right)^{\rho/q}. \end{aligned}$$

Using the assumptions  $|\lambda_1| < 2^\sigma$  and  $\mathbf{a} \in b_{p,q}^s$ , the previous display implies

$$\left(1 - \frac{|\lambda_1|^\rho}{2^{\sigma\rho}}\right) \|\mathbf{a}^{\text{odd}}\|_{b_{p,q}^s}^\rho \leq (|\lambda_0|^\rho + |\lambda_2|^\rho) 2^{-\sigma\rho} \|\mathbf{a}^{\text{even}}\|_{b_{p,q}^s}^\rho.$$

This gives

$$\begin{aligned} \|\mathbf{a}\|_{b_{p,q}^s} &\leq \left( \|\mathbf{a}^{\text{odd}}\|_{b_{p,q}^s}^\rho + \|\mathbf{a}^{\text{even}}\|_{b_{p,q}^s}^\rho \right)^{1/\rho} \\ &\leq \left( \frac{|\lambda_0|^\rho + |\lambda_2|^\rho}{2^{\sigma\rho} - |\lambda_1|^\rho} + 1 \right)^{1/\rho} \|\mathbf{a}^{\text{even}}\|_{b_{p,q}^s}, \end{aligned}$$

which finishes the proof.  $\square$

7.2. *Proof of Theorem 1.6.* We must show (1.24), that is

$$(7.5) \quad \|f\|_{B_{p,q}^s} \lesssim \|f\|_{B_{p,q}^{s,\text{dyad}}}, \quad \text{provided } f \in B_{p,q}^s,$$

which, as we shall see, holds actually in the larger range

$$\frac{1}{p} < s < 1 + \frac{1}{p}.$$

By part (ii) of Proposition 4.9, we know that

$$\|f\|_{B_{p,q}^s} \lesssim \|\{\mathbf{c}_{j,\mu}(f)\}_{j \geq 0, \mu \in \mathbb{Z}}\|_{b_{p,q}^s} + \|\{\langle f, h_{-1,\mu} \rangle\}_{\mu \in \mathbb{Z}}\|_{\ell_p} = A + B.$$

Clearly,  $B$  is bounded by the right hand side of (7.5), so we focus on  $A$ .

Define a sequence  $\mathbf{a} = \mathbf{a}(f) = \{a_{j,\nu}\}$  by

$$a_{j,\nu} = 2^j |\langle f, \tilde{h}_{j,\nu} \rangle|, \quad j \in \mathbb{N}_0, \nu \in \mathbb{Z}.$$

Observe from the definition of the coefficients  $\mathbf{c}_{j,\mu}(f)$  in (1.15) that

$$A \lesssim \|\mathbf{a}\|_{b_{p,q}^s}.$$

Note also from (1.8) that  $a_{j,2\mu} = 2^j |\langle f, h_{j,\mu} \rangle|$ , so in particular

$$\|\mathbf{a}^{\text{even}}\|_{b_{p,q}^s} = \|\{2^j \langle f, h_{j,\mu} \rangle\}_{j \geq 0, \mu \in \mathbb{Z}}\|_{b_{p,q}^s} \leq \|f\|_{B_{p,q}^{s,\text{dyad}}}.$$

Therefore, we have reduced matters to prove that

$$\|\mathbf{a}\|_{b_{p,q}^s} \lesssim \|\mathbf{a}^{\text{even}}\|_{b_{p,q}^s}.$$

We shall do so using Lemma 7.1, so we need to verify the hypothesis (7.3), for suitable scalars  $(\lambda_0, \lambda_1, \lambda_2)$ . This will follow from an elementary property of the spline functions

$$\mathcal{N}_{j,\mu}(x) := \mathcal{N}_{2^j j, \mu}(x) = \mathcal{N}_2(2^j x - \mu),$$

defined in §2.3. Recall that these are piecewise linear functions supported in the intervals  $[2^{-j}\mu, 2^{-j}(\mu + 2)]$ . It is then straightforward to verify that

$$(7.6) \quad \mathcal{N}_{j,\mu}(x) = \frac{1}{2} \mathcal{N}_{j+1,2\mu}(x) + \mathcal{N}_{j+1,2\mu+1}(x) + \frac{1}{2} \mathcal{N}_{j+1,2\mu+2}(x);$$

see Figure 6. We refer to (7.6) as the *refinement identity*.




FIGURE 6. Refinement equation for  $\mathcal{N}_{j,\nu}(x)$ ; see (7.6).

Now, if  $\mu = 2\nu + 1$  is an odd integer, then the integration by parts formula and the refinement identity give

$$\begin{aligned} a_{j,\mu} &= 2^j |\langle f, \tilde{h}_{j,\mu} \rangle| = \frac{1}{2} |\langle f', \mathcal{N}_{j+1,\mu} \rangle| \\ \text{by (7.6)} &\leq \frac{1}{4} |\langle f', \mathcal{N}_{j+2,2\mu} \rangle| + \frac{1}{2} |\langle f', \mathcal{N}_{j+2,2\mu+1} \rangle| + \frac{1}{4} |\langle f', \mathcal{N}_{j+2,2\mu+2} \rangle| \\ \text{by (2.15)} &= \frac{1}{2} a_{j+1,2\mu} + a_{j+1,2\mu+1} + \frac{1}{2} a_{j+1,2\mu+2}, \end{aligned}$$

which coincides with (7.3) with  $(\lambda_0, \lambda_1, \lambda_2) = (1/2, 1, 1/2)$ . So we can apply Lemma 7.1, under the assumption

$$|\lambda_1| = 1 < 2^{s-\frac{1}{p}},$$

which holds precisely when  $s > \frac{1}{p}$ . This completes the proof.  $\square$

7.3. *Proof of Theorem 1.8.* We first show that, if  $1 < p \leq \infty$  and  $f \in W_p^1(\mathbb{R})$ , then

$$\|f\|_{W_p^1} \approx \sup_{j \geq -1} 2^{j(1-1/p)} \left( \sum_{\mu \in \mathbb{Z}} |2^j \langle f, h_{j,\mu} \rangle|^p \right)^{1/p}.$$

In view of Theorem 1.4, this reduces to prove

$$(7.7) \quad \begin{aligned} A &:= \sup_{j \geq -1} 2^{j(1-1/p)} \left( \sum_{\mu \in \mathbb{Z}} |\mathfrak{c}_{j,\mu}(f)|^p \right)^{1/p} \\ &\lesssim \sup_{j \geq -1} 2^{j(1-1/p)} \left( \sum_{\mu \in \mathbb{Z}} |2^j \langle f, h_{j,\mu} \rangle|^p \right)^{1/p}, \end{aligned}$$

whenever  $A$  is finite. The argument is completely analogous to that for proving Theorem 1.6, this time using the spaces  $b_{p,\infty}^1$ . Note, that this argument only needs that  $1 = s > 1/p$ .

Finally, the assertion in (1.31), for  $f \in W_p^1$ , follows now easily from

$$\|f\|_{W_p^1} \lesssim \|f\|_{B_{p,\infty}^{1,\text{dyad}}} \lesssim \|f\|_{F_{p,\infty}^{1,\text{dyad}}} \lesssim \|f\|_{F_{p,2}^1} \equiv \|f\|_{W_p^1},$$

where the third inequality was shown in part (ii) of Theorem 1.7 and for the last three steps we assume  $p < \infty$ .  $\square$

## 8. NECESSARY CONDITION FOR THE EMBEDDINGS FOR $s = 1$

We prove the necessary conditions in Theorem 1.7 for various embeddings into  $B_{p,\infty}^{1,\text{dyad}}$ .

**Lemma 8.1.** *Suppose  $1/2 \leq p < \infty$ . Then*

$$(8.1) \quad B_{p,q}^1 \hookrightarrow B_{p,\infty}^{1,\text{dyad}} \implies q \leq p$$

*Proof.* We shall work with an example that has been used in [13, §6.2] to prove lower bounds for the norms of  $\mathbb{E}_N$  on  $B_{p,q}^1$ . Let  $u \in C_c^\infty$  be supported in  $(1/8, 7/8)$  so that  $u(x) = 1$  on  $[1/4, 3/4]$ . For  $N \gg 1$  and  $N/4 \leq j \leq N/2$  define

$$g_{N,j}(x) = u\left(N\left(x - \frac{2j}{N}\right)\right) e^{2\pi i 2^j x}$$

and let  $f_N(x) = \sum_{N/4 \leq j \leq N/2} 2^{-j} g_{N,j}(x)$ . Then by [13, Lemma 29] (Lemma 6.3 in arxiv:1901:09117) we have  $\|f_N\|_{B_{p,q}^1} \lesssim N^{-(1/p-1/q)}$  for  $p \leq q$ . We show that  $\|f_N\|_{B_{p,\infty}^{1,\text{dyad}}} \gtrsim 1$  for large  $N$ , which will imply that  $B_{p,q}^1$  is not continuously embedded into  $B_{p,\infty}^{1,\text{dyad}}$  when  $q > p$ .

To see this we prove lower bounds for many of the Haar coefficients of  $f_N$  at Haar frequency  $2^N$ . Let  $J^{N,j} = (\frac{2j}{N} + \frac{1}{4N}, \frac{2j}{N} + \frac{3}{4N})$ ; we observe that for fixed  $N$  the intervals  $J^{N,j}$  are disjoint and that  $f_N(x) = e^{2^j 2\pi i x}$  for  $x \in J^{N,j}$ . We get by a Taylor expansion

$$(8.2) \quad \langle f_N, h_{N,\mu} \rangle = 2^{-2N-2} f_N'(2^{-N}(\mu + \frac{1}{2})) + R_{N,\mu}$$

with  $|R_{N,\mu}| \leq 2^{-3N} \sup_{I_{j,\mu}} |f_N''|$ . Let  $\mathcal{Z}^{N,j}$  be the set of all integers  $\mu$  such that  $2^{-N}\mu$  and  $2^{-N}(\mu + 1)$  belong to  $J^{N,j}$ ; then for  $\mu \in \mathcal{Z}_{N,j}$

$$|\langle f_N, h_{j,\mu} \rangle| = |\langle 2^{-j} g_{N,j}, h_{j,\mu} \rangle| = 2\pi 2^{-2N-2} + O(2^{j-3N})$$

and hence

$$\|f_N\|_{B_{p,\infty}^{1,\text{dyad}}} \geq 2^{N(1-1/p)} \left( \sum_{\frac{N}{4} < j < \frac{N}{2}} \sum_{\mu \in \mathcal{Z}_{N,j}} |2^N \langle f_N, h_{j,\mu} \rangle|^p \right)^{1/p}.$$

Since  $\#(\mathcal{Z}^{N,j}) \approx 2^N N^{-1}$  for large  $N$  and  $N/4 < j < N/2$  we obtain  $\|f_N\|_{B_{p,\infty}^{1,\text{dyad}}} \gtrsim 1$ .  $\square$

**Lemma 8.2.** *Suppose  $1/2 \leq p < \infty$ . Then*

$$(8.3) \quad F_{p,q}^1 \hookrightarrow B_{p,\infty}^{1,\text{dyad}} \implies q \leq 2,$$

$$(8.4) \quad B_{p,q}^1 \hookrightarrow B_{p,\infty}^{1,\text{dyad}} \implies q \leq 2.$$

*Proof.* We consider the same example that was used in [12, §7.2.1]. Namely, let  $\psi \in C_c^\infty(0,1)$  with  $\psi \equiv 1$  in  $[1/4, 3/4]$ , and for each large  $N \gg 1$ , let

$$(8.5) \quad \mathfrak{J}_N = \{j \in \mathbb{N} : N/4 < j < N/2\}$$

and

$$(8.6) \quad f_t(x) := \sum_{j \in \mathfrak{J}_N} \frac{r_j(t)}{2^j} e^{2\pi i 2^j x} \psi(x), \quad t \in [0,1],$$

where  $r_j(t)$ ,  $t \in [0,1]$ , are the usual Rademacher functions. Using Lemma 7.3 from [12] one can verify that

$$(8.7) \quad \sup_{t \in [0,1]} \|f_t\|_{F_{p,q}^1} \lesssim N^{1/q};$$

a similar argument also gives

$$(8.8) \quad \sup_{t \in [0,1]} \|f_t\|_{B_{p,q}^1} \lesssim N^{1/q}.$$

Let  $\psi_j(x) = 2^{-j} e^{2\pi i 2^j x} \psi(x)$ , for  $j \in \mathfrak{J}_N$ , and let  $\mathcal{Z}_N$  be the set of all  $\mu \in \mathbb{Z}$  such that  $I_{N,\mu} \subset (1/4, 3/4)$ . Using the Taylor expansion as in (8.2) one sees that

$$(8.9) \quad |\langle \psi_j, 2^N h_{N,\mu} \rangle| = 2\pi 2^{-N-2} + O(2^{j-2N}), \quad \mu \in \mathcal{Z}_N.$$

Now observe that

$$(8.10) \quad \|f_t\|_{B_{p,\infty}^{1,\text{dyad}}} \geq 2^{N(1-1/p)} \left( \sum_{\mu \in \mathcal{Z}_N} |\langle f_t, 2^N h_{N,\mu} \rangle|^p \right)^{\frac{1}{p}}.$$

So, raising to the  $p$ -th power and taking the expectation in the  $t$  variable, we obtain from Khintchine's inequality

$$\begin{aligned} \left( \int_0^1 \|f_t\|_{B_{p,\infty}^{1,\text{dyad}}}^p dt \right)^{\frac{1}{p}} &\geq 2^{N(1-\frac{1}{p})} \left( \sum_{\mu \in \mathcal{Z}_N} \int_0^1 \left| \sum_{j \in \mathfrak{J}_N} r_j(t) \langle \psi_j, 2^N h_{N,\mu} \rangle \right|^p dt \right)^{\frac{1}{p}} \\ &\gtrsim 2^{N(1-\frac{1}{p})} \left( \sum_{\mu \in \mathcal{Z}_N} \left[ \sum_{j \in \mathfrak{J}_N} |\langle \psi_j, 2^N h_{N,\mu} \rangle|^2 \right]^{\frac{p}{2}} \right)^{\frac{1}{p}}. \end{aligned}$$

An application of (8.9), together with the cardinalities of  $\mathfrak{Z}_N$  and  $\mathcal{Z}_N$ , then gives

$$\left( \int_0^1 \|f_t\|_{B_{p,\infty}^{1,\text{dyad}}}^p dt \right)^{\frac{1}{p}} \gtrsim \sqrt{N}.$$

This together with (8.7), (8.8) implies that the inclusions  $B_{p,q}^1 \hookrightarrow B_{p,\infty}^{1,\text{dyad}}$  and  $F_{p,q}^1 \hookrightarrow B_{p,\infty}^{1,\text{dyad}}$  can only hold if  $q \leq 2$ .  $\square$

**Lemma 8.3.** *For  $p = \infty$  we have*

$$(8.11) \quad B_{\infty,q}^1 \hookrightarrow B_{\infty,\infty}^{1,\text{dyad}} \implies q \leq 1.$$

*Proof.* We assume that  $B_{\infty,q}^1 \hookrightarrow B_{\infty,\infty}^{1,\text{dyad}}$ , and we shall prove that necessarily  $q \leq 1$ . Let  $\mathcal{Z}_N$  be as in (8.5) and consider the function

$$f(x) := \sum_{j \in \mathfrak{Z}_N} 2^{-j} e^{2\pi i 2^j x} \psi(x),$$

which is defined as in (8.6), but with all the  $r_j(t)$  set equal to 1. This time we shall assume that  $\psi \in C_c^\infty(-1/2, 1/2)$  with  $\psi = 1$  in  $(-1/4, 1/4)$ . As in (8.8) we have

$$(8.12) \quad \|f\|_{B_{p,q}^1} \lesssim N^{1/q}.$$

On the other hand, note that

$$(8.13) \quad \|f\|_{B_{\infty,\infty}^{1,\text{dyad}}} \geq 2^N |\langle f, 2^N h_{N,0} \rangle|.$$

Arguing as in (8.2) we see that

$$\begin{aligned} \langle f, 2^N h_{N,0} \rangle &= \sum_{j \in \mathfrak{Z}_N} \langle \psi_j, 2^N h_{N,0} \rangle = \sum_{j \in \mathfrak{Z}_N} [2^{-j} 2^{-N-2} \psi_j'(0) + O(2^{j-2N})] \\ &= 2\pi i 2^{-N-2} \text{Card}(\mathfrak{Z}_N) + O(2^{-3N/2}), \end{aligned}$$

which inserted into (8.13) gives

$$(8.14) \quad \|f\|_{B_{\infty,\infty}^{1,\text{dyad}}} \gtrsim N.$$

The lemma is proved after combining (8.12), (8.14) and letting  $N \rightarrow \infty$ .  $\square$

## 9. NECESSARY CONDITION FOR EMBEDDING INTO $B_{p,\infty}^{1/p-1,\text{dyad}}$

**Proposition 9.1.** *Let  $0 < p, q \leq \infty$ . Then*

$$B_{p,q}^{1/p-1} \hookrightarrow B_{p,\infty}^{1/p-1,\text{dyad}} \implies q \leq \min\{1, p\}.$$

*Proof.* We first assume  $1 < p \leq \infty$ . Suppose the embedding  $B_{p,q}^s \hookrightarrow B_{p,\infty}^{s,\text{dyad}}$  holds, with  $s = 1/p - 1$ . By definition of the latter space we have the inequality

$$(9.1) \quad |\langle f, h_{j,\mu} \rangle| \lesssim 2^{j(\frac{1}{p}-1-s)} \|f\|_{B_{p,\infty}^{s,\text{dyad}}},$$

so the assumed embedding would then imply that  $h_{j,\mu}$  defines a bounded linear functional on  $B_{p,q}^{-1+1/p}$  (or in the subspace  $\mathring{B}_{p,q}^{-1+1/p}$  defined by the closure of  $\mathcal{S}$  in the  $B_{p,q}^{-1+1/p}$  norm, in case that  $p$  or  $q$  are  $\infty$ ). By the duality identities of Besov spaces, see [25, §2.11], this means that  $h_{j,\mu} \in B_{p',q'}^{1/p'}$  which cannot be the case if  $q' < \infty$ , i.e. if  $1 < q \leq \infty$ .

Let  $p \leq 1$ . We use an example from [13, §10.1]. We let  $\eta_l(x) = 2^l \eta(2^l x)$  where  $\eta \in C_c^\infty(\mathbb{R})$  is an odd function supported in  $(-1/2, 1/2)$  such that  $\int_0^{1/2} \eta(s) ds = 1$  and such that  $\int_0^{1/2} \eta(s) s^n ds = 0$  for  $n = 1, 2, \dots, M$ , for a sufficiently large integer  $M$ . Let

$$f_N(x) = \sum_{m=1}^{\infty} a_m \eta_{N+m}(x - 2^{-N+5}m).$$

By [13, (85)] we have

$$\|f_N\|_{B_{p,q}^{1/p-1}} \lesssim \left( \sum_{m=1}^{\infty} |a_m|^q \right)^{1/q}.$$

On the other hand a calculation shows  $\langle f_N, h_{N,2^5 m} \rangle = a_m$  and thus

$$\|f_N\|_{B_{p,\infty}^{1/p-1, \text{dyad}}} \geq \left( \sum_{\mu} |\langle f_N, h_{N,\mu} \rangle|^p \right)^{1/p} \gtrsim \left( \sum_{m=1}^{\infty} |a_m|^p \right)^{1/p}$$

which forces  $q \leq p$ . □

## 10. $B_{p,q}^{1/p}$ AND $B_{p,q}^{1/p, \text{dyad}}$ : THE PROOF OF THEOREM 1.10

10.1. *Proof of part (i) of Theorem 1.10.* Let

$$(10.1) \quad f_N = \sum_{j=0}^{N-1} h_{j,0}, \quad \text{for } N = 1, 2, \dots$$

Observe that

$$(10.2) \quad \|f_N\|_{B_{p,\infty}^{s, \text{dyad}}} = \sup_{j \geq 0} 2^{j(s-\frac{1}{p})} \left[ \sum_{\mu \in \mathbb{Z}} |2^j \langle f_N, h_{j,\mu} \rangle|^p \right]^{1/p} = 1, \quad \text{if } s = 1/p.$$

However, using the characterization with differences of order 1 for the  $B_{p,q}^s$ -norm (if  $s \in (0, 1)$ ), see [25, Theorem 2.5.12], we have

$$\begin{aligned} \|f_N\|_{B_{p,\infty}^{1/p}} &\gtrsim 2^{N/p} \|\Delta_{2^{-N}}(f_N)\|_p \\ &\geq 2^{N/p} \left[ \int_{-2^{-N}}^0 \left| \sum_{0 \leq j < N} (h_{j,0}(x + 2^{-N}) - h_{j,0}(x)) \right|^p dx \right]^{\frac{1}{p}}. \end{aligned}$$

Now a simple computation shows that, if  $\delta \in (0, 2^{-j-1}]$ , then

$$h_{j,0}(\cdot + \delta) - h_{j,0}(\cdot) = \mathbb{1}_{[-\delta, 0)} - 2\mathbb{1}_{[2^{-j-1}-\delta, 2^{-j-1})} + \mathbb{1}_{[2^{-j}-\delta, 2^{-j})}$$



and then also

$$\begin{aligned} \Delta_\delta^2 h_{j,0} &= \mathbb{1}_{[-2\delta, -\delta]} - 2\mathbb{1}_{[2^{-j-1}-2\delta, 2^{-j-1}-\delta]} + \mathbb{1}_{[2^{-j}-2\delta, 2^{-j}-\delta]} \\ &\quad - \mathbb{1}_{[-\delta, 0]} + 2\mathbb{1}_{[2^{-j-1}-\delta, 2^{-j-1}]} - \mathbb{1}_{[2^{-j}-\delta, 2^{-j}]} \end{aligned}$$

and therefore, setting  $\delta = 2^{-N}$  we get

$$\Delta_{2^{-N}}^2 h_{j,0}(x) = 1, \text{ for } x \in [-2^{1-N}, -2^{-N}) \text{ and } j = 0, \dots, N-1.$$

Hence, for  $1 \leq p < \infty$

$$(10.3) \quad \|f_N\|_{B_{p,\infty}^{1/p}} \gtrsim 2^{N/p} \|\Delta_{2^{-N}}^2 f_N\|_p \geq N,$$

and hence part (i) of Theorem 1.10 follows.  $\square$

10.2. *Proof of part (ii) of Theorem 1.10.* Consider this time the function

$$f = \sum_{j=0}^{\infty} h_{j,0},$$

and as before  $f_N = \sum_{j=0}^{N-1} h_{j,0}$ , i.e.  $f = \lim_{N \rightarrow \infty} f_N$  with convergence in  $L_p$ . Indeed

$$(10.4) \quad \|f - f_N\|_p \leq \sum_{j \geq N} \|h_{j,0}\|_p = \sum_{j \geq N} 2^{-j/p} \approx 2^{-N/p}.$$

As in (10.2), it is again easy to verify that

$$(10.5) \quad \|f\|_{B_{p,\infty}^{1/p, \text{dyad}}} = 1.$$

We claim that  $f \notin B_{p,\infty}^{1/p}$ . Indeed, for large  $N$  we have

$$(10.6) \quad \begin{aligned} \|f\|_{B_{p,\infty}^{1/p}} &\gtrsim 2^{N/p} \|\Delta_{2^{-N}}(f - f_N) + \Delta_{2^{-N}} f_N\|_p \\ &\geq 2^{N/p} (\|\Delta_{2^{-N}} f_N\|_p - 2\|f - f_N\|_p). \end{aligned}$$

Inserting the bounds (10.3) and (10.4) into (10.6) gives

$$\|f\|_{B_{p,\infty}^{1/p}} \gtrsim 2^{N/p} \|\Delta_{2^{-N}}(f_N)\|_p - O(1) \gtrsim N,$$

which letting  $N \nearrow \infty$  proves the assertion.  $\square$

## 11. SOME PATHOLOGIES OF THE SPACES $B_{p,q}^{s, \text{dyad}}$

We include in this section some pathologies of the spaces  $B_{p,q}^{s, \text{dyad}}$  when  $s > 1$ , or  $s = 1$ ,  $q < \infty$ , or  $s < 1/p - 1$ , which were mentioned in the introduction.

11.1. *Failure of embedding into  $B_{p,q}^{s,\text{dyad}}$  for  $s > 1$  or  $s = 1$ ,  $q < \infty$ .* The following proposition is a simple result on the theme of Brezis' paper [3] on how to recognize constant functions.

**Proposition 11.1.** *Let  $0 < p, q \leq \infty$  and assume that either (i)  $s > 1$ , or (ii)  $s = 1$  and  $q < \infty$ .*

*Then every  $f \in C^1(\mathbb{R}) \cap B_{p,q}^{s,\text{dyad}}(\mathbb{R})$  is a constant function.*

*Remark.* Bočkarëv's results [1] indicate that less restrictive assumptions can be made but we will not pursue the problem of optimal hypotheses here.

*Proof.* We argue as in the proof of Lemma 8.1, now using Taylor's formula in the form

$$\sup_{b \in K} \sup_{|b-y| \leq \varepsilon} |f(y) - f(b) - f'(b)(y-b)| = o(\varepsilon)$$

for any compact  $K$ . Take  $b \equiv b_\mu = 2^{-j}(\mu + \frac{1}{2})$  to see that

$$(11.1) \quad \langle f, h_{j,\mu} \rangle = f'(2^{-j}(\mu + \frac{1}{2}))2^{-2j-2} + o(2^{-2j})$$

with uniformity in the remainder as  $b_\mu$  ranges over a compact set.

Now assume that  $f \in C^1$  and that  $f'$  is not identically zero. Then there is a dyadic interval  $J = [\nu 2^{-\ell}, (\nu+1)2^{-\ell}]$  and  $c > 0$  such that for  $j \geq j_0 > \ell$

$$|\langle f, h_{j,\mu} \rangle| \geq c 2^{-2j}.$$

Hence

$$\begin{aligned} \|f\|_{B_{p,q}^{s,\text{dyad}}} &\geq \left( \sum_{j=j_0}^{\infty} \left[ 2^{j(s-\frac{1}{p})} \left( \sum_{\mu: I_{j,\mu} \cap J} |2^j \langle f, h_{j,\mu} \rangle|^p \right)^{\frac{1}{p}} \right]^q \right)^{\frac{1}{q}} \\ &\geq \left( \sum_{j=j_0}^{\infty} \left[ 2^{j(s-1/p)} 2^{(j-\ell)/p} c 2^{-j} \right]^q \right)^{\frac{1}{q}} \geq c_\ell \left( \sum_{j=j_0}^{\infty} 2^{j(s-1)q} \right)^{1/q} \end{aligned}$$

with  $c_\ell > 0$ . Hence  $\|f\|_{B_{p,q}^{s,\text{dyad}}} = \infty$  when  $s > 1$  or when  $s = 1$  and  $q < \infty$ .

We conclude that for this range we have  $f' \equiv 0$  for every  $f \in C^1 \cap B_{p,q}^{s,\text{dyad}}$  and Proposition 11.1 follows.  $\square$

11.2. *The dyadic Besov-spaces for  $s < 1/p - 1$ : Failure of completeness.*

**Proposition 11.2.** *Let  $0 < p, q \leq \infty$ . If  $s < 1/p - 1$  then the spaces  $B_{p,q}^{s,\text{dyad}}(\mathbb{R})$  are not complete.*

*Proof.* Consider the functions

$$(11.2) \quad f_N = \mathbb{1}_{[0,1)} + \sum_{j=0}^{N-1} 2^j h_{j,0} = 2^N \mathbb{1}_{[0,2^{-N})}, \quad N = 1, 2, \dots$$

It is easily seen that, under the assumption  $s < 1/p - 1$ , then

$$\|f_M - f_N\|_{B_{p,q}^{s,\text{dyad}}} = \left( \sum_{N \leq j < M} [2^{j(s+1-\frac{1}{p})}]^q \right)^{\frac{1}{q}} \rightarrow 0,$$

when  $M > N \rightarrow \infty$ . So,  $\{f_N\}_{N \geq 1}$  is a Cauchy sequence in  $B_{p,q}^{s,\text{dyad}}$ . However, the distributional limit of  $f_N$  is the Dirac measure  $\delta$ , which does not belong to the space  $\mathcal{B}$ .  $\square$

11.3. *Failure of an embedding for  $s = 1/p - 1$ .* A small variation of the last example shows also part (i) of Theorem 1.9 and at the same time the optimality of the condition  $s > 1/p - 1$  in part (ii) of Proposition 4.1 when  $q < \infty$ .

**Proposition 11.3.** *Let  $0 < p, u \leq \infty$ . Then*

$$B_{p,u}^{1/p-1} \not\hookrightarrow B_{p,q}^{1/p-1,\text{dyad}}, \quad 0 < q < \infty.$$

*Proof.* Consider  $f_N$  as in (11.2), and let  $g_N = f_N - f_N(-\cdot)$  be its odd extension. Then, it was shown in [13, Proposition 52] (Proposition 13.3 in arXiv:1901:09117) that

$$\|g_N\|_{B_{p,u}^{1/p-1}} \lesssim 1,$$

for all  $0 < p, u \leq \infty$ . However, it is easily seen that

$$\|g_N\|_{B_{p,q}^{1/p-1,\text{dyad}}} \geq \|f_N\|_{B_{p,q}^{1/p-1,\text{dyad}}} \geq N^{1/q}. \quad \square$$

## REFERENCES

- [1] S. V. Bočkarëv. The coefficients of Fourier series in Haar's system. *Mat. Sb. (N.S.)*, 80 (122):97–116, 1969.
- [2] S. V. Bočkarëv. A method of averaging in the theory of orthogonal series and some problems in the theory of bases. *Proc. Steklov Inst. Math.*, (3):vi+92, 1980. Translated by H. H. McFaden, A translation of Trudy Mat. Inst. Steklov. 146.
- [3] H. Brezis. How to recognize constant functions. A connection with Sobolev spaces. *Uspekhi Mat. Nauk*, 57(4(346)):59–74, 2002.
- [4] H. Brezis, A. Seeger, J. Van Schaftingen, and P.-L. Yung. Family of functions representing Sobolev norms. *Analysis and PDE*, to appear. arxiv 2109.02903.
- [5] H. Brezis, J. Van Schaftingen, and P.-L. Yung. A surprising formula for Sobolev norms. *Proc. Natl. Acad. Sci. USA*, 118(8):Paper No. e2025254118, 6, 2021.
- [6] C. K. Chui. *An introduction to wavelets*, volume 1 of *Wavelet Analysis and its Applications*. Academic Press, Inc., Boston, MA, 1992.
- [7] C. K. Chui and J.-Z. Wang. On compactly supported spline wavelets and a duality principle. *Trans. Amer. Math. Soc.*, 330(2):903–915, 1992.
- [8] N. Derevianko and T. Ullrich. A higher order Faber spline basis for sampling discretization of functions. *J. Approx. Theory*, 257:105449, 38, 2020.
- [9] M. Frazier and B. Jawerth. A discrete transform and decompositions of distribution spaces. *J. Funct. Anal.*, 93(1):34–170, 1990.
- [10] G. Garrigós, A. Seeger, and T. Ullrich. On uniform boundedness of dyadic averaging operators in spaces of Hardy-Sobolev type. *Anal. Math.*, 43(2):267–278, 2017.
- [11] G. Garrigós, A. Seeger, and T. Ullrich. The Haar system as a Schauder basis in spaces of Hardy-Sobolev type. *J. Fourier Anal. Appl.*, 24(5):1319–1339, 2018.

- [12] G. Garrigós, A. Seeger, and T. Ullrich. The Haar system in Triebel-Lizorkin spaces: endpoint results. *J. Geom. Anal.*, 31(9):9045–9089, 2021.
- [13] G. Garrigós, A. Seeger, and T. Ullrich. Basis properties of the Haar system in limiting Besov spaces. In *Geometric aspects of harmonic analysis*, volume 45 of *Springer INdAM Ser.*, pages 361–424. Springer, Cham, [2021] ©2021. ArXiv version in arXiv:1901.09117.
- [14] B. I. Golubov. On Fourier series of continuous functions with respect to a Haar system. *Izv. Akad. Nauk SSSR Ser. Mat.*, 28:1271–1296, 1964.
- [15] K. Gröchenig. Describing functions: atomic decompositions versus frames. *Monatsh. Math.*, 112(1):1–42, 1991.
- [16] S. Jaffard and H. Krim. The unreasonable effectiveness of Haar frames, Preprint 2021.
- [17] G. Kyriazis. Decomposition systems for function spaces. *Studia Math.*, 157(2):133–169, 2003.
- [18] J. R. McLaughlin. Haar series. *Trans. Amer. Math. Soc.*, 137:153–176, 1969.
- [19] I. Novikov and E. Semenov. *Haar series and linear operators*, volume 367 of *Mathematics and its Applications*. Kluwer Academic Publishers Group, Dordrecht, 1997.
- [20] J. Peetre. On spaces of Triebel-Lizorkin type. *Ark. Mat.*, 13:123–130, 1975.
- [21] A. Seeger and T. Ullrich. Haar projection numbers and failure of unconditional convergence in Sobolev spaces. *Math. Z.*, 285(1-2):91–119, 2017.
- [22] A. Seeger and T. Ullrich. Lower bounds for Haar projections: deterministic examples. *Constr. Approx.*, 46(2):227–242, 2017.
- [23] R. Srivastava. Orthogonal systems of spline wavelets as unconditional bases in Sobolev spaces, Preprint 2020. To appear in *Math. Nachr.* arxiv:2002:09980.
- [24] E. M. Stein. *Singular integrals and differentiability properties of functions*. Princeton Mathematical Series, No. 30. Princeton University Press, Princeton, N.J., 1970.
- [25] H. Triebel. *Theory of function spaces*, volume 38 of *Mathematik und ihre Anwendungen in Physik und Technik [Mathematics and its Applications in Physics and Technology]*. Akademische Verlagsgesellschaft Geest & Portig K.-G., Leipzig, 1983.
- [26] H. Triebel. *Theory of function spaces. II*, volume 84 of *Monographs in Mathematics*. Birkhäuser Verlag, Basel, 1992.
- [27] H. Triebel. *Theory of function spaces. III*, volume 100 of *Monographs in Mathematics*. Birkhäuser Verlag, Basel, 2006.
- [28] H. Triebel. *Bases in function spaces, sampling, discrepancy, numerical integration*, volume 11 of *EMS Tracts in Mathematics*. European Mathematical Society (EMS), Zürich, 2010.

GUSTAVO GARRIGÓS, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MURCIA, 30100 ESPINARDO, MURCIA, SPAIN

*Email address:* `gustavo.garrigos@um.es`

ANDREAS SEEGER, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, 480 LINCOLN DRIVE, MADISON, WI, 53706, USA

*Email address:* `seeger@math.wisc.edu`

TINO ULLRICH, FAKULTÄT FÜR MATHEMATIK, TECHNISCHE UNIVERSITÄT CHEMNITZ, 09107 CHEMNITZ, GERMANY

*Email address:* `tino.ullrich@mathematik.tu-chemnitz.de`