

DISCRETE MOVING FRAMES ON LATTICE VARIETIES AND LATTICE BASED MULTISPACES

GLORIA MARÍ BEFFA AND ELIZABETH L. MANSFIELD

ABSTRACT. In this paper, we develop the theory of the discrete moving frame in two different ways. In the first half of the paper, we consider a discrete moving frame defined on a lattice variety, and the equivalence classes of global syzygies that result from the first fundamental group of the variety. In the second half, we consider the continuum limit of discrete moving frames as a local lattice coalesces to a point. To achieve a well-defined limit of discrete frames, we construct multispace, a generalization of the jet bundle that also generalizes Olver's one dimensional construction. Using interpolation to provide coordinates, we prove that it is a manifold containing the usual jet bundle as a submanifold and show that continuity of a multispace moving frame ensures that the discrete moving frame converges to the continuous one as lattices coalesce. We prove further that the discrete invariants and syzygies approximate their smooth counterparts. In our last chapter we discuss two important applications, one to the discrete variational calculus, and the second to discrete integrable systems. Finally, in an appendix, we discuss a more general result concerning equicontinuous families of discretisations of moving frames, which are consistent with a smooth frame.

CONTENTS

1. Introduction	1
2. Background	4
2.1. Moving frames	4
2.2. Lattices	6
3. Moving frames on Lattices and Lattice Varieties	7
3.1. Moving frames, invariants and Maurer–Cartan invariants	7
3.2. Syzygies	11
4. Continuous limits of the discrete picture: lattice based multispace	20
4.1. Multispace in one variable	21
4.2. Multivariate interpolation	22
4.3. Multispaces in several variables	27
4.4. The continuous limit of invariants and syzygies	38
5. Applications	43
5.1. Application to the design of a variational numerical scheme for a shallow water system	43
5.2. Discretizations of completely integrable PDEs	46
Appendix A. Equicontinuous families of discrete frames	53
References	54

This paper was supported by ELM's EPSRC grant EP/H024018/1, GMB's NSF grant DMS #0804541 and GMB's Simons Fellowship.

1. INTRODUCTION

The theory and the applications of Lie group based moving frames is now well established, and provides an “invariant calculus” to study differential systems which are either invariant or equivariant under the action of a Lie group. Associated with the name of Élie Cartan [10], who used *repères mobile* to solve equivalence problems in differential geometry, the ideas go back to earlier works, for example by Cotton [13] and Darboux [14].

Moving frames were further developed and applied in a substantial body of work, in particular to differential geometry and exterior differential systems; see for example papers by Green [19] and Griffiths [20]. From the point of view of symbolic computation, a breakthrough in the understanding of Cartan’s methods for differential systems came in a series of papers by Fels and Olver [16, 17], Olver [50, 51], Hubert [27, 28, 29], and Hubert and Kogan [30, 31], which provide a coherent, rigorous and constructive moving frame method. The resulting differential invariant calculus is the subject of the textbook, [37], and there have been an extensive number of applications since its publication, including to the integration of Lie group invariant differential equations, to the Calculus of Variations and Noether’s Theorem, (see for example [36, 21, 22]), and to integrable systems (for example [39, 42, 43, 44]).

The first results for the computation of discrete invariants using group-based moving frames were given by Olver who called them joint invariants in [51]; modern applications to date include computer vision [49] and numerical schemes for systems with a Lie symmetry [12, 33, 34, 35, 40]. While moving frames for discrete applications as formulated by Olver do give generating sets of discrete invariants, the recursion formulae for differential invariants which were so successful for the application of moving frames to calculus based results, do not generalize well to joint invariants. In particular, joint invariants do not seem to have recursion formulae under the shift operator that are computationally useful. To overcome this problem, the authors, together with Jing Ping Wang, introduced the notion of a *discrete moving frame* which is essentially a sequence of frames [41]. In that paper we prove discrete recursion formulae for a small computable generating sets of invariants, which we call the *discrete Maurer–Cartan invariants*, and investigated their *syzygies*, that is, their recursion relations. The main application to date has been to discrete integrable systems, with the authors of [46] proving that discrete Hamiltonian structures for W_n -algebras can be obtained via a reduction process. We note that a sequence of moving frames was also used in [34] to minimize the accumulation of errors in an invariant numerical method.

In this paper, we extend the theory of discrete moving frames in two ways. The first is to consider a discrete moving frame defined on a lattice variety, which can be thought of as the vertices, or 0-cells, together with their adjacency information, in a discrete approximation of a manifold. We describe their associated cross-sections and define Maurer-Cartan invariants and local syzygies. In Section 3.2 we further classify global syzygies and prove that they are associated to topological aspects of the variety, like representatives of the discrete fundamental group of the lattice variety, with properties like twisting.

The second extension, beginning in Section 4 and for the rest of the paper, is to consider families of discrete frames and how their continuum limits may define

smooth frames. Our interest in this second case is how discrete invariants and their recursion relations limit to differential invariants and their differential syzygies.

In order to provide a general framework, we construct a manifold which we call the *lattice based multispace* and which generalises, in some sense, the curve based multispace of Olver [50]. The multispace is a generalization of the jet bundle and which contains the jet bundle as a submanifold. It also contains the space of lattices as an open subset. The main problem with the definition of the lattice multispace is the fact that multivariate interpolation is not well-defined in general. To avoid this problem we restrict the lattices to sets of points covered by the general construction of deBoor and Ron ([5], [6], [7]), to what we call *corner lattices*. A corner lattice is one with just enough data to guarantee the approximation of a smooth jet. We restrict as well the types of coalescing that can take place to be those along hyperplanes. We show that deBoor and Ron's interpolating family is well defined on corner lattices and is smooth under coalescing, smooth in the sense that the associated Lagrange polynomials converge to the Hermite ones as the vertices of the lattice coalesce. Once the choice of lattice and conditions on coalescing are settled, we can use the interpolating coefficients to define the coordinate system in the multispace manifold. We notice that one can possibly consider other forms of lattices and coalescing, and that our theory will hold true as far as the smoothness of deBoor-Ron's family is preserved.




FIGURE 1. Under coalescence of the points at which the interpolation is calculated, Lagrange interpolation becomes Hermite interpolation, ending with the Taylor approximation to a surface when all the interpolation points coalesce. By taking coordinates for the lattice based multispace to be the grid points and the Lagrange interpolation coefficients, the jet bundle is naturally embedded.

Once the multispace is proved to be a smooth manifold, we can naturally define a group action on it, and hence we can talk about smooth moving frames on multispaces. A moving frame on the lattice based multispace is, simultaneously, a smooth moving frame defined on the jet bundle, and a frame defined on local difference approximations to the derivatives, depending on what point of the multispace the moving frame is evaluated. By defining a moving frame on multispaces, one

has simultaneously the full power of both the smooth and the discrete frames, and the smoothness of the multispace frame will ensure that we can move freely between discrete, discrete/differential and smooth frames, ensuring that the discrete frame converges to the continuous one as the points in the corner lattices coalesce to create the jet. We also show that the continuity carries over to invariants and syzygies as well. Therefore, any smooth geometric construction carried out with a multispace lattice, invariants and syzygies, ensures that the final discrete, or discrete/differential result is an approximation of the corresponding continuous construction.

In section 5 we use the multispace construction in two different applications. The first application is to a class of finite difference variational shallow water systems, which have both the correct continuum limit as well as the necessary symmetries for Noether's theorem to yield conservation laws for energy, and linear and angular momenta, in both the finite difference case and the smooth limit. This is motivated by the desire to achieve an analogue of the conservation of potential vorticity in a numerical approximation to these equations.

The second application concerns discretizations of completely integrable systems. Most well known completely integrable PDEs are linked to some geometric background and the PDE can be interpreted as, for example, the equation induced on invariants by a geometric evolution of curves, or like the Codazzi–Mainardi equations, are associated to the geometry of some type of surface. Discrete lattice systems also have similar interpretations ([2]). The question to ponder is whether or not the same geometric construction performed in the continuous case to generate the PDEs can be carried out in the discrete case, while guaranteeing that the result will be a discretization of the PDE; this might be useful as a base to study the more interesting questions of when the discretization will be also completely integrable. Here we show two such processes: the Boussinesq equation is induced on centro-affine invariants by an evolution of star-shaped curves. We construct the multispace version of the construction to obtain a geometric discretization. We show that a modification of the construction generates an integrable discretization which appeared in [46]. The study of how these different discretizations might be related is underway. In the second example we describe the multispace version of the well-known construction of the Sine-Gordon equation as the Codazzi–Mainardi equations for Euclidean surfaces of negative constant curvature. This interpretation has been widely used to study pseudo-spherical surfaces as generated by solutions of Sine-Gordon, see [11] and [55]. Every step of the construction is guaranteed to discretize the continuous version, while preserving the geometric meaning of the elements involved. The Sine-Gordon is in fact one of several equations describing the surface, but which decouples from the others. In this discretization the equations remains coupled and its integrability is not clear, but the construction itself is a non-trivial example of the use of mixed discrete-smooth moving frames. The connection between multispace and integrability is under study.

Finally, in an appendix, we discuss a more general result concerning the discretisation of smooth moving frames, and the continuum limit of equicontinuous families of discrete moving frames, with an example.

2. BACKGROUND

2.1. Moving frames. Given a Lie group G acting on a manifold M with a left action, so that

$$G \times M \rightarrow M, \quad h \cdot (g \cdot z) = (hg) \cdot z,$$

one can define a left (resp. right) group-based moving frame as a map which is equivariant with respect to the action on M and the left (resp. inverse right) action of G on itself, specifically,

$$\rho : M \rightarrow G, \quad \rho(g \cdot z) = g\rho(z) \quad (\text{resp. } \rho(g \cdot z) = \rho(z)g^{-1}).$$

We call such an equivariant map a left (resp. a right) moving frame. The inverse of a left moving frame is a right one, and vice versa.

Given a group G acting on a manifold M , the existence of a moving frame on the open subset $U \subset M$ is guaranteed if:

- (i) the orbits of the group action all have the same dimension and foliate U ,
- (ii) there is a transverse cross-section \mathcal{K} to the orbits such that for each orbit \mathcal{O} , the intersection $\mathcal{O} \cap \mathcal{K}$ contains a single point, and
- (iii) the group element taking $z \in \mathcal{O}(z)$ (where $\mathcal{O}(z)$ is the orbit through z) to $\mathcal{O}(z) \cap \mathcal{K}$, is unique.

In this case, a right moving frame $\rho : U \rightarrow G$ is given by $\rho(z) \cdot z \in \mathcal{K}$, that is, $\rho(z)$ is the unique element of G taking z to the unique element of $\mathcal{K} \cap \mathcal{O}(z)$. Since \mathcal{K} is transverse to the orbits, the frame defines local co-ordinates given by $z \mapsto (\rho(z), \rho(z) \cdot z) \in G \times \mathcal{K}$.

In the continuous case of moving frames, the manifold M could be the jet space $J^{(\ell)}(\mathbb{R}^p, M)$. In this case it is known ([17]) that provided the action is locally effective on subsets, as ℓ grows the prolonged action of G on $J^{(\ell)}(\mathbb{R}^p, M)$ becomes locally free. The work of Boutin ([8]) discusses what happens for products $M^{\times q}$ as q grows, with G acting with the diagonal action. *In any event, we make the assumption that for large enough dimension, there is a neighborhood of the identity in the group in which a moving frame can be obtained locally.*

A common way to obtain the moving frame is through a normalization process. One can describe normalization equations as those defining the transverse section, \mathcal{K} , to the orbits of the group. If the normalisation equations are given as $\{\Phi = 0\}$, then the conditions above for the existence of a moving frame are the conditions under which the implicit function theorem can be applied to solve $\Phi(g \cdot z) = 0$ for $g = \rho(z)$. Since both $g = \rho(h \cdot z)$ and $g = \rho(z)h^{-1}$ solve $\Phi(g \cdot (h \cdot z)) = 0$, and the implicit function guarantees a unique solution, then $\rho(h \cdot z) = \rho(z)h^{-1}$, that is, ρ is equivariant. Typically, the normalisation equations, for which \mathcal{K} is the zero set, are algebraic. Indeed, in many applications, the cross-section is a co-ordinate plane, so that the normalisation equations involve certain co-ordinates being set to a constant. Since there will be many transverse cross-sections to the orbits, the choice of \mathcal{K} can greatly decrease (or increase) the calculations involved. Part of the ‘‘art’’ of the moving frame in applications is the choice of cross-section, or equivalently, the choice of normalisation equations.

Given a moving frame (left or right) one can generate all possible invariants of the action. Indeed, if ρ is a right moving frame, the expressions

$$\rho(u) \cdot v$$

for any $u, v \in M$ are clearly invariant; their coordinates are called the *basic invariants*. One can easily see that any invariant of the action is a function of these, using the *replacement rule*: If $I : M \rightarrow \mathbb{R}$ is invariant under the action, so that $I(g \cdot v) = I(v)$ for all $g \in G$, then setting $g = \rho(u)$, one obtains

$$I(\rho(u) \cdot v) = I(v).$$

Different choices of the manifold M gives rise to different familiar cases. For example, if M is the jet space $J^{(\infty)}(\mathbb{R}^p, P)$ for some manifold P where G acts, and G acts on M via the natural *prolonged action* given by the chain rule, then ρ would generate moving frames on p -submanifolds and the invariants will be standard differential invariants (for example, curvatures, torsions, etc). If $M = \times_k P$ is the cartesian product of a manifold P where G acts, and G acts on M through the diagonal action, then the invariants are the so-called *joint invariants* (see [51]).

The authors of [41] defined discrete moving frames, essentially a choice of group element associated to each vertex in an equivariant way. The discrete moving frame can be defined to act naturally under the shift operator, greatly simplifying calculations with discrete (difference) equations. We next review this definition.

Let G^N denote the Cartesian product of N copies of the group G . Allow G to act on the left on G^N using the diagonal action $g \cdot (g_r) = (gg_r)$. We also consider what we have called the “right inverse action” $g \cdot (g_r) = (g_r g^{-1})$.

Definition 2.1 (Discrete moving frame). We say a map

$$\rho : M^N \rightarrow G^N$$

is a left (resp. right) *discrete moving frame* if ρ is equivariant with respect to the diagonal action of G on M^N and the left (resp. right inverse) diagonal action of G on G^N . Since $\rho((x_r)) \in G^N$, we will denote by ρ_s its s th component, that is $\rho = (\rho_s)$, where $\rho_s((x_r)) \in G$ for all s . Equivariance means,

$$\rho_s(g \cdot (x_r)) = \rho_s((g \cdot x_r)) = g\rho_s((x_r)) \quad (\text{resp. } \rho_s((x_r))g^{-1})$$

for every s . Clearly, if $\rho = (\rho_s)$ is a left moving frame, then $\hat{\rho} = (\rho_s^{-1})$ is a right moving frame.

As in the original group-based moving frame definition, if $(u_s) \in M^N$, one can define

$$\rho_s \cdot u_r = I_s^r.$$

The coordinates of these invariants for any r generate all other invariants *even when s is fixed* (see [41]). We note that the action induces an action on the coordinate functions, the same as it induces an action on any function, specifically, $g \cdot f(u_r) = f(g \cdot u_r)$. The components of I_s^r will be invariant as I_s^r is, they are called the *basic invariants*.

One can describe a smaller set of generating invariants, the Maurer–Cartan invariants.

Definition 2.2. Let (ρ_s) be a left (resp. right) discrete moving frame evaluated along a twisted N -gon, the element of the group

$$K_s = \rho_s^{-1} \rho_{s+1} \quad (\text{resp. } \rho_{s+1} \rho_s^{-1})$$

is called the left (resp. right) s -Maurer–Cartan matrix for ρ . We call the equation $\rho_{s+1} = \rho_s K_s$ the discrete left s -Frenet–Serret equation.

If $G \in GL(n, \mathbb{R})$, then the entries of the Maurer–Cartan matrices, together with the basic invariants I_s^s , generate all other invariants. See [41] for more details.

2.2. Lattices. Lattices are subsets of \mathbb{Z}^p with a variety of properties. We first define adjacency.

Definition 2.3. Two points $m, n \in \mathbb{Z}^p$ are said to be *adjacent* if

$$\sum_j |m_j - n_j| = 1.$$

Definition 2.4. We say that a subset $\Gamma \subset \mathbb{Z}^p$ is a *connected lattice* if it consists of a single point, or, if between any two points $\mathbf{m}_1, \mathbf{m}_2 \in \Gamma$ there is a path, $\mathbf{m} = \mathbf{n}_1, \mathbf{n}_2, \dots, \mathbf{n}_N = \mathbf{m}_2$ such that \mathbf{n}_i is adjacent to \mathbf{n}_{i+1} for $i = 0, \dots, N - 1$.




FIGURE 2. In (ii) and (iii), Γ is connected. The subsets (i), (iv) and (v) are not connected lattices as there are points not adjacent to any other point.

Natural operators on \mathbb{Z}^p are the well-known shift operators, namely $\{\mathcal{T}_i\}_{i=1}^p$, where

$$\mathcal{T}_i(n_1, \dots, n_i, \dots, n_p) = (n_1, \dots, n_i + 1, \dots, n_p).$$

We will also consider lattices in a manifold.

Definition 2.5. The image of a lattice $\Gamma \subset \mathbb{Z}^p$ in a manifold M by a map $\Phi : \Gamma \rightarrow M$ is denoted by L , and is also called a lattice. We assume this map to be injective, a condition which will be relaxed under controlled conditions in the second part of this paper. The adjacency in L is determined from that of Γ .

We will assume from the beginning that the lattice $L \subset M$ is contained within one coordinate chart for the manifold, so that for all practical purposes we can assume the manifold is \mathbb{R}^n , or often a parametrized surface with the parameter serving as local coordinates. We will also assume that lattices are connected (one can study each connected components separately).

The lattice itself does not need to be covered with one lattice neighborhood, however. We define next a lattice variety, which will allow us to work on lattice models of spheres and tori.

Definition 2.6. A *lattice variety* \mathcal{L} is a set of points that can be covered by subsets \mathcal{L}_α , each of which mapped by an injection ϕ_α to a connected lattice $\mathcal{L}_\alpha \subset \mathbb{Z}^p$ for some fixed p . Every adjacency in \mathcal{L} is contained in at least one of the \mathcal{L}_α . Furthermore, in the overlap $\mathcal{L}_\alpha \cap \mathcal{L}_\beta$, the gluing map $\phi_\alpha \circ \phi_\beta^{-1}$ preserves adjacency. We call $(\mathcal{L}_\alpha, \phi_\alpha)$ a local lattice coordinate system.

Lattice coordinates essentially introduces an order in the lattice (inherited from \mathbb{Z}^p through ϕ_α) so one can clearly define shifts. We say that a shift map is defined at a point in \mathcal{L} , if it is defined in at least one chart. Since the chart interchange maps preserve adjacency, the existence of a shift map is well-defined.

3. MOVING FRAMES ON LATTICES AND LATTICE VARIETIES

Let \mathcal{L} be a lattice variety and let $N = |\Gamma|$ be the number of vertices in \mathcal{L} . Let \mathcal{L}_N be the set of p -lattice varieties in M with N vertices (in much of what follows N could be infinite).

3.1. Moving frames, invariants and Maurer–Cartan invariants. Let G be a group acting on \mathcal{L}_N (for example, if the lattice lives inside a manifold with a group action, the action would be the one induced on the lattice), and for simplicity assume that it is a left action (that is, $g \cdot (h \cdot u) = (gh) \cdot u$. A parallel description can be made for right actions).

A discrete moving frame will associate an element of the group to each vertex in the lattice in an equivariant fashion.

Definition 3.1. [Moving frames on lattices]

Let \mathcal{U} be a subset of \mathcal{L}_N . We say

$$\rho : \mathcal{U} \rightarrow G^N$$

is a left (resp. right) *discrete moving frame* on \mathcal{U} whenever ρ is equivariant with respect to the action of G on \mathcal{L}_N and the left (resp. inverse right) diagonal action of G on G^N . That is, if $\rho = (\rho_i)_{i=1}^N$ denote the components of ρ in G^N , then

$$\rho_i(g \cdot \mathcal{L}) = g\rho_i(\mathcal{L}) \quad (\text{resp. } \rho(g \cdot \rho_i(\mathcal{L})) = \rho_i(\mathcal{L})g^{-1}),$$

with $|\Gamma| = N$.

In general, moving frames exist only locally, which is the reason why we need to restrict its domain in \mathcal{L}_N . Given a lattice variety \mathcal{L} , and a coordinate system indexed by $\Gamma \subset \mathbb{Z}^p$, the moving frame ρ assigns group elements at each vertex (thus N of them). We will call ρ_R *the moving frame at the vertex* $R \in \Gamma$. Note when the index is applied we are assuming the use of local variety coordinates.

It is a simple matter to go from a moving frame to a discrete moving frame by taking a family of cross-sections, one per lattice vertex, as stated in the following result.

Proposition 3.2. *Let $\{\mathcal{S}_R \mid R \in \Gamma\}$, be a family of sections, with \mathcal{S}_R transverse to the orbit of G at u_R , assumed to be indexed locally by Γ . Let $g = (g_R) \in G^N$ be uniquely determined by the condition*

$$(1) \quad g_R \cdot \mathcal{L} \in \mathcal{S}_R$$

for $\mathcal{L} \in \mathcal{L}_N$ and $R \in \Gamma$ in some coordinate system for \mathcal{L} . Then $(g_R) = (\rho_R)$ is a local right discrete moving frame.

The proof of this statement is straightforward from the discussion of the moving frame. We note that a moving frame is defined as an element of the group on the entire lattice, but normalization equations give frames defined only locally. The interchange maps from one domain to another will play a role in what follows.

Remark on Notation. From now on a multi-index will denote the use of local lattice coordinates, while the lack of it will indicate global definitions. Also, we will denote by ρ_R each individual component of ρ , or the moving frame at the vertex u_R .

Example 3.3. Consider 1-lattices - or polygons - in the Euclidean plane. The group $E(2)$ can be identified with the subgroup of $GL(3, \mathbb{R})$ given by

$$(2) \quad g = \begin{pmatrix} 1 & 0 \\ b & \Theta \end{pmatrix}$$

with $\Theta \in O(2)$ and $b \in \mathbb{R}^2$. It acts on \mathbb{R}^2 as

$$\begin{pmatrix} 1 & 0 \\ b & \Theta \end{pmatrix} \begin{pmatrix} 1 \\ x \end{pmatrix} = \begin{pmatrix} 1 \\ \Theta x + b \end{pmatrix}$$

with $x \in \mathbb{R}^2$. We choose as our transverse cross-section, the one given in coordinates by $\rho_n \cdot u_n = 0$ and $\rho_n \cdot u_{n+1} = e_1 \|\Delta u_n\|$, where $\Delta u_n = u_{n+1} - u_n$. If ρ_n is displayed in the matrix representation as in (2), solving the system

$$\rho_n \cdot u_n = \Theta_n u_n + b_n = 0, \quad \rho_n \cdot u_{n+1} = \Theta_n u_{n+1} + b_n = \Theta_n \Delta u_n = \|\Delta u_n\| e_1$$

results in the *left* moving frame

$$\rho_n^{-1} = \begin{pmatrix} 1 & 0 \\ -\Theta_n^{-1} b_n & \Theta_n^{-1} \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ \|\Delta u_n\| e_1 & \frac{u_n}{\|\Delta u_n\|} & -J \frac{u_n}{\|\Delta u_n\|} \end{pmatrix}$$

where $J = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ is the canonical symplectic matrix.

Definition 3.4 (Invariants and basic invariants). We say the function on \mathcal{L}_N

$$I : \mathcal{L}_N \rightarrow \mathbb{R}$$

is a lattice invariant under the action of G if $I(g \cdot \mathcal{L}) = I(\mathcal{L})$ for any $g \in G$, $\mathcal{L} \in \mathcal{L}_N$. A local invariant will have the same property in some coordinate chart.

Given a right moving frame ρ on p -lattices, we call the invariants

$$I_R = \rho_R(\mathcal{L}) \cdot \mathcal{L}$$

the *basic invariants*, where $\rho(\mathcal{L}) = (\rho_R(\mathcal{L}))$ with $\rho_R(\mathcal{L}) \in G$ (the group element at the vertex in position R depends, in general, on all vertices of the lattice). Once we choose coordinates in \mathcal{L} , given by (u_J) , the local invariants are defined to be

$$I_R^M = \rho_R((u_J)) \cdot u_M$$

for $R, M \in \Gamma$. These are clearly invariants of the action.

The basic invariants take their names from the fact that they generate all other local invariants. In fact, they do generate them even when R is fixed.

Proposition 3.5. *If I is any lattice invariant, then I can be written as a function of the basic invariants I_R^N for any fixed $R \in \Gamma$.*

Proof. This is an immediate consequence of so-called *replacement rule*. Let (u_N) represent the vertices of the lattice variety. If $I = I((u_N))$ is an invariant of the action, then $I(g \cdot (u_N)) = I((u_N))$ for all $g \in G$, and in particular for $\rho_R((u_J))$. Thus

$$I(\rho_R((u_J)) \cdot (u_N)) = I(I_R^N) = I((u_N))$$

which shows us how to write I in terms of the basic invariants with R fixed. \square

From this wealth of invariants we will be selecting a few, the so-called Maurer–Cartan invariants. They are the discrete analogue of the invariants defining the classical Frenet–Serret equations and, like their continuous counterpart, they will form a generating system.

From now on we will extend the shift operators in the standard way to algebraic functions of u_J using the properties $\tau_i(u_J u_R) = \mathcal{T}_i u_J \mathcal{T}_i u_R$. We can also apply a shift to ρ_J by sending ρ_J to ρ_{J+e_i} or we could apply it to ρ_J by shifting the variables u_R that ρ_J depends on. But notice that unless the sections in 1 are shifts of each other (i.e. $\mathcal{S}_{R+e_i} = \tau_i \mathcal{S}_R$ for all $i = 1, \dots, p$ and all $R \in \Gamma$), these two operations do not need to produce the same result. Given that in many of our situations and in all of our examples we do assume the sections to be invariant under the shifts, we will abuse the notation and denote all these maps by \mathcal{T}_i , so that, for example, $\mathcal{T}_i \rho_J = \rho_{J+e_i}$.

Definition 3.6. [Maurer–Cartan invariants] Let ρ be a right moving frame along p -lattices. We define the right (R, i) -Maurer–Cartan group element to be $K_{(R,i)}$, the element of the group given by

$$K_{(R,i)} = (\mathcal{T}_i \rho_R) \rho_R^{-1} = \rho_{R+e_i} \rho_R^{-1}.$$

Its local coordinates (or the entries of the matrix, if $G \subset \text{GL}(n, \mathbb{R})$), will be called the (R, i) -Maurer–Cartan invariants.

Theorem 3.7. *Let ρ be any right moving frame. The (R, i) Maurer–Cartan invariants, $i = 1, \dots, p$, $R \in \Gamma$, together with $\rho_R((u_J)) \cdot u_R = I_R^R$, $R \in \Gamma$, generate all other invariants for the action of G on \mathcal{L}_N .*

Proof. The proof is based on what are commonly known as the *recursion formulae*.

Directly from the definitions we get that

$$K_{(R,i)} \cdot I_R^M = (\mathcal{T}_i \rho_R) \rho_R^{-1} \cdot (\rho_R \cdot u_M) = \rho_{R+e_i} \cdot u_M = I_{R+e_i}^M$$

and from

$$(3) \quad K_{(R,i)} \cdot I_R^M = I_{R+e_i}^M$$

we have

$$I_R^M = K_{(R,i)}^{-1} \cdot I_{R+e_i}^M$$

Now, since Γ is connected, given $M \in \mathbb{Z}^p$, any $R \in \mathbb{Z}^p$ is related to $M \in \mathbb{Z}^p$ through either recurrently increasing or decreasing its individual components, using the shift operator. At each step the invariant obtained when increasing or decreasing the components in M is generated by those in previous steps and by Maurer–Cartan invariants. Thus, we can start using I_M^M and reach I_R^M , for any R , using both versions of the recursion formulas. This proves the statement of the theorem. \square

Example 3.8. *The simplest example is the translation group viewed as a subgroup of $\mathrm{GL}(n+1, \mathbb{R})$ and acting on \mathbb{R}^n as*

$$g \cdot u = \pi \left(\begin{pmatrix} 1 & 0 \\ a & I \end{pmatrix} \begin{pmatrix} 1 \\ u \end{pmatrix} \right) = \pi \left(\begin{pmatrix} 1 \\ a + u \end{pmatrix} \right) = a + u,$$

where π is the projection in the last n components. If (u_R) is a p -lattice in \mathbb{R}^n , a transverse section to the orbit of the group at u_R is given by $u_R = 0$. Thus, the moving frame is determined by $g \cdot u_R = 0$, which implies $a = -u_R$. The basic invariants are $I_R^M = \rho_R \cdot u_M = u_M - u_R$, while the Maurer–Cartan matrices are $\rho_{R+e_i} \rho_R^{-1}$ whose only non-constant entries are the Maurer–Cartan invariants $u_R - u_{R+e_i}$, $R \in \Gamma$, $i = 1, \dots, n$. It is straightforward to show that the I_R^M can be written in terms of the Maurer–Cartan invariants $I_R^{R+e_i}$.

Example 3.3 cont. *In the case of the Euclidean plane, we found a moving right moving frame given by*

$$(4) \quad \rho_n = \begin{pmatrix} 1 & 0 \\ -\Theta_n u_n & \Theta_n \end{pmatrix}$$

where

$$\Theta_n^{-1} = \begin{pmatrix} \frac{u_n}{\|\Delta u_n\|} & -J \frac{u_n}{\|\Delta u_n\|} \end{pmatrix}.$$

The basic invariants in this case are given by

$$\rho_n \cdot u_m = \Theta_n(u_m - u_n) = \frac{1}{\|\Delta u_n\|} \begin{pmatrix} u_n \cdot (u_m - u_n) \\ \det(u_n, u_m - u_n) \end{pmatrix}$$

for any n, m . Notice that $\rho_n \cdot u_{n+1} = \|\Delta u_n\| e_1$. The Maurer–Cartan matrix is given by

$$(5) \quad \rho_{n+1} \rho_n^{-1} = \begin{pmatrix} 1 & 0 \\ -\Theta_{n+1} \Theta_n b_n + b_{n+1} & \Theta_{n+1} \Theta_n^{-1} \end{pmatrix}$$

where

$$-\Theta_{n+1} \Theta_n b_n + b_{n+1} = -\Theta_{n+1}(u_{n+1} - u_n) = \frac{1}{\|\Delta u_{n+1}\|} \begin{pmatrix} -u_{n+1} \cdot \Delta u_n \\ \det(u_{n+1}, u_n) \end{pmatrix}$$

and

$$\Theta_{n+1} \Theta_n^{-1} = \frac{1}{\|\Delta u_{n+1}\|} \frac{1}{\|\Delta u_n\|} \begin{pmatrix} u_{n+1} \cdot u_n & \det(u_n, u_{n+1}) \\ -\det(u_{n+1}, u_n) & u_{n+1} \cdot u_n \end{pmatrix} = \begin{pmatrix} \cos \alpha_n & \sin \alpha_n \\ -\sin \alpha_n & \cos \alpha_n \end{pmatrix}.$$

Where α_n is the angle between u_{n+1} and u_n . Therefore, a generating set for the Maurer–Cartan invariants are $\|\Delta u_n\|$ and α_n , for all n . Since the basic invariants are also generated by $\|\Delta u_n\|$ and the angle between u_n and u_m , the Maurer–Cartan invariants generate all the basic ones.

3.2. Syzygies.

3.2.1. *Basic local syzygies.* From the definition of Maurer–Cartan matrix, $K_{(N,i)} = (\mathcal{T}_i \rho_N) \rho_N^{-1}$, we have

$$\mathcal{T}_j K_{(J,i)} = (\mathcal{T}_j \mathcal{T}_i \rho_J) \mathcal{T}_j \rho_J^{-1} = (\mathcal{T}_j \mathcal{T}_i \rho_J) \rho_J^{-1} K_{(J,j)}^{-1}$$

and also

$$\mathcal{T}_i K_{(J,j)} = (\mathcal{T}_i \mathcal{T}_j \rho_J) \mathcal{T}_i \rho_J^{-1} = (\mathcal{T}_i \mathcal{T}_j \rho_J) \rho_J^{-1} K_{(J,i)}^{-1}.$$

Given that shifts commute, we obtain

$$(6) \quad (\mathcal{T}_j K_{(J,i)}) K_{(J,j)} = (\mathcal{T}_i K_{(J,j)}) K_{(J,i)}.$$

This expression gives us a number of algebraic relationships between the different Maurer–Cartan invariants. We will refer to these as *basic local syzygies*, a discrete generalization of the differential syzygies that are satisfied by differential invariants, such as the Codazzi–Mainardi equations for Euclidean invariants defined on surfaces. The above syzygies generate most of the possible algebraic relations among Maurer–Cartan invariants. Further independent syzygies may be created by the topology of the lattice \mathcal{L} .

Definition 3.9. We say two syzygies are equivalent up to basic syzygies if one of them is an consequence of the other and syzygies of the form (6).

To describe global syzygies, we need first to define the discrete fundamental group of the lattice and related material standard in the study of the topology of graphs [47].

3.2.2. Discrete fundamental group. Let L be a p -lattice in M , that is, the image of a map from a connected p -dimensional subset Γ of \mathbb{Z}^p to M . As before, we will denote the image of $J \in \Gamma$ as $\phi(J) = u_J \in M$. For simplicity, let us assume that $\Gamma = \mathbb{Z}^p$, although one can apply much of what we will say next to other cases. Notice that, in principle, we are allowing cases when the map ϕ is not 1-to-1 so L does not need to have a trivial topology.

Definition 3.10 (Paths). We say a subset $P \subset L$ is a *path of length r* joining two points $a, b \in M$, if it can be ordered as $P = \{x_i\}_{i=0}^r$, with $x_0 = a$, $x_r = b$ and x_i adjacent to x_{i+1} , for all $i = 0, \dots, r-1$. We say the path is *closed* if $a = b$; we say it is *simple* if $x_i \neq x_j$ for any $i \neq j$ (except perhaps $x_0 = x_r$ if closed). Notice that by giving the vertices of the path in a certain order we are implicitly assigning an orientation to it. This will be relevant once we associate syzygies to closed paths.

Definition 3.11 (Edge). Given a path γ in a lattice, $\gamma = \{x_i\}_{i=0}^r$, we say the ordered pair $[x_i, x_{i+1}]$ is an *edge* of the path. The ordering gives an orientation of the edge.

Definition 3.12 (Sum of paths). Consider the set of all closed paths with base point a . One can define the *sum* of two such paths by concatenation; that is, if $\{x_i\}_{i=0}^r$ and $\{y_j\}_{j=0}^s$ are two paths, their sum is given by $\{x_i\} + \{y_j\} = \{z_k\}_{k=0}^{r+s+1}$ with

$$z_i = x_i, \quad i = 0, \dots, \quad z_{r+j+1} = y_j, \quad j = 0, \dots, s.$$

If the paths are not closed, but $x_r = y_0$, one can equally define the sum of the paths by concatenation.

Definition 3.13 (Basic homotopy). A transformation of a path $\{x_i\}$, $\phi(\{x_i\}) = \{y_j\}$ is a basic homotopy if $\{y_j\}$ is equal to $\{x_i\}$ except for

- (1) adding or removing a subpath of the form $[x_r, x_{r+1}] + [x_{r+1}, x_r]$;
- (2) changing a subpath of the form $[x_r, x_{r+1}] + [x_{r+1}, x_{r+2}]$ by one of the form $[x_r, y_{r+3}] + [y_{r+3}, x_{r+2}]$, where $x_r, x_{r+1}, x_{r+2}, y_{r+3}$ form a basic square of the lattice.

Transformations (1)-(2) are called the two *basic homotopies*.

Figure 3 shows examples of basic homotopies.




FIGURE 3. Examples of the two basic homotopies applied to the blue paths

Decomposition of two homotopies into basic homotopies

The lattice is missing the inner vertices: the red path wraps around the missing tube while the blue one is homotopically equivalent to the constant path at a .





FIGURE 4. The red and blue paths on the left are homotopically equivalent, with the yellow and green paths showing the sequence of basic homotopies required, while the ones on the right are not.

Definition 3.14. We say the two paths $\{x_i\}_{i=0}^r$ and $\{y_j\}_{j=0}^s$ joining a and b , are *homotopically equivalent* if $x_0 = a = y_0$, $x_r = b = y_s$ and either the paths are equal, or one can be transformed to the other by a finite sequence of basic homotopies.

By construction, homotopy of paths joining a and b is an equivalence relation. Figure 4 shows pairs of homotopic and nonhomotopic paths.

Definition 3.15 (Discrete fundamental group). Consider the space of all closed paths based at a , and let $\pi_1(L, a)$ be the set of homotopy classes of these paths. The operation above endows $\pi_1(L, a)$ with a group structure. We call $\pi_1(L, a)$ the discrete fundamental group of L .

The fundamental group does not depend on the point a chosen, as far as the lattice is connected. (It suffices to join a to a different point b using a path P , and

use P to relate closed paths based on a to those based on b by conjugation, as done in the continuous case.)

3.2.3. *Syzygies associated to closed paths on a lattice and global syzygies.* Assume we have a moving frame along a path. To each edge of the path we can associate a Maurer-Cartan matrix of invariants as follows:

Assume either that $x_i = u_J$ and $x_{i+1} = \mathcal{T}_k u_J = u_{J+e_k}$, or that $x_{i+1} = \mathcal{T}_k^{-1} u_J = u_{J-e_k}$. To $[x_i, x_{i+1}]$ we associate the matrix

$$(7) \quad K_{(J,k)} = \mathcal{T}_k(\rho_J)\rho_J^{-1} = \rho_{J+e_k}\rho_J^{-1}$$

in the first case, and

$$(8) \quad K_{(J-e_k,k)}^{-1} = (\mathcal{T}_k(\rho_{J-e_k})\rho_{J-e_k}^{-1})^{-1} = \rho_{J-e_k}\rho_J^{-1}$$

in the second case.

We note that the choice of Maurer-Cartan matrix depends on the orientation of the path, with an edge being associated to the inverse matrix if the orientation is reversed.

Next we will associate a group element $K(\gamma)$ to each path γ on the lattice, namely, the product of the Maurer-Cartan matrices along the path. From the definition of the Maurer-Cartan matrices in terms of the discrete frame, it will be evident this product telescopes to involve only the discrete frame at the endpoints of the path, provided the discrete frame is defined along the whole path. *Nevertheless, in terms of the components of the Maurer-Cartan matrices, the Maurer-Cartan invariants, the product will not telescope – this is the syzygy.* Evaluating $K(\gamma)$ for closed paths γ , leads to relations on the invariants. It becomes important to find those relations which are non-trivial, in the sense that they are not an algebraic consequence of basic local syzygies, given in Equation 6.

In what follows, we will show that $K(\gamma)$ is a homotopy invariant. Evaluating $K(\gamma)$ on closed paths which are not homotopic to the constant trivial path leads to relations on the Maurer-Cartan invariants which cannot be obtained in terms of the basic local syzygies.

We start with paths which lie in the domain of a discrete frame. This, of course, need not be the case since the existence of moving frames is guaranteed only locally. If we need to cover the lattice with several co-ordinate patches on which discrete moving frames exist, we will obtain invariant transition matrices associated with the cover of the lattice defined by the domains of the discrete moving frames. We discuss this more involved case later in this section.

Definition 3.16. Assume a discrete frame exists along a path γ . The product of the Maurer-Cartan matrices along γ is denoted $K(\gamma)$. Specifically, for the path $\{x_i\}_{r=0}^{i=r}$, we have

$$K(\{x_i\}_{r=0}^{i=r}) = K([x_{r-1}, x_r]) \cdots K([x_0, x_1]),$$

where $K([x_i, x_{i+1}])$ is the Maurer-Cartan element associated to the edge $[x_i, x_{i+1}]$ as in (7)-(8) so that $K([x_i, x_{i+1}]) = K^{-1}[x_{i+1}, x_i]$. If γ is the constant (trivial) path, we define $K(\gamma) = e$.

It is evident that if $\gamma = \gamma_1 + \gamma_2$, then $K(\gamma) = K(\gamma_2) \cdot K(\gamma_1)$. (See Definition 3.12).




FIGURE 5. In this Figure, we see that along the closed path $(x_1, x_2, x_3, x_4 = x_1)$ we have $K_4 K_3 K_2 K_1 = e$, the identity in G . This is equivalent to the basic local syzygy, Eqn. (6).

In Figure 5, we illustrate the basic local syzygy, in the form $K(\{x_1, x_2, x_3, x_1\}) = e$ along a closed path of length four.

Proposition 3.17. *Let γ_1 and γ_2 be two paths joining a and b in the lattice L for which a discrete moving frame exists. Assume that γ_1 is homotopic to γ_2 , then $K(\gamma_1) = K(\gamma_2)$.*

Proof. Since a homotopy is a finite composition of basic homotopies, it suffices to show that if γ_1 and γ_2 differ by a basic homotopy, then $K(\gamma_1) = K(\gamma_2)$.

In the case (1), this is trivial since the only difference between $K(\gamma_1)$ and $K(\gamma_2)$ is a product of the form $K_i K_i^{-1} = e$. In the case (2) it is equally simple since they differ only by a product $K_i K_{i+1}$ appearing in $K(\gamma_1)$ and $K_j K_{j+1}$ appearing in $K(\gamma_2)$, with $K_i K_{i+1} = K_j K_{j+1}$ being a local syzygy since their vertices form a square in the lattice. □

The following Corollary is an immediate consequence of the previous proposition.

Corollary 3.18. *If our lattice variety is covered by one coordinate system and there exists a moving frame defined everywhere, each class $[\gamma]$ of the fundamental group of the lattice defines what we will call a global syzygy of the Maurer-Cartan invariants, in the form $K(\gamma) = e$.*

For example, if a discrete frame is defined on an annular lattice L , then there will be a path not homotopic to the constant path within L , but $K(\gamma) = e$ for all closed paths.

We now turn our attention to paths that move through different domains of discrete frames. We can still associate a group element K to the path by patching local products of Maurer-Cartan matrices, assuming the local neighborhoods where the different discrete frames are defined overlap. Assume we have two subpaths $\{x_0, \dots, x_i\}$ and $\{x_i, x_{i+1}, \dots, x_j\}$ of a closed path, and assume we can find a moving frame ρ_s at x_s , $s = 0, \dots, i$ and a different moving frame $\hat{\rho}_s$ for $s = i, i + 1, \dots, j$ (notice the overlap in the subindices). We define $M(x_i) = \hat{\rho}_i \rho_i^{-1}$ so that $\hat{\rho}_i = M(x_i) \rho_i$. Clearly, since both ρ_i and $\hat{\rho}_i$ are right equivariant, the matrix $M(x_i)$ is invariant. Then, to the path $\gamma = \{x_0, \dots, x_j\}$ we can associate the product of

invariant matrices

$$\begin{aligned} K(\gamma) &= (\hat{\rho}_j \hat{\rho}_{j-1}^{-1})(\hat{\rho}_{j-1} \hat{\rho}_{j-2}^{-1}) \dots (\hat{\rho}_{i+1} \hat{\rho}_i^{-1}) M(x_i) (\rho_i \rho_{i-1}^{-1}) \dots (\rho_1 \rho_0) \\ &= \hat{K}_{j-1} \dots \hat{K}_i M(x_i) K_{i-1} \dots K_0 \end{aligned}$$

with the invariant matrix $M(x_i)$ linking the Maurer-Cartan matrix in one coordinate system to the next.

Definition 3.19. If a vertex x lies in the domains of both the discrete frame ρ and the discrete frame $\hat{\rho}$, we say the group element

$$(9) \quad M(x; \hat{\rho}, \rho) = \tilde{\rho}(x) \rho(x)^{-1}$$

is the transition Maurer-Cartan matrix at the vertex x , associated with the change of frame from ρ to $\hat{\rho}$.

Important Assumptions. We already have the assumption on our co-ordinate charts \mathcal{L}_α , that every edge appears in at least one \mathcal{L}_α , and so every Maurer-Cartan matrix can be written in (at least one) co-ordinate system. We assume further that every edge is in a domain of a discrete frame. In this way, every Maurer-Cartan matrix, every transition matrix, and their local products, can be expressed with respect to a single set of co-ordinates. By taking a refinement of our co-ordinate cover as necessary, we therefore assume that our cover consists of sets which are domains of both frames and co-ordinate charts, and that every edge appears in at least one element of the cover.

If a vertex lies in the intersection of several frame domains, there will be *co-cycle conditions*. For example, if $x \in \text{dom}(\rho_\alpha) \cap \text{dom}(\rho_\beta) \cap \text{dom}(\rho_\delta)$, then clearly

$$(10) \quad M(x; \rho_\delta, \rho_\beta) M(x; \rho_\beta, \rho_\alpha) M(x; \rho_\alpha, \rho_\delta) = e.$$

On the other hand, two equal paths can have different group element $K(\gamma)$ if the choices of either the initial or the final moving frames are different. Thus, our element of the group does not only depend on γ but also on the initial and final choice of moving frame. In this case we will denote the group element above

$$K(\gamma, \alpha, \beta)$$

for the element of the group that starts in $\text{dom}(\rho_\alpha)$ and ends in $\text{dom}(\rho_\beta)$, and where these are the choices of frame for the calculation of the initial and final Maurer-Cartan matrices. By analogy, we will also denote by $M(x; \alpha, \beta)$ the matrix $M(x; \rho_\alpha, \rho_\beta)$.

Lemma 3.20. *$K(\gamma, \alpha, \beta)$ does not depend on the choice of coordinates or moving frames one chooses along γ , only on the initial and final ones.*

Proof. Assume at some point x we make different choices of moving frame at a point x in the overlap of (at least two) different frame domains, so we move from ρ_μ to ρ_η for one path and from ρ_μ to ρ_ν for the other. In that case we introduce the transition factor $M(x; \eta, \mu)$ in one of the lifts, and $M(x; \nu, \mu)$ in the other one, and we continue the different paths using the corresponding choices. At some point we need to come back to a common choice, even if that happens only at the end of the path. But when we expand the different factors of $K(\gamma, \alpha, \beta)$ in terms of moving frames, the intermediate factors all vanish as we saw before, and the difference is only at the beginning and at the end of the product. Thus, without losing




FIGURE 6. Several ways a path can move through an interchange.

generality we can assume that we come back to a common moving frame right after we introduce the split.

That is, a path includes the factor

$$M(x; \beta, \eta)M(x; \eta, \mu),$$

while the other includes

$$M(x; \beta, \nu)M(x; \nu, \mu).$$

But using (10) we have that both these factors are equal to $M(x; \beta, \mu)$, and hence they are equal. \square

We now argue that $K(\gamma, \alpha, \beta)$ is still a homotopy invariant for paths that start and end with the α and β choice of moving frame, even if the path moves through changing domains of discrete frames (and changing co-ordinate systems). Consider Figure 6, in which we assume that $[x, y]$, $[y, z]$ are edges in the domain of ρ , while $[x, t]$ and $[t, z]$ are edges in the domain of $\hat{\rho}$. We consider paths γ_i running from x_0 , in the domain of ρ , to y_0 in the domain of $\hat{\rho}$. If γ_1 goes via x and t , then we must switch from ρ to $\hat{\rho}$ at x . We achieve this by using the matrix $M(x) = \hat{\rho}(x)\rho(x)^{-1}$ (which is invariant as the frames are right frames), and then defining $K(\gamma_1)$ to be

$$\begin{aligned} K(\gamma_1) &= \hat{K}([t, y_0])\hat{K}([x, t])M(x)K([x_0, x]) \\ &= \hat{\rho}(y_0)\hat{\rho}(t)^{-1}(\hat{\rho}(t)\hat{\rho}(x)^{-1})(\hat{\rho}(x)\rho(x)^{-1})\rho(x)\rho(x_0)^{-1} \\ &= \hat{\rho}(y_0)\rho(x_0)^{-1}. \end{aligned}$$

Considering the path γ_2 from x_0 to y_0 via z results in

$$K(\gamma_2) = \hat{K}([t, y_0])\hat{K}([z, t])M(z)K([y, z])K([x_0, y])$$

which is also equal to $\hat{\rho}(y_0)\rho(x_0)^{-1}$ by a similar argument, and thus we have

$$(11) \quad K(\gamma_1) = K(\gamma_2)$$




FIGURE 7. The situation of Figure 6 translated to $\bar{\mathcal{L}}$, the “lift” of \mathcal{L} .

Since we have homotopy invariance of the K element within domains, in this way we can see that even passing through a change of domain, we maintain homotopy invariance.

In order to prove homotopy invariance in general, we construct a “lift” $\bar{\mathcal{L}}$, of \mathcal{L} , with respect to a good cover of \mathcal{L} , using which we can keep track of which discrete frame we are using on our paths in \mathcal{L} . The lift $\bar{\mathcal{L}}$ that we construct is not a lattice variety in general, and does not lie in M , but nevertheless serves our purpose here. To construct $\bar{\mathcal{L}}$, we take the disjoint union of the charts, together with their edges (adjacencies), and for every $x \in \mathcal{L}_\alpha \cap \mathcal{L}_\beta$, we take a new adjacency, or edge, $[x|_\alpha, x|_\beta]$, and let this be a new edge in $\bar{\mathcal{L}}$, with the associated Maurer-Cartan element being $\rho_\beta \rho_\alpha^{-1}$, the transition Maurer-Cartan matrix. See Figure 7, which shows the lifting for the case of Figure 6. Define the projection from $\bar{\mathcal{L}}$ to \mathcal{L} as the natural projection that collapses the different copies of the vertices which lie in intersections of charts. That is

$$\pi : \bar{\mathcal{L}} \rightarrow \mathcal{L}, \quad \pi(x_i) = \begin{cases} x_i & \text{if } x_i \text{ belongs only to one domain} \\ x & \text{if } x_i = x_\alpha \text{ where } x_\alpha \text{ is a lift of } x \in \mathcal{L}_\alpha \end{cases}.$$

A path $\bar{\gamma}$ in $\bar{\mathcal{L}}$ is a lift of a path in \mathcal{L} when the projection of $\bar{\gamma}$ is γ . Lifts may not be unique, as they depend, for example, on the element of $\pi^{-1}(a)$ at which the path begins, where a is the initial vertex of γ .

Changing from one frame to another along a path is, in $\bar{\mathcal{L}}$, simply proceeding from one vertex to another, with the transition Maurer-Cartan elements, Definition 3.19, being the group element associated to the new edge. We note that a change of co-ordinates simply changes the local labelling of the points, and so is less important when considering the group element associated to a path.

We now show that in $\bar{\mathcal{L}}$ we have new basic local syzygies obtained from the transition Maurer-Cartan matrices. These new local syzygies allow us to obtain a result similar to that of Proposition 3.17, but for the more general situation where




FIGURE 8. New edges in $\bar{\mathcal{L}}$ give rise to new local syzygies, here,
 $K([y_\beta, x_\beta])K([x_\beta, x_\alpha]) = K([y_\beta, y_\alpha])K([y_\alpha, x_\alpha])$

we need to change domains of our moving frames. Consider Figure 8. Let the domain of ρ be \mathcal{L}_α and the domain of $\hat{\rho}$ be \mathcal{L}_β . Consider the simple closed path $\gamma = \{x|_\alpha, x_\beta, y|_\beta, y|_\alpha, x|_\alpha\}$. If $K([y_\beta, x_\beta]) = \hat{\rho}(y)\rho(x)^{-1}$, $K([x_\beta, x_\alpha]) = \hat{\rho}(x)\rho(x)^{-1}$, $K([y_\beta, y_\alpha]) = \hat{\rho}(y)\rho(y)^{-1}$, $K([y_\alpha, x_\alpha]) = \rho(y)\rho(x)^{-1}$, the K element for this path is,

$$K(\gamma) = (\rho(x)^{-1}\rho(y)) (\rho(y)\hat{\rho}(y)^{-1}) (\hat{\rho}(y)\hat{\rho}(x)^{-1}) (\hat{\rho}(x)\rho(x)^{-1}) = e$$

showing this path defines a basic syzygy in $\bar{\mathcal{L}}$.

Finally, we define the monodromy of a closed path in \mathcal{L} . When such a path is lifted to $\bar{\mathcal{L}}$ it need not be closed, as it may begin in one frame domain and return in another. Consider the lift $\bar{\gamma}$ of a path γ beginning at x_0 in the domain of the frame ρ_α , and ending at x_0 in the domain of the frame ρ_β . We can close the path in $\bar{\mathcal{L}}$, by adding to $\bar{\gamma}$, the edge $[x_\beta, x_\alpha]$, but in principle this need not happen. Therefore, the element of the group associated to the lift would not be equal to e , but rather to $M(x; \alpha, \beta)$ (or its inverse, depending on the orientation). We call this group element *the monodromy of the lift*.

We can now state the more general Theorem concerning homotopy invariance.

Theorem 3.21. *To each path γ in the lattice variety \mathcal{L} , and to each choice of initial and final moving frame, we can associate a group element, $K(\gamma, \alpha, \beta)$, such that if γ_1 and γ_2 are homotopic in \mathcal{L} , then $K(\gamma_1, \alpha, \beta) = K(\gamma_2, \alpha, \beta)$.*

Proof. We first note that $\bar{\mathcal{L}}$ has essentially a global moving frame, in the sense that every vertex has an equivariant group element associated, namely ρ_α for x_α and ρ_β for x_β , and so on. Hence we can define $K(\gamma)$ for any path, the same way we did previously.

We first note that any lift of the trivial path to a closed path in $\bar{\mathcal{L}}$ has, by the co-cycle conditions, Equation (10), $K(\bar{a}) = e$. When we lift a path γ to $\bar{\mathcal{L}}$, we need to keep track of which discrete frame we are working in, but as we saw before, the element $K(\gamma, \alpha, \beta)$ is only affected by the beginning and end choices.

Assume two paths are homotopically equivalent, and let's lift the homotopy. By construction of $\bar{\mathcal{L}}$, we can assume the endpoints of the lift of the homotopic paths also remain fixed and determined by the α and β choices. As in the proof of Proposition 3.17, it will suffice if we show that two paths that differ by one of the

basic homotopies have the same $K(\gamma, \alpha, \beta)$, even if we need to change the moving frame domain. But this was already proved in the argument concerning Equation (11). \square

The syzygy of any closed path γ , where the domain of the discrete frames are considered to be the same at the end and at the beginning, is $K(\gamma, \alpha, \alpha) = K(\gamma) = e$, while those where $\alpha \neq \beta$ will have a non trivial monodromy $K(\gamma, \alpha, \beta) = M \neq e$. Furthermore, different choices of α give rise to the same syzygy: the group elements are related by conjugation

$$K(\gamma, \alpha, \alpha) = \rho_\alpha \rho_\beta^{-1} K(\gamma, \beta, \beta) \rho_\beta \rho_\alpha^{-1},$$

and the transition matrices $\rho_\alpha \rho_\beta^{-1}$ are essentially a change of coordinates. Indeed, a frame defines a local coordinate system of the form $U \times \mathcal{K}$ where U is a neighbourhood of the identity in G and \mathcal{K} is the cross-section which has invariants for coordinates ([37], Chapter 4.) We thus have the following Corollary.

Corollary 3.22. *Let $[\gamma]$ be the homotopy class of a closed path γ in \mathcal{L} . Each element $[\gamma]$ of $\pi_1(\mathcal{L})$ gives rise to a syzygy on the Maurer-Cartan invariants, in the form $K(\gamma) = e$.*

Notice that associating a syzygy to a closed path with a monodromy is essentially the same as associating a syzygy to the closed path for which the beginning and end moving frames are the same. Indeed, $K(\gamma; \alpha, \alpha) = K(\gamma, \alpha, \beta)M(x; \beta, \alpha)$, and hence $K(\gamma; \alpha, \alpha) = e$ is the same syzygy as $K(\gamma; \alpha, \beta) = M(x; \alpha, \beta)$.

Example 3.23. Consider a bi-periodic lattice $L = \{z_{n,m}\}_{n,m \in \mathbb{Z}}$ with $z_{n+kT_1,m} = z_{n,m}$ and $z_{n,m+kT_2} = z_{n,m}$ for some periods $T_1, T_2 \in \mathbb{Z}$. For simplicity, assume we can find a global moving frame $\rho = (\rho_{n,m})$. The topology of this lattice is comparable to that of a torus, and one can easily show that

$$\pi_1(L) = \mathbb{Z}^2.$$

The two generators of $\pi_1(L)$ (marked in Figure 9 with different colors) are represented by the two global syzygies

$$K_{0,m}K_{1,m} \dots K_{T_1-1,m} = e, \quad K_{n,0}K_{n,1} \dots K_{n,T_2-1} = e.$$

If a global moving frame does not exist, then the product might be equal to a monodromy matrix that will depend on the choice of moving frame at the beginning and the end of the closed path. If we choose the same moving frame, then the syzygy will be independent of the point chosen as beginning and end, and it will be as above.

4. CONTINUOUS LIMITS OF THE DISCRETE PICTURE: LATTICE BASED MULTISPACE

In this section we show how one can construct a continuous moving frame as the limit of a discrete one, and vice-versa, by coordinating the transverse sections that determine them. This is achieved using our *lattice based multispace*, in which derivatives and their finite difference approximations exist in a single manifold containing both the jet bundle and cartesian products of the base space. Both smooth and discrete frames are then part of a single frame on this multispace, and their relationship is given by the continuity of the multispace frame under coalescence. We show in this case that not only moving frames but also discrete invariants and




FIGURE 9. Discrete toroidal lattice variety, with periodicities $T_1 = 8$ and $T_2 = 6$

local discrete syzygies converge to differential invariants and differential syzygies respectively. We will use our multispace constructions to describe discretizations of integrable systems, and to finite difference models of variational systems, in §5. In the Appendix, we will show that more generally, an equicontinuous family of discrete frames will converge to a smooth frame.

First of all, we recall the definition of multispace in one variable, as developed by P.J. Olver in [50]. Olver provides co-ordinates for his space of pointed curves in terms of the Lagrange approximation of the curve via interpolation at specific (given) points. The coordination of the discrete and the smooth pictures is a consequence of the fact that Lagrange interpolation becomes Hermite interpolation under coalescence. In order to provide coordinates for our higher dimensional generalisation of Olver's construction, we need to restrict our 'pointed surfaces' to those where the Lagrange and Hermite interpolations are similarly related and to where the interpolations vary in a smooth manner with respect to the data. The details of the interpolation are critical, since the coefficients of the interpolation polynomial will define the desired coordinates. We use the theory of multivariate polynomial approximation due to de Boor and Ron, [5], [6] and [7], described in §4.2. We then describe our lattice based multispace and prove that it is a manifold. In fact, we detail two related versions of multispace, one containing the jet bundle $J(M, \mathbb{R})$ and one containing the jet bundle $J(U, M)$ where $U \subset \mathbb{R}^p$ for any $p \leq \dim M$. Both arise in the applications. Thereafter we show that the limit of the discrete Maurer Cartan matrices are the smooth Maurer Cartan matrices and that the local syzygies (Equation 6) limit to the so-called zero curvature condition of the smooth Maurer Cartan matrices, in §4.4. We also describe mixed discrete/continuous cases.

4.1. Multispace in one variable. The idea behind the definition of multispace is to create a manifold where both discrete and continuous cases coexist in one overarching smooth construction, where the continuous frame is a limit of the discrete, and the limit of the discrete data is the continuous data. Multispace resembles the jet spaces, but includes also discrete versions of the jet spaces.

Given a manifold M , we let $C^{(n)} = C^{(n)}(M)$ denote the set of all $(n+1)$ -pointed curves contained in M ; that is, the set of $(z_0, \dots, z_n; C)$, where C is a curve and z_i

are $n + 1$ points in C , not necessarily distinct. We denote by

$$n_i = \#\{j \mid z_j = z_i\}$$

the number of points that coincide with z_i .

Definition 4.1. (Multispace for curves) Two $(n + 1)$ -pointed curves

$$\mathbf{C} = (z_0, \dots, z_n; C); \quad \tilde{\mathbf{C}} = (\tilde{z}_0, \dots, \tilde{z}_n; \tilde{C}),$$

have n^{th} order multi-contact if, and only if

$$z_i = \tilde{z}_i, \quad \text{and} \quad j_{n_i-1}C|_{z_i} = j_{n_i-1}\tilde{C}|_{z_i}, \quad \text{for each } i = 0, \dots, n,$$

where $j_k C$ denotes the k th jet of the curve C , and where $\sum_i n_i = n$. The n^{th} order multi-space, denoted $M^{(n)}$, is the set of equivalence classes of $(n + 1)$ -pointed curves in M under the equivalence relation of n^{th} order multi-contact. The equivalence class of an $(n + 1)$ -pointed curve \mathbf{C} is called its n^{th} order multi-jet, and is denoted by $\mathbf{j}_n \mathbf{C} \in M^{(n)}$.

When all the points are distinct, then two curves belong to the same equivalence class whenever they have the distinguished points z_i in common. Thus, we can identify this special subset with the off-diagonal Cartesian product, denoted by $M^{\circ(n+1)}$ in [50]. On the other hand, if all the points coincide, then the class is equal to the jet class. Thus, both extremes can be found in one space, together with all the intermediate cases. In the first part of [50] the main result is the following theorem.

Theorem 4.2. *If M is a smooth m -dimensional manifold, then its n^{th} order multispace $M^{(n)}$ is a smooth manifold of dimension $(n + 1)m$, which contains the off-diagonal part $M^{\circ(n+1)}$ of the Cartesian product space as an open, dense submanifold, and the n^{th} order jet space J^n as a smooth submanifold.*

The topology is inherited from that of the manifold M , and the proof is based on finding coordinate systems in a neighborhood of an equivalence class \mathbf{C} . The coordinate system is given by the classical divided differences and their limits. That is, given a curve C with $n + 1$ distinguished points $\{z_0, \dots, z_n\}$ and with a certain order of contact n_i at each point, there exists a unique polynomial p of degree n such that $p(z_i) = C(z_i)$ and such that $p^{(k)}(z_i) = C^{(k)}(z_i)$ for any $k \leq n_i$ and any $i = 0, \dots, n$. The polynomial is a natural representative of the class \mathbf{C} and its coefficients provide smooth coordinates in a neighborhood of \mathbf{C} . Of particular importance is that the coordinates are smooth under the coalescence of points z_i . For more details, see [50].

In the second part of the paper [50], the author assumes there is a Lie group G acting on the manifold M , and he defines the action of this group on the multispace as that naturally induced by it: the action on the differential part is the prolonged one, and explicit formulae for the action of the group on classical divided differences are given. He also explains how, assuming that one chooses a cross section to the orbit of the group at a point \mathbf{C} , and *requiring the local cross-section to be transverse also to the jet space* (thus defining a cross-section for the prolonged action on J^n), then one can find a moving frame for the action of the group on the multispace with the desired property, that is, the resulting moving frame will be the standard continuous moving frame when restricted to jets, and the discrete one

when restricted to $M^{\circ(n+1)}$. The overarching continuity of the multispace manifold guarantees that one is the continuous limit of the other as the points coalesce.

We note several features of Olver's multi-space for pointed curves:

- (i) the curve \mathbf{C} is not parametrised and the points z_i on the curve need not be labelled in order with respect to some parametrisation,
- (ii) coalescence can take place between any two of the z_i on the curve, see Figure 10,
- (iii) none of the z_i are distinguished in the sense that one of them is a natural base point for a projection of the multispace to M ,
- (iv) the pointed curve $\mathbf{j}_n\mathbf{C}$ is essentially a set of points with a contact condition at each point.




FIGURE 10. Coalescence of points in a pointed curve does not require the points to be adjacent, in some sense, on the curve. Note that there needs to be a well defined tangent of C at the point of coalescence.

In our construction of a higher dimensional lattice based multispace, only a version of property (iv) remains.

Our next section describes an interpolation scheme which can be applied to our geometric construction.

4.2. Multivariate interpolation. One of the main problems with multivariate interpolation is that the solution to the interpolation problem is not unique in general and it might not even exist. For example, if we fix the values of a function $f(x, y)$ at the two points $(1, 2)$, $(-1, 1)$ and we want to find a polynomial in x and y of minimum order, such that it coincides with the function at those points, we can use $f(x, y) = a + bx$ or $f(x, y) = a + by$, and there is no reason why we would chose one over the other. On the other hand, if we fix the value of the function at $(1, 2)$ and $(1, -1)$, then the first choice is not appropriate unless the function has the same value at both points, while the second one works. Thus, the choice of interpolating polynomial might depend on the data, it might not be unique, or even exist, and sometimes there is no reason to favor one choice over a different one. In the $p = 1$ interpolation case none of these problems exist.

Thus, a main question in multivariate interpolation is: is there a family of polynomials which can interpolate the values of a given function and which has properties

like generality, minimal degree, uniqueness and having a well-defined Hermite polynomial (one for which not only the values of the polynomial, but also its derivatives at the points coincide with those of the function) as points coalesce? These are the properties we will need if we want to use them to define smooth coordinates in our multi-space. This question was answered by C. de Boor and A. Ron in [5], [6] and [7]; we describe below their solution to the interpolation problem as it applies to our particular case.

Interpolation Notation Let Π be the set of p -variate polynomials, and Π' its algebraic dual. Let Λ be a subspace of Π' . We will denote by Π_k the subset of polynomials of order less than or equal to k .

Definition 4.3. We say that $P \subset \Pi$ is *correct* for Λ if for any continuous linear functional F on Λ there exist a unique $q \in P$ such that

$$F(\lambda) = \lambda(q)$$

for any $\lambda \in \Lambda$. We also say P interpolates Λ .

The space Π' can be identified with formal power series at the origin o , using the bilinear form

$$(12) \quad \langle q, f \rangle = \sum_k \frac{D^k f(o) D^k q(o)}{k!}.$$

See [5], [6] for more details.

Example 4.4. Lagrange Interpolation. If Λ is spanned by point-evaluations, $\Lambda = \{\lambda_\theta\}_{\theta \in \Theta}$ with $\lambda_\theta(p) = p(\theta)$, $\theta \in \mathbb{R}^p$, finding P correct for Λ solves a *Lagrange interpolation problem*. Indeed, if Θ represents a finite number of points in the parameter space $D \subset \mathbb{R}^p$, and λ_θ is evaluation at an element of Θ , then one can check that the power series representing λ_θ is the Taylor expansion of $e^{\theta \cdot \mathbf{x}}$ (see [5], [6]). An analytic function F defines a continuous linear functional on Λ via (12) and $F(\lambda_\theta) = \langle F, e^{\theta \cdot \mathbf{x}} \rangle = F(\theta)$. Thus, P is correct if for any F there exists $p \in P$ such that

$$F(\lambda_\theta) = F(\theta) = \lambda_\theta(p) = p(\theta)$$

for all $\theta \in \Theta$, which is the definition of Lagrange interpolation, see Example 4.9.

Example 4.5. Hermite Interpolation. If we choose instead the set

$$\Lambda = \{\lambda_{q,\theta}\}_{\theta \in \Theta, q \in V \subset \Pi}$$

where

$$\lambda_{q,\theta}(p) = \lambda_\theta(q(\nabla)(p)) = q(\nabla)(p)(\theta)$$

where ∇ is the gradient vector and $q \in V \subset \Pi$ is a properly chosen polynomial with coefficients reflecting the differential data we need to match (for example, $q(x, y) = x^2$ if we wish to match the second derivative in x), then the associated power series representing $\lambda_{q,\theta}$ is $q(\mathbf{x})e^{\theta \cdot \mathbf{x}}$ (see [5], [6]). Therefore, finding P correct for Λ is equivalent to finding a family of polynomials P such that for any linear functional on Λ represented by an analytic function F , there exists a unique polynomial $p \in P$ with the property

$$F(\lambda_{q,\theta}) = \langle F, q(\mathbf{x})e^{\theta \cdot \mathbf{x}} \rangle = q(\nabla)(F)(\theta) = \lambda_{q,\theta}(p) = q(\nabla)(p)(\theta)$$

for all $\theta \in \Theta$ and all $q \in V$. The different choices of V allow us to find unique polynomials that coincide with F and different choices of derivatives, or combinations

of derivatives, on Θ . This is the solution to *the Hermite interpolation problem*, see *Example 4.10.*

Definition 4.6. Given a formal power series at $x = 0$, call it f , we denote by f_{\downarrow} the homogeneous term in the power expansion of f of lowest order.

For example, if $\theta, \mathbf{x} \in \mathbb{R}^p$, $e_{\downarrow}^{\theta \cdot \mathbf{x}} = 1$.

Definition 4.7. Given a finite subset $\Lambda \subset \Pi'$, we can identify each of its elements with formal power series at the origin using (12), and we can consider $\widehat{\Lambda}$ to be the vector space spanned by Λ as represented by these series. Define the vector space

$$\Lambda_{\downarrow} := \text{span}\{f_{\downarrow}, f \in \widehat{\Lambda}\}.$$

For example, if we consider the planar case, and $\Theta = \{(1, -1), (0, 2)\}$, then $e^{(1, -1) \cdot \mathbf{x}} = 1 + x - y + \dots$ and $e^{(0, 2) \cdot \mathbf{x}} = 1 + 2y + \dots$. Since $e^{(1, -1) \cdot \mathbf{x}} - e^{(0, 2) \cdot \mathbf{x}} = x - 3y + \dots$, we have that Λ_{\downarrow} is the linear space spanned by the polynomials $p_1(x, y) = 1 = e_{\downarrow}^{(1, -1) \cdot \mathbf{x}}$ and $p_2(x, y) = x - 3y = (e^{(1, -1) \cdot \mathbf{x}} - e^{(0, 2) \cdot \mathbf{x}})_{\downarrow}$.

From now on, if Θ is a data set, we say that Λ_{\downarrow} is continuous on Θ if whenever the data Θ' is close to Θ in the standard product topology, then Λ'_{\downarrow} is close to Λ_{\downarrow} in the standard topology of polynomial spaces.

The following theorem is a compilation of results found in [7]. Since our constructions are lattice based, we assume that Λ is defined as in the Lagrangian interpolation associated to a lattice of points $\Theta \subset \mathbb{R}^p$, in which case we also write Λ as Λ^{Θ} . We omit the superscript Θ where the dependence of Λ on Θ is clear.

Theorem 4.8. ([7]) *The space Λ_{\downarrow} has the following properties:*

- (1) **Well defined.** *For any finite Θ , the assignment $\Lambda \rightarrow \Lambda_{\downarrow}$ exists, is unique and Λ_{\downarrow} is correct for Λ .*
- (2) **Continuity.** *Recall that Π_k is the set of p -variate polynomials of degree less or equal to k . If $\Pi_k \subseteq \Lambda_{\downarrow} \subseteq \Pi_{k+1}$, then the assignment $\Lambda \rightarrow \Lambda_{\downarrow}$ is continuous with respect to Θ .*
- (3) **Coalescence \implies Osculation.** *That is, the Lagrange interpolation becomes the Hermite interpolation under coalescence, provided the coalescence is well controlled, so that data points coalesce along embedded curves.*
- (4) Λ_{\downarrow} *is closed under differentiation and it is spanned by a homogeneous basis.*
- (5) **Minimal degree.** Λ_{\downarrow} *has minimal degree.*
- (6) **Monotonicity.** *If $\Theta \subset \Theta'$, then $\Lambda_{\downarrow}^{\Theta} \subset \Lambda_{\downarrow}^{\Theta'}$.*
- (7) **Cartesian product \implies tensor product.** *If Θ and Θ' are two data sets, then*

$$(13) \quad \Lambda_{\downarrow}^{\Theta \times \Theta'} = \Lambda_{\downarrow}^{\Theta} \otimes \Lambda_{\downarrow}^{\Theta'}.$$

- (8) **Constructible.** *The space Λ_{\downarrow} can be constructed in finitely many arithmetic steps.*

Notice that once an interpolating family is chosen, the actual interpolation problem reduces to solving a linear system of equations. Indeed, one would choose a linear combination of a basis generating Λ_{\downarrow} and write a linear system for the coefficients using the values of the function we wish to interpolate on the interpolating data. The solution of the linear system will define the proper combination of the basis and hence the interpolating polynomial for the function.

It is essential that our construction of multispace ensures that the interpolation problem satisfies Properties (2) and (3) of the above Theorem. To quote de Boor and Ron [7], concerning Property (2) in the above Theorem, (note that Λ_\downarrow is denoted as Π_Θ in the original [7])

“... If $\Theta \subset \mathbb{R}^3$ consists of three points, then one would choose $\Pi_\Theta \subset \Pi_1$ (as our scheme does) but if one of the three points approaches some point between the two other points, this choice has to change in the limit and hence cannot change continuously. As it turns out, our scheme is continuous at every Θ for which $\Pi_k \subseteq \Pi_\Theta \subseteq \Pi_{k+1}$, for some k .”

Next, we quote de Boor and Ron [7], concerning Property (3) in the above Theorem.

“... If, eg, a point spirals in on another, then we cannot hope for osculation. But if, eg, one point approaches another along a straight line, then we are entitled to obtain, in the limit, a match at that point also of the directional derivative in the direction of that line.”

These limitations on continuity and coalescence mean that in our construction of our multispace, we cannot be as free in our choice of generalisation of the one dimensional pointed curves used to construct Olver’s one dimensional multispace, as might seem possible. We return to this discussion in §4.3.

Example 4.9. *In the Lagrange interpolation case, assume $\Theta \subset \mathbb{R}^p$ is given, as before, by $(1, 2)$ and $(-1, 1)$, and assume $\{\lambda_\theta\}_{\theta \in \Theta}$ are the associated point-evaluation functionals. Thus, we have two series generating $\widehat{\Lambda}$, namely*

$$e^{x+2y} = 1 + x + 2y + o(\|\mathbf{x}\|), \quad e^{-x+y} = 1 - x + y + o(\|\mathbf{x}\|).$$

A basis for the vector space generated by these two series are the series of $e^{x+2y} = 1 + x + 2y + o(\|\mathbf{x}\|)$ and $e^{x+2y} - e^{-x+y} = 2x + y + o(\|\mathbf{x}\|)$, and so

$$\Lambda_\downarrow = \text{span}\{1, 2x + y\}.$$

If we choose as Θ the points $(1, 2)$ and $(1, -1)$, then the generators of $\widehat{\Lambda}$ are the same as before, but

$$\Lambda_\downarrow = \text{span}\{1, y\}$$

as expected.

If we choose four points of the form $(1, 2), (1 + \epsilon, 2), (-1, 1), (-1, 1 + \epsilon)$, then the four series generating $\widehat{\Lambda}$ are

$$f_1 = 1 + (1 + \epsilon)x + 2y + \frac{1}{2}((1 + \epsilon)x + 2y)^2 + o(\|\mathbf{x}\|^2), \quad f_2 = 1 - x + (1 + \epsilon)y + \frac{1}{2}(-x + (1 + \epsilon)y)^2 + o(\|\mathbf{x}\|^2).$$

and

$$f_3 = 1 + x + 2y + \frac{1}{2}(x + 2y)^2 + o(\|\mathbf{x}\|^2), \quad f_4 = 1 - x + y + \frac{1}{2}(-x + y)^2 + o(\|\mathbf{x}\|^2).$$

Since $(f_1 - f_3)_\downarrow = \epsilon x$ and $(f_2 - f_4)_\downarrow = \epsilon y$, if $\epsilon \neq 0$, then $1, x$ and y will be three of four generators for Λ_\downarrow . A fourth will be given by

$$\left(\frac{1}{\epsilon}(2(f_1 - f_3) + f_2 - f_4) - f_3 + f_4 \right)_\downarrow = (2 + \epsilon)x^2 - \frac{1 - \epsilon}{2}y^2,$$

and hence

$$\Lambda_{\downarrow} = \text{span}\left\{1, x, y, (2 + \epsilon)x^2 - \frac{1 - \epsilon}{2}y^2\right\}$$

will generate the interpolating polynomials.

Example 4.10. In the Hermite interpolation case, assume we would like to instead find interpolating polynomials that coincide with a function at $(1, 2)$ and $(-1, 1)$, and, say, with its partial with respect to x at $(1, 2)$ and with respect to y at $(-1, 1)$. In this case, the polynomials q_i generating the Hermite data are, at $(1, 2)$, $q_1(x, y) = x$, and at $(-1, 1)$ $q_2(x, y) = y$. One can see (see [5], [6]) that $\lambda_{q_1, (1, 2)}$ is represented by the analytic function $f(\mathbf{x}) = xe^{x+2y}$, while $\lambda_{q_2, (-1, 1)}$ is defined by $g(\mathbf{x}) = ye^{-x+y}$. Since $\Lambda = \text{span}\{\lambda_{(1, 2)}, \lambda_{(-1, 1)}, \lambda_{q_1, (1, 2)}, \lambda_{q_2, (-1, 1)}\}$, $\hat{\Lambda}$ has four generators, namely

$$f_1 = xe^{x+2y} = x + x^2 + 2yx + o(\|\mathbf{x}\|^2), \quad f_2 = ye^{-x+y} = y - xy + y^2 + o(\|\mathbf{x}\|^2).$$

and

$$f_3 = e^{x+2y} = 1 + x + 2y + \frac{1}{2}(x+2y)^2 + o(\|\mathbf{x}\|^2), \quad f_4 = e^{-x+y} = 1 - x + y + \frac{1}{2}(-x+y)^2 + o(\|\mathbf{x}\|^2).$$

Some simple and direct calculations show that $(f_3)_{\downarrow} = 1$, $(f_1)_{\downarrow} = x$, $(f_2)_{\downarrow} = y$ and $(2f_1 + f_2 - f_3 + f_4)_{\downarrow} = 2x^2 - \frac{1}{2}y^2$. Thus,

$$\Lambda_{\downarrow} = \text{span}\left\{1, x, y, 2x^2 - \frac{1}{2}y^2\right\}.$$

The interpolating polynomials in the previous example converge to these as $\epsilon \rightarrow 0$.

We are now ready to define a lattice based multi-space in several variables.

4.3. Multispaces in several variables. We define two related versions of multi-space, the first containing the jet bundle $J^{\ell}(M, \mathbb{R})$ and the second containing the jet bundle $J^{\ell}(U, M)$ where $U \subset \mathbb{R}^p$ is open.

We first recall that a point in the jet bundle $J^r(M, \mathbb{R})$ is represented by a triple $[x, f, U]_r$ where $x \in U \subset M$, the set U is open, and $f : U \rightarrow \mathbb{R}$, is a C^r function. We say that the triple $[x, f, U]_r \sim [x', f', U']_r$ if $x = x'$ and if, in some coordinate chart containing x , f and f' have the same derivatives up to order r , ([26], page 60.) The equivalence class $[x, f, U]_r$ is known as the r -jet of f at x . If $T_r(f)(x)$ is the order r Taylor polynomial of the (sufficiently smooth) function f at x , then $[x, f, U]_r \sim [x, T_r(f)(x), U]_r$, so we speak of $T_r(f)(x)$ as being the r -jet of f at x . Further, the coefficients of the r th order Taylor polynomials form local coordinates of the jet bundle $J^r(M, \mathbb{R})$. It is this construction that we generalise first.

To construct our multi-space which both contains and generalises the jet space $J^{\ell}(M, \mathbb{R})$, we proceed as follows:

1. We first define the kinds of lattices Γ that we will take as the models of domains for a mixed discrete-continuous jet at a point in M . They will be sets of points in \mathbb{Z}^p with ‘‘directional multiplicities’’ or more precisely, ‘‘required contact conditions’’ attached. Our model lattices will come equipped with a base point. We show further that these models have the properties required for the de Boor and Ron interpolation of functions on them to be smooth, both as their image in M is varied and under coalescence.
2. Next, for a model lattice $\Gamma \subset \mathbb{Z}^p \subset \mathbb{R}^p$, we let $U \subset \mathbb{R}^p$ be an open set, diffeomorphic to the unit disc in \mathbb{R}^p , containing Γ . Let ϕ be a diffeomorphic map of U into M , and $f : \phi(U) \rightarrow \mathbb{R}$ a function. Our multi-jet will then be an equivalence

For any p the lattice is a connected lattice containing $\ell + 1$, $(p - 1)$ -dimensional disjoint corner lattices of lengths $0, 1, \dots, \ell$. Figure 12 shows four corner lattices for $p = 2$ of lengths 4, 3, 2 and 2 (clockwise from the first quadrant). Figure 13 shows a forward $d = 3$ corner distribution. We will consider corner lattices with a distinguished point x_0 .




FIGURE 12. Corner lattices will allow the definition of finite differences needed to approximate a Taylor polynomial.




FIGURE 13. A forward $p = 3$ corner lattice of length $\ell = 3$.

Corner distributions contain exactly enough points to define interpolating polynomials that will converge to Taylor polynomials upon coalescing. For example, if $p = 2$, and $\ell = 2$, and $u_{0,0}$ is the base point, a possible interpolating polynomial will have coefficients which are a linear combination of the terms

$$1, \Delta_x f(u_{0,0}), \Delta_y f(u_{0,0}), \Delta_x^2 f(u_{0,0}), \Delta_y^2 f(u_{0,0}), \Delta_y \Delta_x f(u_{0,0})$$

where $\Delta_x|_{u_{0,0}}$ is the operator $\Delta_x|_{u_{0,0}}(f) = f(u_{1,0}) - f(u_{0,0})$, and similarly with $\Delta_y|_{u_{0,0}}$. To be able to uniquely determine an interpolating polynomial with those coefficients we will need to use all the vertices in a corner distribution like the one in the first quadrant of figure 12, with length $\ell = 2$ instead of 3. Different corner




FIGURE 14. Corner lattices under repeated coalescence of hyperplanes. An arrow indicates a zeroth and a first order contact condition is required at that point in the interpolation, an arc indicates a zeroth, first and second order contact condition, a plane indicates all zeroth, first and second order contact conditions in the plane are required. The squared points are the base points.

lattices will produce different types of interpolating polynomials, using forward, backwards or other types of differences.

From now on we will assume that all lattices Γ have a corner distribution of length ℓ with base point x_0 . When several lattices are involved we will denote the base point of Γ by x_Γ and the base point of $\phi(\Gamma)$ by $u_\Gamma = \phi(x_\Gamma)$.

Lemma 4.12. *Assume our data Θ is given by evaluating a function on the points of a corner lattice Γ of length ℓ . Then*

$$\Lambda_\downarrow = \Pi_\ell$$

Proof. Consider the monomials corresponding to polynomials of order ℓ . That is all monomials of the form x^I where $|I| \leq \ell$ is a multi-index $I = (i_1, \dots, i_p)$ and $x^I = x_1^{i_1} \dots x_p^{i_p}$. We want to show that these monomials are generators for Λ_\downarrow . Thus, we want to show that they generate f_\downarrow , where f is any possible linear combination of $e^{\theta \cdot x}$, $\theta \in \Theta$. Therefore, it suffices to show that the coefficients of the monomials x^I , $|I| \leq \ell$, in the Taylor expansions of $e^{\theta \cdot x}$ form an invertible matrix. Notice that we have the same number of monomials as points and hence the matrix is square.

Next, notice that the coefficients of these Taylor expansions are multiples of the monomials themselves evaluated at the point θ (since we are simply substituting x_i by $\theta_i x_i$ in the expansion). Therefore, the matrix of coefficients is given by a multiple of the matrix with rows (θ^I) , $\theta \in \Theta$, where θ^I has the different monomials in some prescribed order. This means that if the matrix were not to be invertible, we would have a combination $\sum_I a_I \theta^I = 0$ for $a_I \in \mathbb{R}$ which will be valid for all $\theta \in \Theta$. That is, the points in Θ lie in an algebraic variety of order ℓ .

But this is not possible: our points lie on $\ell + 1$ distinct hyperplanes, and ℓ of them contain enough points to make the hyperplane unique (only one of them contains a single point and does not determine it). That means the polynomial must factor through the ℓ linear equations that define the hyperplanes and must have at least order ℓ . But the extra single point left does not belong to any of the hyperplanes, and hence to ensure the point also lies in the variety we will need to use a polynomial of order at least $\ell + 1$. Therefore the matrix of coefficients is invertible and

$$\Lambda_{\downarrow} = \Pi_{\ell}.$$

□

This lemma ensures that property (2) in Theorem 4.8 is satisfied when we use corner lattices. Next, we consider coalescence of the point in the lattice, leading to a change in the lattice, an increase in the contact order and to our Lagrange interpolation becoming a Hermite interpolation. We restrict the coalesce to be along coordinate hyperplanes in the model lattices; we call these kinds of coalescence *hyperplane coalescence*. See Figure 14. Forbidden coalescences are also illustrated in Figure 15. We note that hyperplane coalescence maintains the basepoint of the lattice, although not in general the contact condition there. Coalescence means, in effect, that we consider some lattice points to not be distinct.




FIGURE 15. Examples of forbidden coalescence. We restrict coalescence to being along coordinate hyperplanes, which maintains a coordinate structure.

Since we want the multispace to be closed under hyperplane coalescence, we consider coalesced model lattices to come equipped with certain required contact conditions specified at particular points of the lattice. Suppose two points, u_0 and $u_1 = u_0 + h\mathbf{v}$ coalesce as $h \rightarrow 0$. Then the interpolation goes from matching the values of the function f at u_0 and u_1 , to matching the values $f(u_0)$ and $D(f)|_{u_0}(\mathbf{v})$. See Figure 11.

Repeated coalescence leads to higher order Hermite interpolation problems. If you begin with a corner lattice and coalesce along hyperplanes repeatedly, you will arrive at a single point at which the interpolation for a function f is simply the Taylor polynomial for f , with the order of the Taylor polynomial being the length of the corner lattice. In this way, the jet bundle $J^\ell(M, \mathbb{R})$ is a subset of our multispace. See Figure 14.

From now on we will abuse the notation and denote the (coalesced) model lattice also as Γ and the contact conditions on it by $\mathcal{C}(\Gamma)$. Under the map $\phi : U \supset \Gamma \rightarrow M$, $\phi(\Gamma)$ inherits contact conditions which we will denote by $\mathcal{C}(\phi(\Gamma))$.

The result of Lemma 4.12 remains unchanged under coalescence, as shown in the next lemma.

Lemma 4.13. *Assume Γ is a hyperplane coalesced corner lattice of length ℓ . Then $\Lambda_\downarrow = \Pi_\ell$.*

Proof. We can show this by induction. Assume that $\widehat{\theta}_j = \theta_j + h e_i$ approaches θ_j along the e_i direction, that is, as $h \rightarrow 0$, for $j \in \mathcal{J}$, (that is for those data points in a hyperplane). Except for the coalescing of a corner, we would have more than one point coalescing since we are using limits of hyperplanes.

Once more we want to prove that the coefficients of the monomials x^I in the expansion of $\exp(\theta \cdot \mathbf{x})$, for any $\theta \neq \widehat{\theta}_j$ and $x_i \exp(\theta_j \cdot \mathbf{x})$ for any $j \in \mathcal{J}$ define an invertible matrix. As in the proof of Lemma 4.12, the coefficient of x^I in $\exp(\theta_i \cdot \mathbf{x})$ is given by $\theta_i^I / I!$, and one can directly check that the coefficient of x^I in $x_i \exp(\theta_j \cdot \mathbf{x})$ is given by

$$\frac{1}{I!} \theta_j^{I - e_i} k_i$$

if $I = (k_1, \dots, k_p)$ and $I - e_i = (k_1, \dots, k_i - 1, \dots, k_p)$, for $j \in \mathcal{J}$. Assume that the matrix formed by these coefficients is not invertible. It means that there is a polynomial of degree ℓ of the form

$$(14) \quad \sum_{|I| \leq \ell} a_I \mathbf{x}^I$$

that vanishes on all $\theta \in \Theta$ except for $\widehat{\theta}_j$. As we saw in the proof of Lemma 4.12, we would need an ℓ -order polynomial to describe a polynomial vanishing on those points, given by the product of ℓ linear equations representing the ℓ hyperplanes (one of them will not be unique if it corresponds to a corner, since it is determined only by one point, but we can just make any choice). Now, if the lemma were not true, we would additionally need to satisfy the relation

$$\sum_{|I| \leq \ell} a_I \theta_j^{I - e_i} k_i = 0$$

where $I = (k_1, \dots, k_p)$. This is simply the derivative of the polynomial (14) with respect to x_i . Thus, the polynomial (14) needs to have order of contact 2 at the points on the hyperplane that coalesced, Typically one would need a second order polynomial along the hyperplane to achieve that, which would increase the degree of (14) to $\ell + 1$. Higher orders of contact would result from a higher number of coalescing hyperplanes and the order of the polynomial would generically increase accordingly, proving the Lemma. □




FIGURE 16. The data for a multispace element. The Figure shows functions f and f' which agree on the contact conditions at the lattice points, $\phi(\Gamma)$. We have $[\Gamma, f, \phi, U] \sim [\Gamma, f', \phi, U]$.

The definition of elements of our multispace as equivalence classes of functions which agree on images of lattices, is illustrated in Figure 16.

Definition 4.14. We define the multispace $(M, \mathbb{R})_p^{(\ell)}$ as the set of equivalence classes of quadruples $[\Gamma, f, \phi, U]$, where $(\Gamma, \mathcal{C}(\Gamma))$ is a (possibly hyperplane coalesced) corner lattice of length ℓ ; U is an open set of \mathbb{R}^p for some $p \leq \dim(M)$, diffeomorphic to the unit disc and containing Γ ; the map $\phi : U \rightarrow M$ is an embedding of U into a single coordinate chart of M , and the map $f : \phi(U) \rightarrow \mathbb{R}$ is smooth of order ℓ . We say two quadruples are equivalent, $[\Gamma, f, \phi, U] \sim [\Gamma', f', \phi', U']$, if the base points of $\phi(\Gamma)$, $\phi'(\Gamma')$ agree, that is $\phi(x_\Gamma) = \phi'(x_{\Gamma'})$, if $\phi(\Gamma) = \phi'(\Gamma')$, $\mathcal{C}(\phi(\Gamma)) = \mathcal{C}(\phi'(\Gamma'))$ and the contact conditions evaluate to zero on $f - f'$, that is, both f and f' satisfy the same contact conditions on each vertex of $\phi(\Gamma)$. (We note the contact conditions are linear.)

4.3.2. *The Main Theorem: multispaces are manifolds which contain the jet bundle.* Denote by $\mathcal{L}_p^\ell(M, \mathbb{R})$ the subset of $(M, \mathbb{R})_p^{(\ell)}$ given by non-coalesced lattices, that is a lattice with zero contact condition in every vertex. Denote also by $J_p^\ell(M, \mathbb{R})$ the space of regular jets of maps from p -dimensional submanifolds of M to \mathbb{R} . The main purpose of this section is to prove the following theorem.

Theorem 4.15. *Let M be a manifold of dimension m . Then, there exists a topology and a differential structure that makes $(M, \mathbb{R})_p^{(\ell)}$ into a smooth manifold of*

dimension $(m+1)\binom{p+\ell}{p}$, with the jet space $J_p^\ell(M, \mathbb{R})$ as a smooth submanifold and with $\mathcal{L}_p^\ell(M, \mathbb{R})$ as an open submanifold.

Before we start proving the theorem we recall that the main difficulty in defining interpolating polynomials is determining the family of polynomials with which we choose to interpolate. Once this is determined, the actual interpolating coefficients are simply given by the solution of a linear system of equations defined by the equality conditions we need to satisfy at the chosen points. Therefore, they will change smoothly insofar as the linear system (and hence the family of interpolating polynomials) changes continuously. Lemmas 4.12 and 4.13 together with Theorem 4.8 show that we are indeed in the smooth regime of the de Boor and Ron interpolation method.

Proof. First of all, let us show that $(M, \mathbb{R})_p^{(\ell)}$ is a bundle over the manifold M . Let us call x_Γ the base point of Γ . The fibre over $u \in M$ is the set of equivalence classes, $[\Gamma, f, \phi, U]$ where u satisfies $\phi(x_\Gamma) = u$. Define $\pi : (M, \mathbb{R})_p^{(\ell)} \rightarrow M$ to be the projection map

$$(15) \quad \pi([\Gamma, f, \phi, U]) = \phi(x_\Gamma) = u.$$

Let us first restrict to $\mathcal{L}_p^\ell(M, \mathbb{R})$. In order to have a well defined coordinate labelling on the lattice, we need to describe an ordering on the vertices. The first point will always be the base point. Although any ordering will do, we can order them by induction: if the lattice is one dimensional we move from smallest coordinate to largest coordinate in \mathbb{R} . If the lattice is two dimensional, then we order the hyperplanes from smallest length to largest length, and then order in each hyperplane as in the one dimensional case. In addition to its place in the ordering of the lattice points, each point in the lattice will have a coordinate in M . Since the image of the lattice lies in a single coordinate chart, these coordinates are consistent across the lattice, once the chart is designated. Thus the dimension of the set of embedded lattices is $m\binom{p+\ell}{p}$. This set can be considered as a submanifold of M^N , where $N = \binom{p+\ell}{p}$, with the subspace topology, and evidently has coordinates in terms of the coordinates on M .

If the lattices are coalesced, the coalesced vertices will have repeated coordinates, as many as the multiplicity requires. The set of coalesced lattices will have measure zero since it is the level set of some functions depending on coordinates.

If we now consider the relevant polynomial interpolation p_f of a (sufficiently smooth) function f on the embedded image of the corner lattice, we find that it can be described by $\binom{p+\ell}{p}$ coefficients. Since, by construction, $[\Gamma, f, \phi, U] \sim [\Gamma, p_f, \phi, U]$, we may add these coefficients to the coordinates of the equivalence class. That these coefficients can be used as coordinates follows from knowing that we are in the regime where the de Boor and Ron interpolation is unique and depends smoothly on the data, and from the fact that the coordinates of interpolations on the coalesced lattices are well behaved limit points of those on the non coalesced lattices.

A simple counting tells us that we have, in total, $(m+1)\binom{p+\ell}{p}$ coordinates needed to specify the equivalence class in each element of the fibre.

The bundle $(M, \mathbb{R})_p^{(\ell)}$ is taken to be the disjoint union of the fibres, and π as in (15) is its projection. We may take the usual bundle topology given by the

smoothness of the local trivializations defined by the coordinates

$$V \subset (M, \mathbb{R})_p^{(\ell)} \rightarrow \mathbb{R}^{m \binom{p+\ell}{p}} \times \mathbb{R}^{\binom{p+\ell}{p}}$$

which turns the space into a manifold. The map π is clearly smooth since the basepoint is simply the first element in the list of lattice points.

We have already shown that $J_p^\ell(M, \mathbb{R}) \subset (M, \mathbb{R})_p^{(\ell)}$, indeed, it is embedded as the submanifold whose first set of coordinates is the diagonal Δ in M^N , where $N = \binom{p+\ell}{p}$. We note the standard jet space is $J^\ell(M, \mathbb{R}) = J_m^\ell(M, \mathbb{R})$.

Notice that if we perturb slightly an uncoalesced lattice Γ , the lattice will remain uncoalesced, and so the subspace of classes of the form $[\Gamma, f, \phi, U]$, with Γ an uncoalesced corner lattice, is open. \square

Remark 4.16. Although we have required the image of a lattice in M to be within a single coordinate chart of M , this restriction is perhaps not vital. The generalisation requires, firstly, keeping track of which chart as well as the coordinate given by the chart, in the lattice part of the coordinates for the multispace element. Secondly, it requires the construction of interpolations which agree on intersections of coordinate charts.

We now construct our second multispace, $(\Omega, M)_p^\ell$, where $\Omega \subset \mathbb{R}^p$, $p \leq \dim M$, which represents local approximations to embedded parametrized p -submanifolds in M .

Definition 4.17. We define the multispace $(\Omega, M)_p^{(\ell)}$ to be the set of equivalence classes of triples $[\Gamma, \phi, U]$ where $\Gamma \subset U \subset \Omega \subset \mathbb{R}^p$ is a (possibly coalesced) lattice of length ℓ ; U is an open set of $\Omega \subset \mathbb{R}^p$, diffeomorphic to the unit disc, where $\phi : U \rightarrow M$ is smooth of order ℓ and where $\phi(U)$ is contained in a single coordinate chart of M . We say two triples are equivalent, $[\Gamma, \phi, U] \sim [\Gamma', \phi', U']$ if the base points of Γ, Γ' agree; if $\phi(\Gamma) = \phi'(\Gamma')$; if $\mathcal{C}(\phi(\Gamma)) = \mathcal{C}(\phi'(\Gamma'))$; and if $\phi(U)$ and $\phi'(U')$ have the same order of contact (as submanifolds) at $\phi(\Gamma)$ as indicated by $\mathcal{C}(\phi(\Gamma))$.

Theorem 4.18. *Let M be a manifold of dimension m . There exists a topology and a differential structure that makes $(\Omega, M)_p^{(\ell)}$ into a smooth manifold of dimension $2m \binom{p+\ell}{p}$.*

Proof. The proof of this theorem is almost identical to that of theorem 4.15. In local coordinates in M , we can write $\phi = (\phi_1, \dots, \phi_m)$. We would then apply the process in theorem 4.15 to produce $m \binom{p+\ell}{p}$ coordinates that determine the lattice, plus $m \binom{p+\ell}{p}$ coordinates determining the interpolating polynomial for ϕ_k with data Γ , for each $k = 1, \dots, m$. The remainder of the proof is identical to that of theorem 4.15. \square

The main purpose of our multispace construction is to show that a frame on a multispace is simultaneously a frame on the jet bundle and a frame on the set of all local lattice based discretisations. We now proceed to discuss how the group action on M induces a group action on multispace. We then show that a moving frame on multispace is simultaneously a smooth and a discrete frame, with the smooth frame being the limit of the discrete, and that the discrete Maurer Cartan invariants and their syzygies coalesce to the smooth ones.

4.3.3. *The action of a group on $(M, \mathbb{R})_p^{(\ell)}$.* Let G be a group acting on $M \times \mathbb{R}$,

$$G \times M \times \mathbb{R} \rightarrow M \times \mathbb{R}.$$

Recall the equivalence classes in $(M, \mathbb{R})_p^{(\ell)}$ have the form $[\Gamma, f, \phi, U]$ where $\Gamma \subset U \subset \mathbb{R}^p$, $\phi : U \rightarrow M$ and $f : \phi(U) \rightarrow \mathbb{R}$. Then for each $g \in G$ there is an induced action on the map $(\phi, f) : U \times \phi(U) \rightarrow M \times \mathbb{R}$ given by $(g \cdot (\phi, f))(x, z) = g \cdot (\phi(x), f(z))$.

The action of G on $(M, \mathbb{R})_p^{(\ell)}$ is given by

$$g \cdot [\Gamma, f, \phi, U] = [\Gamma, (g \cdot (\phi, f))_2, (g \cdot (\phi, f))_1, U]$$

We note that $(g \cdot (\phi, f))_1$ may be an embedding only for g in a neighbourhood of the identity. In this case we would have a local group action as in [52]. Since the action of G on $M \times \mathbb{R}$ preserves the order of contact, this action is independent of the representative of the class and is thus well-defined on $(M, \mathbb{R})_p^{(\ell)}$.

If, in a particular application G acts only on M , one can extend to an action on $M \times \mathbb{R}$ by taking $g \cdot (z, t) = (g \cdot z, t)$, that is the identity action on the \mathbb{R} coordinate.

If Γ is a single point so that $[\Gamma, f, \phi, U]$ is an element of $J_p^\ell(M, \mathbb{R})$, the induced action is the standard prolonged action, that is, as induced by the chain rule, while if the lattice Γ is uncoalesced and the group does not act on the parameter \mathbb{R} , the action is the diagonal action on M^N where $N = \binom{p+\ell}{p}$.

4.3.4. *The action of a group on $(\Omega, M)_p^{(\ell)}$.* Let G be a group acting on M

$$G \times M \rightarrow M$$

and let $[\Gamma, \phi, U] \in (\Omega, M)_p^{(\ell)}$ be an element of our multispace. Since $\phi : U \subset \Omega \rightarrow M$, we may define $g \cdot \phi : U \rightarrow M$ by $(g \cdot \phi)(x) = g \cdot (\phi(x))$. Define the action of G on $M_p^{(\ell)}$ to be

$$g \cdot [\Gamma, \phi, U] = [\Gamma, g \cdot \phi, U].$$

Again, since the action of G on $U \times M$ preserves the order of contact, this action is independent of the representative of the class and is thus well-defined on $(\Omega, M)_p^{(\ell)}$. Further, the action restricted to $J_p^\ell(\Omega, M) \subset (\Omega, M)_p^{(\ell)}$ is the standard prolonged action induced by the chain rule.

Notice that one could consider more general actions on $M \times U$, but we will omit it here to avoid further complications.

4.3.5. *Moving frames on $(M, \mathbb{R})_p^{(\ell)}$, $(\Omega, M)_p^{(\ell)}$.* We are now in a familiar situation: we have a smooth manifold $(M, \mathbb{R})_p^{(\ell)}$, or $(\Omega, M)_p^{(\ell)}$ and the action of a Lie group G on it. Thus, we can investigate the use of the standard moving frames method developed in [17] to establish the existence of a moving frame for the multispace.

Assume the action of the group G on $(M, \mathbb{R})_p^{(\ell)}$, or $(\Omega, M)_p^{(\ell)}$, is free and regular, so that the existence of a local moving frame is guaranteed. Let us choose a point in $L \in (M, \mathbb{R})_p^{(\ell)}$ or $(\Omega, M)_p^{(\ell)}$ and let \mathcal{S} be a section transverse to the orbit of G through L . Using the standard moving frame method, we would get a local moving frame ρ , defined for all $\hat{L} \in (M, \mathbb{R})_p^{(\ell)}$, or $(\Omega, M)_p^{(\ell)}$ in some neighbourhood of L , as the group element such that $\rho \cdot \hat{L} \in \mathcal{S}$. That is, ρ is an equivariant continuous map

$$\rho : \mathcal{U} \subset (M, \mathbb{R})_p^{(\ell)} \rightarrow G \quad (\text{resp. } \rho : \mathcal{U} \subset (\Omega, M)_p^{(\ell)} \rightarrow G)$$

where \mathcal{U} is an open neighborhood of L .

Remark 4.19. We note that there are results detailing conditions under which an action on a jet bundle will become free and regular for a sufficiently large prolongation, that is, by considering sufficiently high order derivative terms [17]. A discussion of the related results for a product action, under a sufficiently large number of products, is given by Boutin [8]. We conjecture that similar results will hold for actions on multispace.

Before proving our results, we give a simple example. The first example refers to the multispace, $(\mathbb{R}^2, \mathbb{R})_1^{(\ell)}$.

Example 4.20. We consider the two dimensional group $G = \mathbb{R}^+ \ltimes \mathbb{R}$ acting on $M \times \mathbb{R} = \mathbb{R}^2 \times \mathbb{R}$ as $(\lambda, \epsilon) \cdot (x, y, u(x, y)) = (x, y, \lambda u + \epsilon)$. We take ϕ to be the identity map for simplicity. At the corner lattice $\Gamma = \{(x_0, y_0), (x_0 + h, y_0), (x_0, y_0 + k)\}$ with (x_0, y_0) the base point, the multi jet coordinates are the lattice coordinates, and the coefficients of the linear interpolant to some function u on these three points. The interpolant is

$$\begin{aligned} p(u)(x, y) &= A + B \cdot (x - x_0) + C \cdot (y - y_0) \\ &= u(x_0, y_0) + \frac{u(x_0 + h, y_0) - u(x_0, y_0)}{h} (x - x_0) \\ &\quad + \frac{u(x_0, y_0 + k) - u(x_0, y_0)}{k} (y - y_0). \end{aligned}$$

Thus the coordinates coming from the interpolant are

$$(A, B, C) = \left(u(x_0, y_0), \frac{u(x_0 + h, y_0) - u(x_0, y_0)}{h}, \frac{u(x_0, y_0 + k) - u(x_0, y_0)}{k} \right).$$

We see that the coefficients are functions of u at the lattice points and so the induced group action on these coordinates is the natural action on functions. We thus have

$$(\lambda, \epsilon) \cdot (A, B, C) = (\lambda A + \epsilon, \lambda B, \lambda C).$$

Theorem 4.21. Assume ρ is a local moving frame for the action of G on $(M, \mathbb{R})_p^{(\ell)}$ (resp. $(\Omega, M)_p^\ell$) determined by a section transverse to an orbit of G . Assume that the section is also transverse to the orbit through a point $L \in J_p^{(\ell)}(M, \mathbb{R})$ (resp. $J^\ell(\Omega, M)$), that is, the domain of the multispace frame includes L .

Denote by $\rho(Q)$ the multispace frame at Q , and by $\rho(L)$ the smooth moving frame which is obtained by the restriction of the normalisation equations for ρ to the jet space, evaluated at L . If L and Q have the same base point, then as Q coalesces to L ,

$$\rho(Q) \rightarrow \rho(L).$$

Proof. We prove the result for $(M, \mathbb{R})_p^{(\ell)}$, the other case being similar. Notice that $\rho(L)$ is the standard moving frame on jet spaces obtained through a section transverse to the prolonged orbits, which is the multispace section restricted to the jet bundle. We note that $J_p^\ell(M, \mathbb{R})$ is a submanifold of $(M, \mathbb{R})_p^{(\ell)}$, invariant under the action of the group, so that the orbit of G through a point $L \in J_p^\ell(M, \mathbb{R})$ is equal to the prolonged orbit. Furthermore, if a section is transverse to the orbit within $(M, \mathbb{R})_p^{(\ell)}$, its intersection with $J_p^\ell(M, \mathbb{R})$ will also be transverse to the orbit within the submanifold $J_p^\ell(M, \mathbb{R})$.

The proof is now immediate from the fact that the moving frame on $(M, \mathbb{R})_p^{(\ell)}$ is a smooth map. \square

In Appendix A, we will show a different convergence theorem for families of discrete frames.

Example 4.22. *Continuing with our previous example 4.20, a moving frame ρ is defined by $(\lambda, \epsilon) \cdot (A, B, C) = (0, 1, *)$ where $*$ will be the invariant, $(\lambda, \epsilon) \cdot C|_\rho$. This yields $(\lambda, \epsilon)|_\rho = \left(\frac{h}{u(x_0+h, y_0) - u(x_0, y_0)}, -\frac{hu(x_0, y_0)}{u(x_0+h, y_0) - u(x_0, y_0)} \right)$ or in the standard matrix representation of this group,*

$$(16) \quad \rho = \begin{pmatrix} \frac{h}{u(x_0+h, y_0) - u(x_0, y_0)} & -\frac{hu(x_0, y_0)}{u(x_0+h, y_0) - u(x_0, y_0)} \\ 0 & 1 \end{pmatrix}.$$

It is readily seen that as $h \rightarrow 0$, we obtain the correct induced group action on u_x , and that the limiting frame

$$(17) \quad \rho(x, u, u_x) = \begin{pmatrix} 1 & -u \\ u_x & u_x \\ 0 & 1 \end{pmatrix}$$

is obtained both from the limit of the normalisation equations, $(\lambda, \epsilon) \cdot (u_x, u) = (1, 0)$ as well as being the limit of the frame itself. In other words, both the discrete and the smooth cases are handled by the one calculation of the frame on multispace.

4.4. The continuous limit of invariants and syzygies. We return to our discussion of discrete moving frames in the previous section, in which we have a lattice variety \mathcal{L}_N embedded in some manifold, and a discrete frame is a map $\rho : \mathcal{L}_N \rightarrow G^N$ where N is the number of points in the lattice. Suppose now that adjacent vertices in the lattice variety begin to coalesce. Under what conditions will the discrete frame converge to a smooth frame, $\bar{\rho}$? Furthermore, under what kinds of conditions will we have

$$(18) \quad \rho_{n+1}\rho_n^{-1} \sim \bar{\rho}(z + \epsilon v)\bar{\rho}(z)^{-1} \sim \exp(\epsilon(\mathcal{D}\bar{\rho})\bar{\rho}^{-1})$$

where \mathcal{D} is an invariant differential operator? And when will the discrete local syzygies converge to the local differential syzygies?

Example 4.23. *Let us go back to our running example 4.20. Setting $\rho_i = \rho(x_i, y_i, u(x_i, y_i), u(x_i + h_i, y_i), u(x_i, y_i + k_i))$ and calculating $\rho_1\rho_0^{-1}$, we have, using Equation (16),*

$$\rho_1\rho_0^{-1} = \begin{pmatrix} \frac{h_1(u(x_0+h_0, y_0) - u(x_0, y_0))}{h_0(u(x_1+h_1, y_1) - u(x_1, y_1))} & \frac{h_1(u(x_0, y_0) - u(x_1, y_1))}{u(x_1+h_1, y_1) - u(x_1, y_1)} \\ 0 & 1 \end{pmatrix}$$

If we now set $(x_1, y_1) = (x_0 + \bar{h}, y_0)$ and $h_0 = h_1 = h$ (say), so that the second lattice is the shift of the first, by $(\bar{h}, 0)$, then

$$\begin{aligned} & \left. \frac{d}{d\bar{h}} \right|_{\bar{h}=0} \rho_1 \rho_0^{-1} \\ &= \begin{pmatrix} -\frac{u_x(x_0 + h, y_0) - u_x(x_0, y_0)}{u(x_0 + h, y_0) - u(x_0, y_0)} & -\frac{hu_x(x_0, y_0)}{u(x_0 + h, y_0) - u(x_0, y_0)} \\ 0 & 0 \end{pmatrix} \\ &\xrightarrow{h \rightarrow 0} \begin{pmatrix} -\frac{u_{xx}|_{(x_0, y_0)}}{u_x|_{(x_0, y_0)}} & -1 \\ 0 & 0 \end{pmatrix} \\ &= \rho_x \rho^{-1} |_{J^2(M)} \end{aligned}$$

Alternatively, using the method we will apply in Example 4.27, we have setting $u(x + \bar{h}, y) = u(x, y) + \bar{h}u_x(x, y) + \mathcal{O}(\bar{h}^2)$, that

$$\begin{aligned} (19) \quad & \left. \rho_1 \rho_0^{-1} \right|_{J^2(M)} \\ &= \begin{pmatrix} 1 - \frac{u_x(x_0 + h, y_0) - u_x(x_0, y_0)}{u(x_0 + h, y_0) - u(x_0, y_0)} \bar{h} + \mathcal{O}(\bar{h}^2) & \frac{hu_x(x_0, y_0)}{u(x_0 + h, y_0) - u(x_0, y_0)} \bar{h} + \mathcal{O}(\bar{h}^2) \\ 0 & 1 \end{pmatrix} \\ &\sim \exp \left(\bar{h} \begin{pmatrix} -\frac{u_x(x_0 + h, y_0) - u_x(x_0, y_0)}{u(x_0 + h, y_0) - u(x_0, y_0)} & -\frac{hu_x(x_0, y_0)}{u(x_0 + h, y_0) - u(x_0, y_0)} \\ 0 & 0 \end{pmatrix} \right) \\ &\sim \exp(\bar{h} \rho_x \rho^{-1}) \end{aligned}$$

as above. We note that $\partial/\partial x$ is an invariant operator since the independent variables are invariant under the action, so that $\rho_x \rho^{-1}$ is invariant. Cases where the independent variables participate in the action require more care, as we indicate below.

Our first theorem concerns the convergence of the discrete Maurer–Cartan invariants to the smooth Maurer–Cartan invariants of the smooth frame.

We consider the case where the discrete frame (ρ_R) on the lattice variety $\mathcal{L}_N \subset M$ can be viewed as a multispace frame ρ , with

$$\rho_J = \rho([\Gamma_J, \phi, U]), \quad u_J = \phi(x_J),$$

where x_J is the basepoint of Γ_J

Assume the point $\mathcal{T}_i x_J$ is also part of Γ_J so that $\mathcal{T}_i u_J = \phi(\mathcal{T}_i x_J)$. If we have a path from $\mathcal{T}_i(u_J)$ to u_J in the multispace indicating their coalescence, we can use the multispace frame to differentiate $\mathcal{T}_i(\rho_J)\rho_J^{-1}$ with respect to the path parameter at u_J .

Theorem 4.24. *Let a multispace \mathcal{M} with the embedded jet bundle \mathcal{J} be given. Let a path in \mathcal{M} be given, $\epsilon \mapsto [\Gamma(\epsilon), \phi, U]$ for $0 \leq \epsilon \leq 1$. Let $u(\epsilon)$ be the base point of $\phi(\Gamma(\epsilon))$. Assume that both paths and the coalescence $\phi(\Gamma(\epsilon)) \rightarrow \phi(\Gamma_0)$ lie in the domain of a multispace frame ρ . Set $\rho(\epsilon) = \rho([\Gamma(\epsilon), \phi, U])$, that is, ρ evaluated at the point $[\Gamma(\epsilon), \phi, U]$. If $\mathbf{v} = d/d\epsilon|_{\epsilon=0} u(\epsilon)$, then*

$$(20) \quad \lim_{\Gamma(0) \rightarrow \Gamma_0} \left[\left. \frac{d}{d\epsilon} \right|_{\epsilon=0} \rho(\epsilon) \rho^{-1}(0) \right] = (D(\mathbf{v})\rho) \rho^{-1}$$

where $D(\mathbf{v}) = \sum v_i \partial / \partial x_i$ is the directional derivative.

If the independent variables participate in the group action, then $(D(\mathbf{v})\rho)\rho^{-1}$ will not be invariant. We recall that a smooth frame on a jet bundle yields a canonical, maximal set of invariant differential operators. Indeed, on a manifold with coordinates \mathbf{x} , if $g \cdot \mathbf{x} = \tilde{\mathbf{x}}$, then we define

$$(21) \quad \mathcal{D}_i = \left. \frac{\partial}{\partial \tilde{x}_i} \right|_{g=\rho} = \left(\left. \frac{\partial \tilde{\mathbf{x}}}{\partial \mathbf{x}} \right|_{ij} \right)^{-T} \left. \frac{\partial}{\partial x_j} \right|_{g=\rho}$$

Rewriting the partial derivatives in (20) in terms of the invariantized derivatives, by inverting equation (21), yields an invariant expression which will satisfy equation (18). Indeed, suppose that $d/d\epsilon = D(\mathbf{v}) = \sum_i v_i \partial / \partial x_i$. Inverting Equation (21) yields expressions of the form $\partial / \partial x_i = \sum_k A_{ik} \mathcal{D}_k$, so that

$$\frac{d}{d\epsilon} = \sum_i v_i \sum_k A_{ik} \mathcal{D}_k = \sum_k \left(\sum_i v_i A_{ik} \right) \mathcal{D}_k = \sum_k \bar{v}_k \mathcal{D}_k = \mathcal{D}(\bar{\mathbf{v}})$$

where the last equality defines the vector $\bar{\mathbf{v}}$, and $\mathcal{D}(\bar{\mathbf{v}})$ is the invariantised directional derivative.

The theorem follows from standard calculations with smooth moving frames. The invariantisation of the right hand side of Equation (20) can be calculated using the cross section that determines the frame and the infinitesimal action, see [37] for a detailed discussion.

Example 4.25 (Special Euclidean group action on curves in the plane). *The group is $SO(2) \times \mathbb{R}^2$ with the standard linear action of translation and rotation of curves on the plane, specifically,*

$$\begin{pmatrix} x(t) \\ y(t) \end{pmatrix} \mapsto \begin{pmatrix} \tilde{x}(t) \\ \tilde{y}(t) \end{pmatrix} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} x(t) - a \\ u(t) - b \end{pmatrix}.$$

This is the inverse (or right) action and so the equivariance of the frame is $\rho(g \cdot z) = g\rho(z)$.

The multispace frame calculation is as follows. We take an order 2 interpolation as we wish to achieve a multispace approximation of the curvature. If we interpolate the curve $(x(t), u(t))$ at $\Gamma = \{t_0, t_0 + h_1, t_0 + h_2\}$ with base point t_0 , we get

$$\begin{aligned} p(x(t)) &= A(x) + B(x) \cdot (t - t_0) + \frac{1}{2} C(x) \cdot (t - t_0)^2 \\ &= x(t_0) + \frac{(h_1^2 - h_2^2)x(t_0) + h_2^2 x(t_0 + h_1) - h_1^2 x(t_0 + h_2)}{h_1 h_2 (h_2 - h_1)} (t - t_0) \\ &\quad + \frac{(h_2 - h_1)x(t_0) - h_2 x(t_0 + h_1) + h_1 x(t_0 + h_2)}{h_1 h_2 (h_2 - h_1)} (t - t_0)^2 \end{aligned}$$

and similarly for $p(u(t)) = A(u) + B(u) \cdot (t - t_0) + \frac{1}{2}C(u) \cdot (t - t_0)^2$. The induced action on the coefficients is that induced on $(x(t), u(t))$ so for example

$$\begin{aligned} g \cdot B(u) &= \frac{(h_1^2 - h_2^2)g \cdot u(t_0) + h_2^2 g \cdot u(t_0 + h_1) - h_1^2 g \cdot u(t_0 + h_2)}{h_1 h_2 (h_2 - h_1)} \\ &= \frac{1}{h_1 h_2 (h_2 - h_1)} [(h_1^2 - h_2^2)(\cos \theta(x(t_0) - a) + \sin \theta(u(t_0) - b)) \\ &\quad + h_2^2(\cos \theta(x(t_0 + h_1) - a) + \sin \theta(u(t_0 + h_1) - b)) \\ &\quad - h_1^2(\cos \theta(x(t_0 + h_2) - a) + \sin \theta(u(t_0 + h_2) - b))] \end{aligned}$$

The normalisation equations $g \cdot A(x) = 0$, $g \cdot A(u) = 0$ and $g \cdot B(u) = 0$ yield the frame at Γ to be

$$a = x(t_0), \quad b = u(t_0), \quad \tan \theta = \frac{(h_1^2 - h_2^2)u(t_0) + h_2^2 u(t_0 + h_1) - h_1^2 u(t_0 + h_2)}{(h_1^2 - h_2^2)x(t_0) + h_2^2 x(t_0 + h_1) - h_1^2 x(t_0 + h_2)}$$

In the limit as $h_2 \rightarrow h_1$, we have

$$\tan \theta \rightarrow \frac{u_t(t_0 + h_1) - 2u(t_0 + h_1) + 2u(t_0)}{x_t(t_0 + h_1) - 2x(t_0 + h_1) + 2x(t_0)}$$

and then finally as $h_1 \rightarrow 0$, we have $\tan \theta \rightarrow u_t/x_t$ as expected, indeed, yielding the smooth frame as determined by the limit of the normalisation equations.

If we calculate the invariant matrix $\rho([\Gamma', u, \phi, U])^{-1} \rho([\Gamma, u, \phi, U])$, recalling we have a right action (we are taking ϕ to be the identity map), we obtain the discrete multispace Maurer-Cartan invariants for this frame. Further, $\rho \cdot B(x) \rightarrow (x_t^2 + u_t^2)^{1/2}$ so that we may treat $\rho \cdot B(x)$ as the multispace approximation to the infinitesimal arc length.

In the above example, we used an invariant parameter t to describe the curve $(x(t), u(t))$. If instead we parametrise the curve as $(x, u(x))$, so that the parameter participates in the group action and the operator $\partial/\partial x$ is not invariant, then greater care is required. For example, the group action on the interpolation curve, written as $p(u)(x) = A(x, u) + B(x, u)(x - x_0) + C(x, u)(x - x_0)^2$, looks like $g \cdot p(u)(x) = A(\tilde{x}, \tilde{u}) + B(\tilde{x}, \tilde{u})(x - \tilde{x}_0) + C(\tilde{x}, \tilde{u})(x - \tilde{x}_0)^2$. Solving for the multispace frame in this case seems nontrivial. Such examples will be examined elsewhere.

Now assume we have four lattice points, u_J , $\mathcal{T}_1 u_J$, $\mathcal{T}_2 u_J$ and $\mathcal{T}_2 \mathcal{T}_1 u_J = \mathcal{T}_1 \mathcal{T}_2 u_J$, and that we have two paths connecting u_J with $\mathcal{T}_2 \mathcal{T}_1 u_J = \mathcal{T}_1 \mathcal{T}_2 u_J$ via each of the $\mathcal{T}_1 u_J$ and $\mathcal{T}_2 u_J$ respectively. If we can associate the discrete frame with a multispace frame and differentiate the local syzygy associated with the discrete frame with respect to the path parameters at u_J , we obtain the differential syzygy associated with the multispace frame at u_J . Indeed, let a discrete frame (ρ_J) at the points u_J , $\mathcal{T}_1 u_J$, $\mathcal{T}_2 u_J$ and $\mathcal{T}_1 \mathcal{T}_2 u_J$ be associated with the multispace frame ρ so that $\rho_J = \rho(z_J)$ for some $z_J = [\Gamma_J, \phi, U]$ in the relevant multispace \mathcal{M} , $u_J = \phi(x_J)$. Then, the local syzygy is

$$(22) \quad \mathcal{T}_1(K_2)K_1 = \mathcal{T}_2(K_1)K_2, \quad K_1 = \rho(\mathcal{T}_1 z_J) \rho(z_J)^{-1}, \quad K_2 = \rho(\mathcal{T}_2 z_J) \rho(z_J)^{-1},$$

where $\mathcal{T}_i z_J = [\mathcal{T}_i \Gamma_J, \phi, U]$ and $\mathcal{T}_i \Gamma_J$ is the shift of Γ_J by \mathcal{T}_i , with base point $\mathcal{T}_i x_J$ and $\mathcal{T}_i u_J = \phi(\mathcal{T}_i x_J) = \phi(x_{J+e_i})$. Here we are assuming that the shifts of Γ_J remain within U . Now let $z_J^i(\epsilon_i)$ be paths in the multispace lying within U , with

$z_J^i(0) = z_J$, $z_J^i(1) = \mathcal{T}_i z_J$, $z_J^i(\epsilon_i) = [\Gamma_J^i(\epsilon_i), \phi, U]$ and $u_J^i(\epsilon_i) = \phi(x_J^i(\epsilon_i))$, $x_J^i(\epsilon_i)$ the base point of $\Gamma_J^i(\epsilon_i)$. We denote

$$\mathbf{v} = \frac{d}{d\epsilon_1} \Big|_{\epsilon_1=0} u_J^1(\epsilon_1), \quad \mathbf{w} = \frac{d}{d\epsilon_2} \Big|_{\epsilon_2=0} u_J^2(\epsilon_2),$$

and

$$K_{\mathbf{v}} = \frac{d}{d\epsilon_1} \Big|_{\epsilon_1=0} \rho(z_J^1(\epsilon_1)) \rho(z_J)^{-1}, \quad K_{\mathbf{w}} = \frac{d}{d\epsilon_2} \Big|_{\epsilon_2=0} \rho(z_J^2(\epsilon_2)) \rho(z_J)^{-1}$$

Theorem 4.26. *After differentiating twice, once each with respect to the path parameters ϵ_1 and ϵ_2 and coalescing the lattice of the multispace point z_J to its associated jet point, and under the conditions just stated, the local syzygy, equation (22), becomes the continuous basic syzygy associated to $\rho(z_J)$*

$$(23) \quad \mathcal{D}(\mathbf{w})K_{\mathbf{v}} = \mathcal{D}(\mathbf{v})K_{\mathbf{w}} + [\mathcal{D}(\mathbf{w}), \mathcal{D}(\mathbf{v})] \rho \rho^{-1} + [K_{\mathbf{v}}, K_{\mathbf{w}}]$$

Proof. The core of the proof is standard. We assume a matrix representation of the frame, and note that Taylor's Theorem is valid.

To ease the notation, we set $\epsilon_1 = h$ and $\epsilon_2 = k$, and simplify to where $\mathbf{v} = d/dh|_{h=0} \phi(x_0 + h\mathbf{e}_1)$ and $\mathbf{w} = d/dk|_{k=0} \phi(x_0 + k\mathbf{e}_2)$ in local coordinates, so that we evaluate our frame at the multispace elements with lattice basepoints at $u_J = \phi(x_0)$, $\mathcal{T}_1 u_J = u_{J,1} = \phi(x_0 + h\mathbf{e}_1)$, $\mathcal{T}_2 u_J = u_{J,2} = \phi(x_0 + k\mathbf{e}_2)$ and $\mathcal{T}_2 \mathcal{T}_1 u_J = u_{J,1,2} = u_{J,2,1} = \phi(x_0 + h\mathbf{e}_1 + k\mathbf{e}_2)$. We denote the partial derivative operators $\partial/\partial\epsilon_i$ as ∂_i .

Denoting $\mathcal{T}_i \rho_J = \rho_{J,i}$ we have

$$K_{J,1} = \rho_{J,1} \rho_J^{-1}, \quad K_{J,2} = \rho_{J,2} \rho_J^{-1},$$

and for sufficiently small h and k , there will exist, dropping the index J for clarity, matrices X_1 and X_2 in the Lie algebra of G such that

$$K_1 = \exp(hX_1 + \mathcal{O}(h^2)), \quad K_2 = \exp(kX_2 + \mathcal{O}(k^2)).$$

We have

$$(24) \quad \partial_1 \rho \cdot \rho^{-1} = X_1, \quad \partial_2 \rho \cdot \rho^{-1} = X_2$$

and

$$\mathcal{T}_2 X_1 = X_1 + k\partial_2 X_1 + \mathcal{O}(h^2), \quad \mathcal{T}_1 X_2 = X_2 + k\partial_2 X_2 + \mathcal{O}(k^2).$$

Then

$$\begin{aligned} \mathcal{T}_2 K_1 &= \exp(h\mathcal{T}_2 X_1 + \mathcal{O}(h^2)) \\ &= \exp(hX_1 + hk\partial_2 X_1 + \mathcal{O}(k^2, h^2)) \\ \mathcal{T}_1 K_2 &= \exp(k\mathcal{T}_1 X_2 + \mathcal{O}(k^2)) \\ &= \exp(kX_2 + hk\partial_1 X_2 + \mathcal{O}(k^2, h^2)) \end{aligned}$$

Applying the Baker Campbell Haussdorff formula, [57],

$$\log(\exp(X)\exp(Y)) = X + Y + \frac{1}{2}[X, Y] + \text{higher order brackets},$$

we have

$$\begin{aligned} \log(\mathcal{T}_2 K_1 \cdot K_2) &= hX_1 + hk\partial_2 X_1 + kX_2 + \frac{1}{2}[hX_1 + hk\partial_2 X_1, kX_2] + \mathcal{O}(k^2, h^2) \\ &= hX_1 + kX_2 + hk(\partial_2 X_1 + \frac{1}{2}[X_1, X_2]) + \mathcal{O}(k^2, h^2), \\ \log(\mathcal{T}_1 K_2 \cdot K_1) &= kX_2 + hk\partial_1 X_2 + hX_1 + \frac{1}{2}[kX_2 + hk\partial_1 X_2, hX_1] + \mathcal{O}(k^2, h^2) \\ &= hX_1 + kX_2 + hk(\partial_1 X_2 + \frac{1}{2}[X_2, X_1]) + \mathcal{O}(k^2, h^2). \end{aligned}$$

Equating the two formulae by imposing the local syzygy, differentiating with respect to both h and k and then sending $h, k \rightarrow 0$, yields, after a slight rearrangement,

$$(25) \quad \partial_2 X_1 - \partial_1 X_2 = [X_2, X_1].$$

Finally, we need to rewrite Equation (25) in terms of the invariant differential operators. The formula given in Equation (21) shows that the partial derivatives ∂_i can be written as a linear sum of the invariant operators with invariant coefficients. We must then back substitute for the ∂_i , including rewriting the $X_i = \partial_i \rho \cdot \rho^{-1}$ in terms of the $\mathcal{D}_j \rho \cdot \rho^{-1}$. The final result yields the extra terms in the case that the invariant operators do not commute. \square

Example 4.27. *We conclude the running example, 4.20. We set the points (x_i, y_i) , $i = 1, 2, 3$ to be $(x_1, y_1) = (x_0 + \bar{h}, y_0)$, $(x_2, y_2) = (x_0, y_0 + \bar{k})$ and $(x_3, y_3) = (x_0 + \bar{h}, y_0 + \bar{k})$. We then calculate the four matrices, $K_{10} = \rho_1 \rho_0^{-1} = \exp(X_{10})$, $K_{31} = \rho_3 \rho_1^{-1} = \exp(X_{31})$, $K_{20} = \rho_2 \rho_0^{-1} = \exp(X_{20})$ and $K_{32} = \rho_3 \rho_2^{-1} = \exp(X_{32})$. Direct calculation gives, setting $\Delta_1 F = F(x_0 + h, y_0) - F(x_0, y_0)$ for a function $F \in \{u, u_x, u_y, u_{xy}\}$ in the formulae to ease the notation,*

$$\begin{aligned} X_{10} &= -\bar{h} \begin{pmatrix} \frac{\Delta_1 u_x}{\Delta_1 u} & \frac{u_x(x_0, y_0)}{\Delta_1 u} \\ 0 & 0 \end{pmatrix} + \mathcal{O}(\bar{h}^2) \\ X_{20} &= -\bar{k} \begin{pmatrix} \frac{\Delta_1 u_y}{\Delta_1 u} & \frac{u_y(x_0, y_0)}{\Delta_1 u} \\ 0 & 0 \end{pmatrix} + \mathcal{O}(\bar{k}^2) \\ X_{31} &= X_{20} - \widetilde{h\bar{k}X_{31}} + \mathcal{O}(\bar{h}^2, \bar{k}^2) \\ &= X_{20} - \bar{h}\bar{k} \begin{pmatrix} \frac{\Delta_1 u_x \Delta_1 u_y + \Delta_1 u_{xy} \Delta_1 u}{(\Delta_1 u)^2} & h \frac{u_y \Delta_1 u_x + u_{xy} \Delta_1 u}{(\Delta_1 u)^2} \\ 0 & 0 \end{pmatrix} + \mathcal{O}(\bar{h}^2, \bar{k}^2) \\ X_{32} &= X_{10} - \widetilde{h\bar{k}X_{32}} + \mathcal{O}(\bar{h}^2, \bar{k}^2) \\ &= X_{10} - \bar{h}\bar{k} \begin{pmatrix} \frac{\Delta_1 u_x \Delta_1 u_y + \Delta_1 u_{xy} \Delta_1 u}{(\Delta_1 u)^2} & h \frac{u_x \Delta_1 u_y + u_{xy} \Delta_1 u}{(\Delta_1 u)^2} \\ 0 & 0 \end{pmatrix} + \mathcal{O}(\bar{h}^2, \bar{k}^2) \end{aligned}$$

where this defines $\widetilde{X_{31}}$ and $\widetilde{X_{32}}$. The local syzygy is $K_{31}K_{10} = K_{32}K_{20}$, and applying the Baker Campbell Hausdorff formula to this yields

$$X_{10} + X_{31} + \frac{1}{2}[X_{31}, X_{10}] = X_{32} + X_{20} + \frac{1}{2}[X_{32}, X_{20}] + \mathcal{O}(\bar{h}^2, \bar{k}^2).$$

The equation for the lower order terms simplifies to

$$(26) \quad \widetilde{X_{32}} - \widetilde{X_{31}} = [X_{10}, X_{20}].$$

This last equation is straightforward to verify. Finally, taking the limit as $h \rightarrow 0$, Equation (26) yields the differential syzygy for ρ evaluated on the jet bundle,

$$\frac{\partial}{\partial y} \rho_x \rho^{-1} - \frac{\partial}{\partial x} \rho_y \rho^{-1} = [\rho_y \rho^{-1}, \rho_x \rho^{-1}],$$

where recall ρ on the jet bundle is given in Equation (17).

Similar relationships exist when we take limits only in one of the variables, producing an evolution of discrete submanifolds. For example, if $p = 2$ and S has coordinates (x, y) , then if $K_{J,2}^i = \rho_{J+e_2}(z_J^i)\rho(z_J)^{-1}$ and $N_J = \frac{d}{dx}(K_J)K_J^{-1}$, then when we take limits in the calculation in the proof of the theorem as $x_{J+e_1} \rightarrow x_J$ we have

$$\frac{d}{dx}(K_{(J,2)}) = N_{J+e_2} - K_{(J,2)}N_JK_{(J,2)}^{-1}$$

which is a mixed syzygy that often appears describing invariant evolutions of polygons in terms of coordinates in their moduli spaces, as in [41] and [46]. Among these evolutions one often finds completely integrable discretizations of well known completely integrable PDEs. These results are really key to some of the applications in our next section.

5. APPLICATIONS

5.1. Application to the design of a variational numerical scheme for a shallow water system. This example is motivated by the need for finite difference versions of variational shallow water problems which are invariant under the so-called particle relabelling symmetry. We consider the base space to have coordinates (a, b, t) , where (a, b) is the fluid particle label at time $t = 0$. The two dimensional dependent variable space is $(x, y) = (x(a, b, t), y(a, b, t))$, which is the position of the fluid particle at time t , so that $(x(a, b, 0), y(a, b, 0)) = (a, b)$. The particle relabelling action is given by

$$g \cdot a = A(a, b), \quad g \cdot b = B(a, b), \quad A_a B_b - A_b B_a = 1$$

together with $g \cdot x = x$, $g \cdot y = y$ and $g \cdot t = t$. It can be seen that the particle relabelling group is the group of area preserving diffeomorphisms of the (a, b) plane (or at least the domain of interest in the (a, b) plane). Further, it is known that the invariants of this group action are $\Delta = x_a y_b - x_b y_a$ and its derivatives under the invariant differential operators,

$$\frac{\partial}{\partial x} = \frac{y_b}{\Delta} \frac{\partial}{\partial a} - \frac{y_a}{\Delta} \frac{\partial}{\partial b}, \quad \frac{\partial}{\partial y} = -\frac{x_b}{\Delta} \frac{\partial}{\partial a} + \frac{x_a}{\Delta} \frac{\partial}{\partial b}.$$

The aim is to design a multispace version of the Lagrangian for variational shallow water problems, which have the form,

$$(27) \quad \mathcal{L}[x, y] = \int L(x, y, x_t, y_t, x_a y_b - x_b y_a) da db dt.$$

This family of Lagrangians are each invariant under translations in time, translations in both a and b , rotations in the (a, b) plane, and more generally, the full particle relabelling group. Noether's Theorem [48, 1, 52] then yields conservation of energy, linear and angular momenta, and potential vorticity.

If we take the simplest corner lattice with base point (a_0, b_0, t_0) to be

$$\Gamma = \{(a_0, b_0, t_0), (a_1, b_1, t_0), (a_2, b_2, t_0), (a_3, b_3, t_1)\}$$

then the (linear) interpolation of x is given by

$$x(a, b, t) \sim x(a_0, b_0, t_0) + \mathcal{M}(x_a)(a - a_0) + \mathcal{M}(x_b)(b - b_0) + \mathcal{M}(x_t)(t - t_0)$$

where this defines the coefficients $\mathcal{M}(x_K)$, and we have

$$\mathcal{M}(x_a) = \frac{1}{\mathcal{A}} \begin{vmatrix} 1 & x(a_0, b_0, t_0) & b_0 \\ 1 & x(a_1, b_1, t_0) & b_1 \\ 1 & x(a_2, b_2, t_0) & b_2 \end{vmatrix}, \quad \mathcal{M}(x_b) = \frac{1}{\mathcal{A}} \begin{vmatrix} 1 & a_0 & x(a_0, b_0, t_0) \\ 1 & a_1 & x(a_1, b_1, t_0) \\ 1 & a_2 & x(a_2, b_2, t_0) \end{vmatrix}$$

and

$$\mathcal{M}(x_t) = \frac{1}{(t_1 - t_0)\mathcal{A}} \begin{vmatrix} 1 & a_0 & b_0 & x(a_0, b_0, t_0) \\ 1 & a_1 & b_1 & x(a_1, b_1, t_0) \\ 1 & a_2 & b_2 & x(a_2, b_2, t_0) \\ 1 & a_3 & b_3 & x(a_3, b_3, t_2) \end{vmatrix}$$

where

$$\mathcal{A} = \begin{vmatrix} 1 & a_0 & b_0 \\ 1 & a_1 & b_1 \\ 1 & a_2 & b_2 \end{vmatrix}$$

is the area, $|(a_1 - a_0, b_1 - b_0) \wedge (a_2 - a_0, b_2 - b_0)|$. The interpolant for y is similar, with $y(a_i, b_i, t_i)$ replacing $x(a_i, b_i, t_i)$ in the above formulae.

We consider the Lie group $SL(2) \times \mathbb{R}^2$ acting on the (a, b) plane as the standard (right) equiaffine action,

$$\begin{pmatrix} a \\ b \end{pmatrix} \mapsto \begin{pmatrix} g \cdot a \\ g \cdot b \end{pmatrix} = \begin{pmatrix} \delta & -\beta \\ -\gamma & \alpha \end{pmatrix} \begin{pmatrix} a - \epsilon_1 \\ b - \epsilon_2 \end{pmatrix}$$

so that $(\epsilon_1, \epsilon_2) \in \mathbb{R}^2$ is the translation vector, and $\alpha\delta - \beta\gamma = 1$, and that $g \cdot x = x$, $g \cdot y = y$ and $g \cdot t = t$. This group is contained within the particle labelling symmetry group, and is just big enough to obtain the area invariant, which we do next.

The induced action on the coefficients in the interpolants is given by, for example,

$$g \cdot \mathcal{M}(x_a) = \frac{1}{\mathcal{A}} \begin{vmatrix} 1 & x(a_0, b_0, t_0) & g \cdot b_0 \\ 1 & x(a_1, b_1, t_0) & g \cdot b_1 \\ 1 & x(a_2, b_2, t_0) & g \cdot b_2 \end{vmatrix},$$

noting that $g \cdot \mathcal{A} = \mathcal{A}$, indeed, \mathcal{A} is an invariant as is easily seen.

We take the normalisation equations $g \cdot (a_0, b_0) = (0, 0)$, $g \cdot \mathcal{M}(x_a) = 1$, $g \cdot \mathcal{M}(x_b) = 0$ and $g \cdot \mathcal{M}(y_a) = 0$. Then the multispace frame $\mathcal{M}(\rho)(\Gamma)$ is $(\epsilon_1, \epsilon_2) = (a_0, b_0)$ and

$$\begin{pmatrix} \delta & -\beta \\ -\gamma & \alpha \end{pmatrix} = \begin{pmatrix} \mathcal{M}(x_a) & \mathcal{M}(x_b) \\ \frac{\mathcal{M}(y_a)}{\mathcal{M}(\Delta)} & \frac{\mathcal{M}(y_b)}{\mathcal{M}(\Delta)} \end{pmatrix}$$

where $\mathcal{M}(\Delta) = \mathcal{M}(x_a)\mathcal{M}(y_b) - \mathcal{M}(x_b)\mathcal{M}(y_a)$. Evaluating the remaining coefficient on the frame, we obtain the invariant,

$$\begin{aligned} I(\mathcal{M}(y_b)) &= g \cdot \mathcal{M}(y_b) \Big|_{\rho} \\ &= \mathcal{M}(\Delta) \\ &= \frac{1}{\mathcal{A}} \begin{vmatrix} 1 & x(a_0, b_0, t_0) & y(a_0, b_0, t_0) \\ 1 & x(a_1, b_1, t_1) & y(a_1, b_1, t_1) \\ 1 & x(a_2, b_2, t_2) & y(a_2, b_2, t_2) \end{vmatrix} \end{aligned}$$

Calculating the continuum limit of $\mathcal{M}(\Delta)$ we obtain $x_a y_b - x_b y_a$, which is Δ , the area invariant, as expected. Further, the continuum list of the frame is

$$\rho \rightarrow \begin{pmatrix} x_a & x_b \\ \frac{y_a}{x_a y_b - x_b y_a} & \frac{y_b}{x_a y_b - x_b y_a} \end{pmatrix}$$

This is the smooth frame obtained with the smooth limit of the normalisation equations, that is, $\{\tilde{x}_a = 1, \tilde{x}_b = 0, \tilde{y}_a = 0\}$.

We observe that both $\mathcal{M}(x_t)$ and $\mathcal{M}(y_t)$ are invariant under the equiaffine action.




FIGURE 17. Length one lattices stacked as a mesh, for a finite difference variational problem. Shown here is single time slice for the shallow water problem.

Thus, we propose the multispace analogue of the Lagrangian (27) to be

$$(28) \quad \mathcal{M}(\mathcal{L}[x, y]) = \sum_{\Gamma} L(x, y, \mathcal{M}(x_t), \mathcal{M}(y_t), \mathcal{M}(\Delta)) \mathcal{A}(t_1 - t_0)$$

where the sum is over all corner lattices stacked into a mesh, as in Figure 17. The factor $\mathcal{A}(t_1 - t_0)$ is the multispace approximation of the volume form, $dadbdt$, and is needed to obtain the correct continuum limits for the conservation laws for energy and the linear momenta.

Finite difference Euler Lagrange equations and Noether's conservation laws can be calculated in the standard way [32, 23]; the details and the results of this calculation will be explored elsewhere. Here we simply demonstrate that the multispace Lagrangian, (28) is invariant under a discrete analogue of the particle labelling symmetry. Indeed, looking at Figure 17, one can use a different element of $SL(2) \times \mathbb{R}^2$ on each corner lattice, provided that certain consistency conditions hold, specifically, that if a vertex is in the intersection of two lattices, that their image under the two different group elements is the same. The Lagrangian (28) is clearly invariant under this discretisation of the particle relabelling group, the discretisation being a subgroup of $(SL(2) \times \mathbb{R}^2)^R$ where R is the number of corner lattices on a time slice of the mesh. Finally, if one takes the collection of frames consisting of the multispace frame restricted to each vertex, then we have a discrete moving frame, using which the discrete Euler Lagrange equations and discrete conservation laws can be studied [53].

5.2. Discretizations of completely integrable PDEs. The geometry of curves and surfaces have been linked to integrable systems repeatedly in the literature, see [24], [56], [44], for example. A drawback of the application of the results in this paper to finding completely integrable discretizations of completely integrable systems is that one needs to choose a type of approximation (forward, backwards, linear or higher order, etc) a priori to find the limit. On the other hand, any two choices of discrete moving frames (be the one associated to a certain type of limit or any different one) will always be associated by a gauge transformation. This means that if one finds a discrete integrable system associated to any given choice of moving frame, one might be able to relate it to a different choice and perhaps link it to the continuous case. This was done in [46], where the authors found discrete integrable systems that were the discretization of Adler-Gel'fand-Dikii integrable evolutions, both of them linked to the projective geometry of curves and polygons. The authors of [46] also found a way to obtain two Hamiltonian structures associated to the discrete system through a reduction process, a process that was later extended to other semisimple homogeneous spaces in [45]. Different approaches were used, for example in [2] and in other works of these authors to construct completely integrable discretizations of integrable systems with the use of lattice models in Euclidean, projective and conformal geometry. Their approach is quite different from the one used in [46] and in this paper, in that they choose lattices with different geometric properties to achieve discretizations. The following are just some examples of the connection between continuous and discrete models, in the equi-centro affine and the Euclidean space. Both of these examples use mixed multispace discrete/continuous models, and although the first example only uses discrete coordinates to construct the multispace moving frame, the second example will make full use of both.

5.2.1. Integrable discretizations of Boussinesq equations. In this example we make use of mixed differential/difference coordinates in the multispace. For reasons that will become clear later, we will also assume that the lattice variety has a monodromy in the discrete variable (a global property). That is, $u_{n+T}(t) = \mathbf{m} \cdot u_n(t)$ for any $n \in \mathbb{Z}$, with T the period and some monodromy $\mathbf{m} \in \text{SL}(3, \mathbb{R})$. This ensures that the invariants will be T -periodic in n .

First we describe the situation when we are in continuous jets with two parameters (x, t) . It is well known (see for example [9]) that the Boussinesq equations

$$(q_0)_t + \frac{1}{6}q_1''' + \frac{2}{3}q_1q_1' = 0, \quad (q_1)_t - 2q_0' = 0$$

can be obtained as the evolution induced on equi-centro-affine curvatures by a certain evolution of curves. Let our manifold be \mathbb{R}^3 with $M = \text{SL}(3, \mathbb{R})$ acting linearly on it. Within M consider parametrized surfaces on (x, t) . We will define the following cross-section:

$$(29) \quad \rho \cdot u = e_3, \quad \rho \cdot u' = e_2, \quad \rho \cdot u'' = de_1$$

where the prime denotes d/dx and e_i are the standard unit vectors in \mathbb{R}^3 . Clearly $d = \det(u, u', u'')$. This defines uniquely a *right* moving frame whose *left* companion is given by

$$\rho^{-1} = \left(\frac{1}{d}u'', u', u\right).$$

Let us assume that

$$(30) \quad d = \det(u'', u', u) = 1,$$

that is we will parametrize the surface so that the curves associated with t fixed are parametrized by the equi-centro-affine arc-length (these curves are in one-to-one correspondence with projective curves, see [9]). In that case the left x -Maurer–Cartan matrix associated to it is given by

$$Q = \rho(\rho^{-1})' = \begin{pmatrix} 0 & 1 & 0 \\ k_1 & 0 & 1 \\ k_2 & 0 & 0 \end{pmatrix}$$

where $u''' = k_1 u' + k_2 u$. Next we will gauge this frame to a different left frame $\widehat{\rho} = \rho^{-1}g$ by the element

$$g = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -k_1 & 0 & 1 \end{pmatrix}.$$

The resulting x -Maurer–Cartan matrix is given by

$$K = g^{-1}g' + g^{-1}Qg = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ b & a & 0 \end{pmatrix}$$

where $a = k_1$ and $b = k_2 - k_1'$. Gauging the system can be seen as changing the coordinates, the results can always be gauged back to the original setting.

We will next consider the syzygy

$$(31) \quad u_t - u'' + \frac{2}{3}au = 0$$

which describes a precisely chosen evolutionary equation for curves whose flow will be tracing our parametrized surface. With this condition, the left t -Maurer–Cartan matrix is easily seen to be given by

$$N = \widehat{\rho}^{-1}\widehat{\rho}'_t = \begin{pmatrix} -(w_1 + 1/3a) & w_0 & 1 \\ v_1 & w_1 & 0 \\ v_2 & w_2 & \frac{1}{3}a \end{pmatrix}$$

for some entries v_i, w_i . The local basic syzygies (or the compatibility condition between x and t) are given by

$$K_t = N_x + [K, N]$$

and they can be used to solve for N so that

$$w_0 = 0, \quad w_1 = \frac{1}{3}a, \quad w_2 = b + \frac{1}{3}a', \quad v_1 = b + \frac{2}{3}a', \quad v_2 = -b' - \frac{2}{3}a''.$$

We can further find two more syzygies given by

$$a_t - 2b' - a'' = 0, \quad b_t + \frac{2}{3}a''' + b'' - \frac{2}{3}a'a.$$

This system of equations is equivalent to the Boussinesq equation. Indeed, if $a = q_1$ and $b = \frac{1}{2}q_1' - q_0$, we have

$$(q_0)_t + \frac{1}{6}q_1''' + \frac{2}{3}q_1q_1' = 0, \quad (q_1)_t - 2q_0' = 0$$

which is the standard Boussinesq equation.

Assume now that we move in the multispace away from a continuous jet to a mixed discrete/continuous multispace submanifold, where x is now discrete and t is continuous. Let us choose lattices containing (x_n, t) , (x_{n+1}, t) , (x_{n+2}, t) and such that $x_{r+1} - x_r = \Delta x_r = c$ is constant for any r . That is, we will restrict to lattices with sides of equal length. The cross-section (29) evaluated on lattices of this form will be given by

$$\rho \cdot u_n = e_3, \quad \rho \cdot (u_{n+1} - u_n)/c = e_2, \quad \rho \cdot \left(\frac{1}{c^2}(u_{n+2} - 2u_{n+1} + u_n)\right) = e_1$$

where $a = (1/c^3) \det(u_{n+2}, u_{n+1}, u_n)$ and $u_i = u(x_i, t)$. The left moving frame associated to this cross section is given by

$$\begin{aligned} \rho^{-1} &= \left(\frac{1}{c^2}(u_{n+2} - 2u_{n+1} + u_n), \frac{1}{c}(u_{n+1} - u_n), u_n\right) \\ &= (u_{n+2} \quad u_{n+1} \quad u_n) \begin{pmatrix} c^{-2} & 0 & 0 \\ -2c^{-2} & c^{-1} & 0 \\ c^{-2} & -c^{-1} & 1 \end{pmatrix} \end{aligned}$$

which can clearly be gauged to

$$\eta = (u_{n+2}, u_{n+1}, u_n).$$

The multispace subspace (30), when restricted to our partially coalesced lattices, becomes

$$(32) \quad \det(u_{n+2}, u_{n+1}, u_n) = c^3$$

for all n . Let us introduce one last gauge by the matrix

$$g = \begin{pmatrix} 1 & 0 & 0 \\ -a_{n-1} & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

If $u_{n+3} = k_1^n u_{n+2} + k_2^n u_{n+1} + u_n$, then, the discrete x -Maurer-Cartan matrix associated to $\rho = \eta g$ is given by

$$K_n = \begin{pmatrix} 0 & 1 & 0 \\ b_n & a_n & 1 \\ 1 & 0 & 0 \end{pmatrix}$$

where $b_n = k_2^n$ and $a_n = k_1^{n-1}$. As before, the t -Maurer-Cartan matrix is given by

$$N_n = \widehat{\rho}_n^{-1}(\widehat{\rho}_n)_t = \begin{pmatrix} -(w_1^n + r_2^n) & w_0^n & r_0^n \\ v_1^n & w_1^n & r_1^n \\ v_2^n & w_2^n & r_2^n \end{pmatrix}$$

and the local basic syzygy is

$$K_n^{-1}(K_n)_t = N_{n+1} - K_n^{-1}N_nK_n.$$

This syzygy solves for N_n in terms of r_i

$$\begin{aligned} w_0^n &= r_1^{n+1}; & w_1^n &= r_2^{n+1} + a_n r_1^{n+1} + b_n r_0^{n+1}; & w_2^n &= r_0^{n+1}; \\ v_1^n &= r_0^{n-1} + b_{n-1} r_1^n; & v_2^n &= r_1^{n+2} - a_n r_0^{n+1}, \end{aligned}$$

and it provides the condition for preserving the restriction to the multispace submanifold (32), namely

$$(33) \quad r_2^{n+2} + r_2^{n+1} + r_2^n + a_{n+1} r_1^{n+2} + b_{n+1} r_0^{n+2} + b_n r_0^{n+1} = 0.$$

If the map $r_2^n \rightarrow r_2^{n+2} + r_2^{n+1} + r_2^n = (\tau^2 + \tau + 1)r_2^n$ is invertible (which is true if $N \neq 3s$ for any s as shown in [46]), this condition solves for r_2^s in terms of r_1 and r_0 . The syzygy also describes $(a_n)_t$ and $(b_n)_t$ to satisfy

$$\begin{aligned} (a_n)_t &= (1 + a_n b_{n+1})r_0^{n+2} - (a_n b_n r_0^{n+1} + r_0^{n-1}) \\ &\quad + (b_n + a_n a_{n+1})r_1^{n+2} - (b_{n-1}r_1^n + a_n^2 r_1^{n+1}) + a_n r_2^{n+2} - a_n r_2^n \\ (b_n)_t &= r_1^{n+3} - (b_n a_{n+1} r_1^{n+2} + r_1^n) - (a_{n+1} + b_n b_{n+1})r_0^{n+2} \\ &\quad - b_n^2 r_0^{n+1} + a_n r_0^n - (b_n r_2^{n+2} + 2b_n r_2^{n+1}). \end{aligned}$$

If further we impose the syzygy

$$(34) \quad (u_n)_t + \frac{1}{b_{n-1}} (u_{n+2} - a_n u_{n+1}) + \frac{2}{3} u_n = 0$$

then we can see that $r_0^n = \frac{1}{b_{n-1}}$, $r_1^n = 0$ and $r_2^n = -(\tau^2 + \tau + 1)^{-1}(\tau + 1)\frac{b_n}{b_n} = -\frac{2}{3}$ is the solution of (33) for these choices. Then

$$\begin{aligned} (a_n)_t &= \frac{1}{b_{n+1}} - \frac{1}{b_{n-2}} \\ (b_n)_t &= -\frac{a_{n+1}}{b_{n+1}} + \frac{a_n}{b_{n-1}} \end{aligned}$$

and the changes

$$\alpha_n = -\frac{1}{b_n b_{n+1} b_{n+2}}, \quad \beta_n = -\frac{a_{n+1}}{b_n b_{n+1}}$$

transform this equation in the integrable discretization of Boussinesq

$$\begin{aligned} (\alpha_n)_t &= \alpha_n(\beta_{n+2} - \beta_{n-1}) \\ (\beta_n)_t &= \alpha_{n-1} - \alpha_n + \beta_n(\beta_{n+1} - \beta_{n-1}). \end{aligned}$$

This system appears in [25].

It is not clear to us how to systematically connect integrable discrete systems and evolutions of polygons as given by (34). In the continuous case there is a general link between Hamiltonian evolutions at the level of the invariants and evolution of curves on geometric manifolds which are homogeneous of the form G/H or $(G \rtimes \mathbb{R}^n)/G$ with G semisimple (see [42], [43]), but the situation in the discrete case is not so clear in general. In particular, the syzygies (31) and (34) are not the restriction of the same syzygy on different points in the multispace, even when we account for all the different changes introduced by gauges. Still, it is widely known that certain evolutions of polygons result in completely integrable discrete systems (see, for example, [3] in the Euclidean case and [2] in more cases with further restrictions on the lattices). The multispace allows us to construct geometrically without the need to account for the limits. We include one more example along these lines and further use of multispace in this area will appear elsewhere.

5.2.2. Discretization of the Sine-Gordon equation. It is well known that the Codazzi–Mainardi equations for Euclidean surfaces in \mathbb{R}^3 with constant negative Gauss curvature includes a Sine-Gordon equation, a well known completely integrable system, that decouples from the rest of the determining equations for the surface. The Codazzi–Mainardi equations are simply syzygies for a well-chosen moving frame, hence using the multispace framework we will be able to find a discretization of the Sine-Gordon equation with strong geometric meaning as determining mixed lattice/surfaces with negative Gauss curvature. It is not clear to us if the discretization

below is completely integrable as it becomes part of a system of equations defining the lattice/surface, rather than decoupling to discretize Sine-Gordon individually. A discrete Sine-Gordon equation on lattices was also defined in [2], although the conditions that the authors imposed on their lattices are not impose here. Further study on the connection between both approaches will appear elsewhere.

We review the continuous case first.

5.2.3. *Sine-Gordon as syzygy of Euclidean surfaces with constant negative curvature.* Let G be the Euclidean group represented as the subgroup of $GL(4, \mathbb{R})$

$$(35) \quad g = \begin{pmatrix} 1 & 0 \\ v & \Theta \end{pmatrix}$$

where $v \in \mathbb{R}^3$ and $\Theta \in SO(3)$. The group acts in \mathbb{R}^3 with the standard action $g \cdot u = \Theta u + v$ which coincides with the one induced by $g \begin{pmatrix} 1 \\ u \end{pmatrix}$. Let $u(x, y)$ be a parametrized surface and assume that x and y are normalized to measure the arc-length in the x and y direction. That is, assume that $\|u_x\| = \|u_y\| = 1$. Let us define a moving frame through the normalizations

$$g \cdot u = 0, \quad g \cdot u_x = e_1, \quad g \cdot u_y = \cos \alpha e_1 + \sin \alpha e_2$$

where α is the angle formed by u_x and u_y . Solving the equations we obtain that $v = -\Theta u$ and

$$\Theta^T = \left(u_x \quad \frac{1}{\sin \alpha} (u_y - \cos \alpha u_x) \quad N \right)$$

where $N = \frac{1}{\sin \alpha} (u_x \times u_y)$ is the standard normal unit vector determined by the parametrization. Using the traditional notation $u_{xx} = \Gamma_{11}^1 u_x + \Gamma_{11}^2 u_y + eN$, $u_{xy} = \Gamma_{12}^1 u_x + \Gamma_{12}^2 u_y + fN$, $u_{yy} = \Gamma_{22}^1 u_x + \Gamma_{22}^2 u_y + gN$, we can write the Maurer–Cartan matrices as

$$\widehat{N} = \rho_x \rho^{-1} = \begin{pmatrix} 0 & 0 \\ -(\Theta u)_x & \Theta_x \end{pmatrix} \begin{pmatrix} 1 & 0 \\ u & \Theta^T \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ -e_1 & \Theta_x \Theta^T \end{pmatrix}$$

$$\widehat{K} = \rho_y \rho^{-1} = \begin{pmatrix} 0 & 0 \\ -\cos \alpha e_1 - \sin \alpha e_2 & \Theta_y \Theta^T \end{pmatrix}$$

where

$$(36) \quad N = \Theta_x \Theta^T = \begin{pmatrix} 0 & \sin \alpha \Gamma_{11}^2 & e \\ -\sin \alpha \Gamma_{11}^2 & 0 & \frac{1}{\sin \alpha} (f - \cos \alpha e) \\ -e & \frac{-1}{\sin \alpha} (f - \cos \alpha e) & 0 \end{pmatrix}$$

$$(37) \quad K = \Theta_y \Theta^T = \begin{pmatrix} 0 & \sin \alpha \Gamma_{12}^2 & f \\ -\sin \alpha \Gamma_{12}^2 & 0 & \frac{1}{\sin \alpha} (g - \cos \alpha f) \\ -f & \frac{-1}{\sin \alpha} (g - \cos \alpha f) & 0 \end{pmatrix}.$$

Substituting these values in the local syzygy $\widehat{K}_x = \widehat{N}_y + [\widehat{K}, \widehat{N}]$ and selecting the \mathbb{R}^3 component, we get the equation

$$0 = \begin{pmatrix} -\cos \alpha \\ -\sin \alpha \\ 0 \end{pmatrix}_x - \begin{pmatrix} 0 \\ -\sin \alpha \Gamma_{12}^2 \\ -f \end{pmatrix} + \cos \alpha \begin{pmatrix} 0 \\ -\sin \alpha \Gamma_{11}^2 \\ -e \end{pmatrix} + \sin \alpha \begin{pmatrix} \sin \alpha \Gamma_{11}^2 \\ 0 \\ \frac{-1}{\sin \alpha} (f - \cos \alpha e) \end{pmatrix}.$$

The last entry is trivial, the first solves for the value

$$\Gamma_{11}^2 = \frac{-1}{\sin \alpha} \alpha_x,$$

while the second one simplifies to $\Gamma_{12}^2 = 0$, whenever $\sin \alpha \neq 0$. The $\text{SO}(3)$ portion of the syzygy is given by

$$(38) \quad K_x = N_y + [K, N].$$

If we write down the equation that does not involve derivatives of the second fundamental form (the equations defined by the (1, 2) entry), we have

$$(\sin \alpha \Gamma_{11}^2)_y = \frac{-f}{\sin \alpha} (f - \cos \alpha e) + \frac{e}{\sin \alpha} (g - \cos \alpha f),$$

which becomes

$$\alpha_{xy} = -\frac{1}{\sin \alpha} (eg - f^2).$$

If \mathbf{K} is the Gauss curvature, we know that $\mathbf{K} = \frac{eg-f^2}{EG-F^2}$, where $E = \|u_x\| = G = \|u_y\| = 1$ and $F = u_x \cdot u_y = \cos \alpha$. Thus, the equation becomes

$$\alpha_{xy} = -\sin \alpha \mathbf{K}$$

which is the Sine–Gordon equation whenever \mathbf{K} is constant and negative.

Notice that this equation is not enough to determine the surface. Indeed, solving for α only determines the first fundamental form (or metric), and the knowledge of the Gauss curvature does not suffice to determine the second fundamental form. Indeed, one would need two more equations to do so, given by the two remaining entries (1, 3) and (2, 3) of the $\text{SO}(3)$ portion of the local syzygy, i.e. (38). Thus, the surface is determined upon solving a system of 3 equations, one of which decouples and is equal to Sine–Gordon.

5.2.4. *A discrete Sine–Gordon evolution as a syzygy of a mixed difference-discrete lattice.* Assume we have a smooth family of polygons, or a mixed (1, 1)-lattice (one continuous direction and one discrete one) of the form y_r, y_{r+1} in the y direction and continuous 1-jet in the x direction. As far as we use the same multispace cross-section, we will have guaranteed that discrete invariant data approximates the continuous one. Thus, consider the transverse section

$$g \cdot u_r = g \cdot u(x, y_r) = 0; \quad g \cdot_p u_x = \|u_x\| e_1; \quad \frac{1}{y_1 - y_0} (g \cdot \mathcal{T}u_r - g \cdot u_r) = \frac{\|\Delta u_r\|}{\Delta y_0} w_r$$

where g is as in (35); $\mathcal{T}u_r = u_{r+1} = u(x, \mathcal{T}y_r) = u(x, y_{r+1})$; \cdot_p is the prolonged action given by $g \cdot_p u_x = \Theta u_x$; and w_r is a unit vector with $w_r = \cos \alpha_n e_1 + \sin \alpha_n e_2$ where $\alpha_n = \alpha_n(x)$ is the angle between u_x and Δu_r , $\Delta u_r = (\tau - 1)u_r$. From here, the multispace cross section defines $v_r = -\Theta_r u_r$, $\Theta_r u_x = \|u_x\| e_1$ and $\Theta_r \Delta u_r = \|\Delta u_r\| w_r$.

With these choices the right moving frame becomes

$$\rho_r = \begin{pmatrix} 1 & 0 \\ v_r & \Theta_r \end{pmatrix}, \quad v_r = -\Theta_r u_r$$

with

$$\Theta_r^T = \begin{pmatrix} t_r & n_r & b_r \end{pmatrix}$$

$$t_r = \frac{(u_r)_x}{\|(u_r)_x\|}; \quad n_r = \frac{1}{\sin \alpha_r} \left(\frac{\Delta u_r}{\|\Delta u_r\|} - \cos \alpha_r t_r \right); \quad b_r = t_r \times n_r.$$

From now on, and for convenience, we will drop the subindex to denote position unless the situation is confusing, indicating a change in position by the application of the shift operator ($\mathcal{T}^k u = u_{n+k}$). We will calculate the *left* Maurer–Cartan

matrices, the more geometrically significant one (those in the continuous case are right ones). The left Maurer–Cartan matrices are given by

$$\widehat{K} = \rho \mathcal{T} \rho^{-1} = \begin{pmatrix} 1 & 0 \\ -\Theta u & \Theta \end{pmatrix} \begin{pmatrix} 1 & 0 \\ \mathcal{T} u & \mathcal{T} \Theta^T \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ \|\Delta u\| w & K \end{pmatrix}$$

where $K = \Theta \mathcal{T} \Theta^{-1}$; and by

$$\widehat{N} = \rho(\rho^{-1})_t = \begin{pmatrix} 1 & 0 \\ -\Theta u & \Theta \end{pmatrix} \begin{pmatrix} 0 & 0 \\ u_x & (\Theta^T)_x \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ \|u_x\| e_1 & N \end{pmatrix}$$

where $N = \Theta(\Theta^{-1})_x$. The local syzygies are given by

$$(39) \quad (\widehat{K})_x = \widehat{K} \mathcal{T} \widehat{N} - \widehat{N} \widehat{K}.$$

As we did in the continuous case, and to ensure convergence, we will restrict to the submanifold of the multispace defined locally by

$$\|u_x\| = 1 \quad \frac{\|\Delta u_n\|}{|\Delta y_n|} = 1.$$

For simplicity we will restrict further to those lattices where $|\Delta y_r| = \epsilon$ for all r (and hence $\|\Delta u_r\| = \epsilon$).

Equation (39) breaks into two equations, namely

$$(40) \quad w_t = K e_1 - N w - e_1$$

$$(41) \quad K_t = K \mathcal{T} N - N K.$$

Assume K factorizes as

$$(42) \quad K = Y \begin{pmatrix} \Omega & 0 \\ 0 & 1 \end{pmatrix}$$

with $\Omega \in \text{SO}(2)$, for some $Y = \exp \begin{pmatrix} 0 & y \\ -y^T & 0 \end{pmatrix}$ (this is always possible when $\mathcal{T} \Theta$ is closed enough to Θ so that K is closed enough to the identity). Assume further that

$$\Omega = \begin{pmatrix} \cos k & -\sin k \\ \sin k & \cos k \end{pmatrix} = (v \quad \widehat{v})$$

v and \widehat{v} denoting the two columns of Ω . We will denote with a hat the transformation $\widehat{v} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} v = Jv$, and so $\widehat{w} = Jw$. Finally, denote N by

$$(43) \quad N = \begin{pmatrix} \Lambda & z \\ -z^T & 0 \end{pmatrix}, \quad \Lambda = \begin{pmatrix} 0 & \nu \\ -\nu & 0 \end{pmatrix}, \quad z = \begin{pmatrix} z_1 \\ z_2 \end{pmatrix}.$$

With this notation (40) can be rewritten and simplified to equations

$$(44) \quad -s y \cdot v = z \cdot w$$

$$(45) \quad w \cdot e_1 = w \cdot v + c y \cdot v \quad y \cdot w$$

$$(46) \quad \alpha_x = -\nu + \frac{1}{\sin \alpha} (1 - \cos k - c y \cdot v \quad y_2)$$

where $s = \frac{1}{\|y\|} \sin \|y\|$, $c = \frac{1}{\|y\|^2} (\cos \|y\| - 1)$, and as usual $y^T = (y_1, y_2)$.

The remaining three equations that will determine the lattice/surface are given by the three entries in the $\mathfrak{so}(3)$ component (41). We will only reproduce the portion

corresponding to Sine-Gordon, that is the $(2, 1)$ entry of (41). After some long, but straightforward algebraic manipulations, the equation becomes

$$(47) \quad k_x + c \det(y, y_x) = \mathcal{T}\nu - \cos \|y\| \nu + s \det(y, z).$$

Looking at (37) and (36), and comparing it to (42) and (43), we see that

$$s \det(y, z)$$

discretizes the determinant of the first two entries of the last column in both (37) and (36). That is, $s \det(y, z)$ discretizes

$$\det \begin{pmatrix} f & e \\ \frac{1}{\sin \alpha} (g - \cos \alpha f) & \frac{1}{\sin \alpha} (f - \cos \alpha e) \end{pmatrix} = -\sin \alpha \mathbf{K}$$

where \mathbf{K} is the Gauss curvature of the surface. Therefore, we can define

$$\mathbf{K} = -\frac{s}{\sin \alpha} \det(y, z) = -\frac{\sin \|y\|}{\|y\| \sin \alpha} \det(y, z)$$

to be the discrete Gauss curvature for the lattice/surface. Then, (47) becomes

$$k_x + c \det(y, y_x) = \mathcal{T}\nu - \cos \|y\| - \sin \alpha \mathbf{K}.$$

This will be a discretization of Sine-Gordon, together with the other equations in the system.

APPENDIX A. EQUICONTINUOUS FAMILIES OF DISCRETE FRAMES

In this Appendix, we use the Arzela-Ascoli Theorem to give a general convergence result for an equicontinuous family of moving frames. This provides a rigorous foundation to a variety of examples involving the discretisation of a smooth frame.

Let M be a manifold, and let G be a Lie group with local metric d . The set G^M consists of all continuous maps from M to G , and we give it the compact-open topology, defined as that generated by finite intersections of the so-called subbasic sets,

$$(A, V) = \{f \in G^M \mid f(A) \subset V\}$$

where $A \subset M$ is open and $V \subset G$ is compact. A sequence of maps converging in this topology is uniformly convergent on compact subsets.

Definition A.1. A family $\mathcal{F} \subset G^M$ is said to be *equicontinuous* at $y_0 \in M$ if for all $\epsilon > 0$ there exists a neighbourhood $\mathcal{U}(y_0) \subset M$ such that for all $\rho \in \mathcal{F}$,

$$\rho(\mathcal{U}(y_0)) \subset B(\rho(y_0), \epsilon) = \{g \in G \mid d(g, \rho(y_0)) < \epsilon\}.$$

Theorem A.2. *Suppose that a family of left (resp. right) moving frames $\mathcal{F} \subset G^M$ satisfies the following:*

- (1) \mathcal{F} is equicontinuous on M , and
- (2) the set

$$\{\rho(y) \mid \rho \in \mathcal{F}\}$$

has compact closure for each y .

Then $\overline{\mathcal{F}}$ is a compact and equicontinuous family of moving frames.

Proof. We give the proof for \mathcal{F} a family of left frames, the proof for right frames is analogous. The conditions of the Theorem are precisely those of the Arzela-Ascoli Theorem, ([15], XII, Theorem 6.4), which yields that the family $\overline{\mathcal{F}}$ is compact and equicontinuous. We need only show that its elements are also equivariant with respect to the group action on M . Fix $y \in M$ and $f \in \overline{\mathcal{F}}$, and let $\epsilon > 0$ be given. By the definition of the closure of the set \mathcal{F} and the continuity of the group action, there is a $\rho \in \mathcal{F}$ and a neighbourhood U of the identity $e \in G$ such that for $g \in U$ we have both $d(f(g \cdot y), \rho(g \cdot y)) < \frac{1}{2}\epsilon$ and $d(g \cdot f(y), g \cdot \rho(y)) < \frac{1}{2}\epsilon$. Then

$$\begin{aligned} d(g \cdot f(y), f(g \cdot y)) &< d(f(g \cdot y), \rho(g \cdot y)) + d(\rho(g \cdot y), g \cdot f(y)) \\ &= d(f(g \cdot y), \rho(g \cdot y)) + d(g \cdot \rho(y), g \cdot f(y)) \\ &< \epsilon, \end{aligned}$$

so that f is equivariant, as required. \square

Example A.3. Consider the scaling and translation action of \mathbb{R}^2 given on a equivariant family of Lipschitz continuous curves $(x, y(x))$ in the plane by $g \cdot (x, y(x)) = (x, \exp(\lambda)y + k)$. A smooth frame is given by $g \cdot y = 0$, $g \cdot y_x = 1$, or $(\exp(\lambda), k)|_\rho = (1/y_x, -y/y_x)$. Suppose now we wish to discretise this frame in a way that is compatible with the smooth frame and with forward difference, that is $y_{n+1} = y_n + \alpha y_x$. Then the discrete frame ρ_n would be obtained by the normalisation equations, $g \cdot y_n = 0$, $g \cdot y_{n+1} = \alpha$, so that

$$\rho_n = \left(\frac{\alpha}{y_{n+1} - y_n}, \frac{-\alpha y_n}{y_{n+1} - y_n} \right).$$

This family of frames is straightforwardly seen to be equicontinuous, to have the smooth frame as its continuum limit, and to have the smooth Maurer Cartan invariants as the limit of the discrete ones, provided the parameter α scales as the mesh size $x_{n+1} - x_n$.

REFERENCES

- [1] George Bluman and Sukeyuki Kumei, *Symmetries and Differential Equations*, volume 154 of *Applied Mathematical Sciences*. Springer Verlag, New York, 1989
- [2] A.I. Bobenko and Y.B. Suris. Discrete differential geometry. Consistency as integrability, *Graduate Studies in Mathematics*, Vol. **98**. AMS, 2008.
- [3] A. Bobenko, U. Pinkall, *Discrete surfaces with constant negative Gaussian curvature and the Hirota equation*, J. Diff. Geometry **43** (1996), 527–611.
- [4] C. de Boor, *Polynomial interpolation in several variables*, in: Studies in Computer Science: In Honour of Samuel D. Conte. J.R. Rice and R. DeMillo (eds.), Plenum Press, 1994 and Springer Science+Business Media New York, 1994, pp87–110.
- [5] C. de Boor and A. Ron, *On Multivariate Polynomial Interpolation*, Constr. Approx. **6** (1990), 287–302.
- [6] C. de Boor and A. Ron, *The least solution for the polynomial interpolation problem*, Mathematische Zeitschrift **210** (1992), 347–378.
- [7] C. de Boor and A. Ron, *Computational aspects of polynomial interpolation in several variables*, Mathematics of Computation **58** (1992), 705–727.
- [8] M. Boutin, *On orbit dimensions under a simultaneous Lie group action on n copies of a manifold*, J. Lie Theory **12** (2002), 191–203.
- [9] A. Calini, T. Ivey and G. Marí Beffa, *An integrable flow for starlike curves in centroaffine space*, Symmetry, Integrability and Geometry: Methods and Applications **9** (2013), 022.
- [10] É. Cartan, *Oeuvres complètes*, Gauthier-Villiers, 1952–55.
- [11] S.S. Chern and K. Tenenblat, *Pseudo-spherical surfaces and evolution equations*, Stud. Appl. Math. **74** (1986), 55–83.

- [12] M. Chhay and A. Hamdouni, *A new construction for invariant numerical schemes using moving frames*, C. R. Acad. Sci. Meca. **338** (2010), 97–101.
- [13] É. Cotton, "Genralisation de la theorie du tridre mobile", Bull. Soc. Math. France **33** (1905), 1–23.
- [14] G. Darboux, *Lecons sur la theorie générale des surfaces et des applicaitons géométrique du calcul infinitesimal*, Gauthier-Villars, 1887.
- [15] J. Dugunji, *Topology*, Allyn and Bacon Inc., Boston, 1966.
- [16] M. Fels and P.J. Olver, *Moving Coframes I*, Acta Appl. Math. **51** (1998), 161–213.
- [17] M. Fels, P.J. Olver, *Moving coframes. II. Regularization and theoretical foundations* Acta Appl. Math. (1999), pp 127–208.
- [18] M. Gasca and J.I. Maeztu, *On Lagrange and Hermite interpolation in \mathbb{R}^k* , Numerische Mathematik **39** (1982), 1–14.
- [19] M. L. Green, *The moving frame, differential invariants and rigidity theorems for curves in homogeneous spaces*, Duke Math. J. **45** (1978), 735–779.
- [20] P. Griffiths, *On Cartan's methods of Lie groups and moving frames as applied to uniqueness and existence questions in differential geometry*, Duke Math. J. **41** (1974), 775–814.
- [21] T.M.N. Goncalves and E.L. Mansfield, *On Moving frames and Noether's Conservation Laws*, Studies in Applied Mathematics **128** (2011), 1–29
- [22] T.M.N. Goncalves and E.L. Mansfield, *Moving Frames and Conservation Laws for Euclidean Invariant Lagrangians*, Studies in Applied Mathematics **130** (2013), 134–166
- [23] M.S. Hickman, Willy A. Hereman, *Computation of densities and fluxes of nonlinear differential/difference equations*, Proceedings of the Royal Society A **457** (2003), 2705–2729.
- [24] R. Hasimoto, *A soliton on a vortex filament*, J. Fluid Mech. **51** (1972), 477–485.
- [25] K. Hikami, R. Inoue, *Classical lattice W algebras and integrable systems*, J. Phys. A: Math. Gen. **30** (1997), 6911–6924.
- [26] M.W. Hirsch, *Differential Topology*, Springer Verlag, New York, 1976.
- [27] E. Hubert, *Differential Algebra for Derivations with Nontrivial Commutation Rules*, J. of Pure and Applied Algebra **200** (1-2) (2005), 163–190.
- [28] E. Hubert, *Differential invariants of a Lie group action: syzygies on a generating set*, J. Symbolic Computation **44**(4) (2009a), 382–416.
- [29] E. Hubert, *Generation properties of Maurer–Cartan invariants*, (2009b) Preprint [hal:inria-00194528/en]
- [30] E. Hubert and I.A. Kogan, *Smooth and Algebraic Invariants of a Group Action. Local and Global Constructions*, Foundations of Computational Mathematics **7** (4) (2007a), 345–383.
- [31] E. Hubert and I. A. Kogan, *Rational Invariants of a Group Action. Construction and Rewriting*, Journal of Symbolic Computation **42** (2007b), 203–217.
- [32] P.E. Hydon and E.L. Mansfield, *A variational complex for difference equations*, Foundations of Computational Mathematics **4** (2004), 187–217.
- [33] P. Kim and P.J. Olver, *Geometric integration via multi-space*, Regular and Chaotic Dynamics **9**(3) (2004), 213–226.
- [34] P. Kim, *Invariantization of Numerical Schemes Using Moving Frames*, BIT Numerical Mathematics **47**(3) (2007), 525.
- [35] P. Kim, *Invariantization of the Crank-Nicolson Method for Burgers' Equation*, Physica D: Nonlinear Phenomena **237**(2) (2008), 243.
- [36] I.A. Kogan and P.J. Olver, *Invariant Euler-Lagrange equations and the invariant variational bicomplex*, Acta Appl. Math. **76** (2003), 137–193.
- [37] E.L. Mansfield, *A practical guide to the invariant calculus*, Cambridge Monographs on Applied and Computational Mathematics Volume 26, Cambridge University Press, 2010.
- [38] E.L. Mansfield and P.E. Hydon, *Difference Forms*, Foundations of Computational Mathematics **8** (2008), 427–467.
- [39] E.L. Mansfield and P. van der Kamp, *Evolution of curvature invariants and lifting integrability*, J. Geometry and Physics **56** (2006), 1294–1325.
- [40] E.L. Mansfield and P.E. Hydon, *Towards approximations of difference equations that preserve integrals*, Proc. 2001 International Symposium on Symbolic and Algebraic Manipulation (IS-SAC 2001) ed., B. Mourrain, ACM, New York, (2001), 217–222.
- [41] E.L. Mansfield, G. Marí Beffa, and J.P. Wang, *Discrete moving frames and integrable systems*, Foundations of Computational Mathematics **13**, Issue 4 (2013), 545–582.

- [42] G. Marí Beffa, *Geometric Hamiltonian structures on flat semisimple homogeneous manifolds*, the Asian Journal of Mathematics **12**, 1 (2008) 1–33.
- [43] G. Marí Beffa, *Poisson geometry of differential invariants of curves in some nonsemisimple homogeneous spaces*, Proc. Amer. Math. Soc. **134** (2006), 779–791.
- [44] G. Marí Beffa, *On bi-Hamiltonian flows and their realizations as curves in real semisimple homogeneous manifolds*, Pacific Journal of Mathematics **247** (2010), 163–188.
- [45] G. Marí Beffa, *Hamiltonian evolutions of twisted polygons in parabolic manifolds: the Lagrangian Grassmannian*, Pacific Journal of Mathematics **270** (2014), 287–317.
- [46] G. Marí Beffa, and J.P. Wang, *Hamiltonian evolutions of twisted gons in RP^n* , Nonlinearity **26** (2013), 2515–2551.
- [47] J. Munkres, *Topology*, Prentice-Hall, New Jersey, Second edition, 2000.
- [48] Emmy Noether, *Invariant variation problems*, Transport Theory Statist. Phys., 1(3):186–207, 1971. Translated from the German (Nachr. Akad. Wiss. Göttingen Math.-Phys. Kl. II 1918, 235–257).
- [49] P.J. Olver, *Moving frames – in geometry, algebra, computer vision, and numerical analysis*, Foundations of Computational Mathematics, R. DeVore, A. Iserles and E. Suli, eds., London Math. Soc. Lecture Note Series, vol. 284, Cambridge University Press, Cambridge, 2, 2001, 67–297.
- [50] P.J. Olver, *Geometric Foundations of Numerical Algorithms and Symmetry*, Appl. Alg. Engin. Comp. Commun. **11** (2001), 417–436.
- [51] P.J. Olver, *Joint invariant signatures*, Foundations of Computational Mathematics **1** (2001), 3–67.
- [52] P.J. Olver, *Applications of Lie groups to differential equations*, Graduate Texts in Mathematics 107, Springer Verlag, New York, Second edition, 1993.
- [53] L. Peng, E.L. Mansfield and P.E. Hydon, *Discrete moving frames and finite difference variational problems*, in preparation.
- [54] U. Pinkall, *Hamiltonian flows on the space of star-shaped curves*, Result. Math. **27** (1995), 328–332.
- [55] E.G. Reyes, *Pseudo-spherical surfaces and integrability of evolution equations*, J. Diff. Eq. **147** (1998) 195–230.
- [56] C.L. Terng, K. Uhlenbeck, *Schrödinger flows on Grassmannians, Integrable systems, geometry and topology*, AMS/IP Stud. Adv. Math, AMS, Providence (2006), 235–256.
- [57] L.W. Tu, *Une courte démonstration de la formule de Campbell-Hausdorff*, Journal of Lie Theory **14** (2004), 501– 508.