

CONTINUOUS CRAIG INTERPOLATION

H. JEROME KEISLER

ABSTRACT. We prove analogues of the Craig interpolation theorem for the continuous model theory of metric structures.

1. INTRODUCTION

In this paper we will prove two analogues of the Craig Interpolation Theorem for the continuous model theory of metric structures as developed in [BBHU] (2008). The model theory of metric structures is currently an active area of research with many applications to analysis, so it is important to clarify what happens to Craig interpolation in that setting.

In classical first order model theory, the Craig Interpolation Theorem ([Cr] 1957) says that:

For every sentence φ in a vocabulary V and ψ in a vocabulary W such that $\varphi \models \psi$, there is a sentence θ in the common vocabulary $V \cap W$ such that $\varphi \models \theta$ and $\theta \models \psi$.

The sentence θ is called a (Craig) *interpolant* of φ and ψ . Craig's proof used proof-theoretic methods. A closely related result, the Robinson Consistency Theorem ([Ro1] 1956), says that:

For every theory T_V in a vocabulary V , T_W in a vocabulary W , and complete theory T in the common vocabulary $V \cap W$, if each of $T \cup T_V$ and $T \cup T_W$ is consistent then $T \cup T_V \cup T_W$ is consistent.

Around 1959, several people noticed that the Craig Interpolation Theorem can be proved fairly easily from the Robinson Consistency Theorem, and vice versa (see Feferman [Fe] (2008) and Robinson [Ro2] (1963), pp. 114–117).

This paper was partly motivated by ongoing work with my son Jeffrey M. Keisler. In [KK1] (2012) and [KK2] (2014) we applied the first order Craig Interpolation Theorem to questions arising from the field of decision analysis. The present paper will enable the extension of that work from a discrete to a continuous setting.

In the introduction to the paper [BYP] (2010), Ben Yaacov and Petersen wrote that “continuous first-order logic satisfies a suitably

Date: January 15, 2025.

phrased form of Craig’s interpolation theorem.” (We will return to that point later). The paper [BYP] did not prove or even state a form of the Craig Interpolation Theorem for continuous logic. However, the paper [BYP] did develop a notion of formal proof for continuous logic that could perhaps be used to prove interpolation theorems. A continuous analogue of Beth’s Theorem was stated and proved in [BBHU] (Theorem 9.32). See also the monograph [Fa] (2021)).

We assume the reader is familiar with [BBHU]. In order to state our main results we give a few reminders to fix notation. Instead of the equality predicate symbol, the logic in [BBHU] has a distinguished binary predicate symbol d , called the *distance predicate*. We fix a set L of finitary predicate and function symbols such that $d \in L$, and a *metric signature* \mathbb{L} that specifies a modulus of uniform continuity with respect to d for each predicate or function symbol in L . A *vocabulary* V is a set $V \subseteq L$ such that $d \in V$. Atomic formulas with vocabulary V are the same as in first order logic. $[0, 1]$ -valued structures are like first order structures except that the atomic formulas take values in $[0, 1]$ instead of $\{\top, \perp\}$.

Continuous V -formulas and V -sentences are built from atomic formulas with vocabulary V using sup, inf as quantifiers, and n -ary continuous functions $C: [0, 1]^n \rightarrow [0, 1]$ (where $n \in \mathbb{N}$) as connectives. Each constant $r \in [0, 1]$ also counts as a V -sentence (and a 0-ary connective). In a $[0, 1]$ -valued structure \mathcal{M} with vocabulary V , each V -sentence φ has a truth value $\varphi^{\mathcal{M}} \in [0, 1]$, and φ is called true in \mathcal{M} if $\varphi^{\mathcal{M}} = 0$.

A *metric structure* is a $[0, 1]$ -valued structure \mathcal{M} with a vocabulary $V \subseteq L$ such that the interpretation of d in \mathcal{M} is a complete metric, and \mathcal{M} respects the bounds of uniform continuity in \mathbb{L} .

Convention 1.1. V and W will always denote vocabularies, φ will always be a V -sentence, and ψ will always be a W -sentence.

We will prove the following two analogues of Craig Interpolation for metric structures.

Definition 1.2. Let $\varepsilon \in (0, 1]$. A *weak ε -interpolant* of φ and ψ is a $V \cap W$ -sentence θ such that $\varphi = 0 \models \theta = 0$ and $\theta = 0 \models \psi \leq \varepsilon$.

Intuitively, θ is a sentence in the common language such that θ is true whenever φ is true, and ψ is almost true whenever θ is true.

Theorem 1.3. (*Weak Interpolant*) Suppose $\varphi = 0 \models \psi = 0$. Then for each $\varepsilon \in (0, 1]$, φ and ψ have a weak ε -interpolant.

Definition 1.4. Let $\varepsilon \in (0, 1]$. A *strong ε -interpolant* of φ and ψ is a $V \cap W$ -sentence θ such that $\models \varphi \geq \theta$ and $\models \theta \geq \psi - \varepsilon$.

Theorem 1.5. (*Strong Interpolant*) *Suppose that $\models \varphi \geq \psi$. Then for each $\varepsilon \in (0, 1]$, φ and ψ have a strong ε -interpolant.*

Theorems 1.3 and 1.5 can be compared with two interpolation theorems for linear continuous logic in [Ba] (2014). Linear continuous logic is like the continuous logic of [BBHU] except that the only connectives are linear functions, and the space of truth values is \mathbb{R} rather than $[0, 1]$ (see [BM] (2023)). The statements of Theorems 1.3 and 1.5 are similar to those of Propositions 6.7 and 6.8 in [Ba]. [Ba] gives direct proofs of both of Proposition 6.7 and 6.8 that resemble the model theoretic proof of the first order Craig interpolation theorem.

In the present setting of metric structures, the proof of the Weak Interpolant Theorem 1.3 again resembles the model theoretic proof of the first order Craig interpolation theorem. But the proof of the Strong Interpolant Theorem 1.5 is more difficult and uses the Weak Interpolant Theorem.

Before going on, we note that in the other direction, the Weak Interpolant Theorem is an easy consequence of the Strong Interpolant Theorem, because $\models \varphi \geq \psi$ trivially implies $\varphi = 0 \models \psi = 0$. Thus every strong ε -interpolant of φ and ψ is a weak ε -interpolant of φ and ψ .

The following corollary of the Weak Interpolant Theorem shows that except for extreme cases, the notion of a strong interpolant is strictly stronger than the notion of a weak interpolant.

Corollary 1.6. *Suppose $\models \varphi \geq \psi$, $\varepsilon > 0$, and there is a metric structure \mathcal{M} with vocabulary $V \cup W$ such that either $\varepsilon < \psi^{\mathcal{M}}$ or $\varphi^{\mathcal{M}} < 1$. Then there is a weak ε -interpolant of φ and ψ that is not a strong ε -interpolant of φ and ψ .*

Proof. By the Weak Interpolant Theorem, there is a weak ε -interpolant θ (of φ and ψ). If θ is already not a strong ε -interpolant, we are done. Assume instead that θ is a strong ε -interpolant. Let C be a strictly increasing continuous function from $[0, 1]$ into $[0, 1]$ such that $C(0) = 0$. Since θ is a weak ε -interpolant, the sentence $C(\theta)$ is also a weak ε -interpolant.

Case 1: $\varepsilon < \psi^{\mathcal{M}}$. Since θ is a strong ε -interpolant, $\theta^{\mathcal{M}} \geq \psi^{\mathcal{M}} - \varepsilon > 0$. Then we may take C such that $0 < C(\theta^{\mathcal{M}}) < \psi^{\mathcal{M}} - \varepsilon$, so the sentence $C(\theta)$ is not a strong ε -interpolant.

Case 2: $\varphi^{\mathcal{M}} < 1$. Since θ is a strong ε -interpolant, $1 > \varphi^{\mathcal{M}} \geq \theta^{\mathcal{M}}$. Then we may take C such that $1 > C(\theta^{\mathcal{M}}) > \varphi^{\mathcal{M}}$, so again the sentence $C(\theta)$ is not a strong ε -interpolant. \square

It is instructive to compare weak and strong interpolants in the case that $\models \varphi \geq \psi$. A strong ε -interpolant of φ and ψ is a $V \cap W$ -sentence θ such that for each $r \in [0, 1]$ and each \mathcal{M} we have

$$\varphi^{\mathcal{M}} \leq r \Rightarrow \theta^{\mathcal{M}} \leq r, \quad \theta^{\mathcal{M}} \leq r \Rightarrow (\psi^{\mathcal{M}} - \varepsilon) \leq r.$$

Thus $\theta^{\mathcal{M}}$ is always between $\varphi^{\mathcal{M}}$ and $\psi^{\mathcal{M}} - \varepsilon$, so “almost” between $\varphi^{\mathcal{M}}$ and $\psi^{\mathcal{M}}$. For each r , the Weak Interpolant Theorem for the sentences¹ $\varphi \div r$ and $\psi \div r$ gives us a $V \cap W$ -sentence θ_r (that may depend on r) such that for each \mathcal{M} ,

$$\varphi^{\mathcal{M}} \leq r \Rightarrow \theta_r^{\mathcal{M}} \leq r, \quad \theta_r^{\mathcal{M}} \leq r \Rightarrow (\psi^{\mathcal{M}} - \varepsilon) \leq r.$$

To prove the Strong Interpolant Theorem, we will have to construct a sentence θ_r that does not depend on r .

This is illustrated in Figure 1. The horizontal axis represents the class of all metric structures \mathcal{M} with vocabulary $V \cup W$, and the vertical axis represents the set of all $r \in [0, 1]$. The upper curve may be regarded as the “graph” of the function $\mathcal{M} \mapsto \varphi^{\mathcal{M}}$, and similarly for the other two curves.²

The region below the upper curve is the class of pairs (\mathcal{M}, r) where $r \leq \varphi^{\mathcal{M}}$, and the region below the lower curve is the class of pairs (\mathcal{M}, r) where $r \leq (\psi^{\mathcal{M}} - \varepsilon)$. The region below the middle curve is the class of pairs (\mathcal{M}, r) where $r \leq \theta^{\mathcal{M}}$ for strong interpolants and $r \leq \theta_r^{\mathcal{M}}$ for weak interpolants.

Our proof of the Strong Interpolant Theorem will go roughly as follows. Let $\varepsilon = 2^{-n}$ for some n . Let \mathbb{D}_n be the set of all multiples of ε in $[0, 1]$ (so \mathbb{D}_n has cardinality 2^n). For each $r \in \mathbb{D}_n$, use the Weak Interpolant Theorem to get the sentence θ_r defined in the above paragraphs. The hard part of the proof will be to find a continuous function C with 2^n variables such that the sentence $C(\langle \theta_r \rangle_{r \in \mathbb{D}_n})$ is a strong ε -interpolant of φ and ψ .

As mentioned above, the paper [BYP] stated that “continuous first-order logic satisfies a suitably phrased form of Craig’s interpolation theorem.” When asked what a suitable form of the Craig Interpolation Theorem for continuous logic would be, Ben Yaacov [BY] (2022) proposed the following, which in the present setting is easily seen to be equivalent to the Weak Interpolant Theorem above (and to Corollary 3.3 below).

¹ $\varphi \div r$ denotes the sentence $\max(\varphi - r, 0)$.

²The picture is intended to illustrate the idea, but there are actually too many models to fit on a real line and the “curves” can be very wild.




FIGURE 1. Interpolation when $\models \varphi \geq \psi$.

If $\{\varphi, \psi\}$ is inconsistent then there is a sentence θ in the common vocabulary such that $\{\varphi, \theta \div 1/2\}$ is inconsistent and $\{\psi, 1/2 \div \theta\}$ is inconsistent.

Ben Yaacov [BY] pointed out that the above statement implies that a uniform limit of sentences in the common vocabulary could serve as an interpolant. This is also equivalent to the Weak Interpolant Theorem (see Corollary 3.5 below).

In Section 2 we will prove an analogue of the Robinson Consistency Theorem for metric structures (see Theorem 2.7 below) using results in the literature about saturated and special metric structures. In Section 3 we will prove the Weak Interpolant Theorem from the continuous Robinson Consistency Theorem. In Section 4 we will prove the Strong Interpolant Theorem from the Weak Interpolant Theorem, using an argument that does not have a first order counterpart.

I thank Itai Ben Yaacov and Jeffrey M. Keisler for helpful discussions related to this paper.

2. ROBINSON CONSISTENCY THEOREM

We first recall some basic notation and facts about metric structures that will be needed in this and the following sections. We will then prove a continuous analogue of the Robinson Consistency Theorem.

By a *V-theory* we mean a set of *V*-sentences. We say that \mathcal{M} is a *metric model* of *T*, in symbols $\mathcal{M} \models T$, if \mathcal{M} is a metric structure, *T* is a *V-theory* where *V* is the vocabulary of \mathcal{M} , and every $\theta \in T$ is true (has truth value 0) in \mathcal{M} . For *V*-theories *T* and *U*, $T \models U$ means that every metric model of *T* is a metric model of *U*. We write $T \models \varphi \geq \theta$ if $\varphi^{\mathcal{M}} \geq \theta^{\mathcal{M}}$ for every metric model $\mathcal{M} \models T$ (similarly for \leq and $=$). Thus $T \models \{\varphi\}$ if and only if $T \models \varphi = 0$. Note that to the right of the \models symbol we can have either a set of sentences or an inequality or equation between a pair of sentences.

T is *consistent* if *T* has at least one metric model, and *T* is *inconsistent* if *T* has no metric models. If \mathcal{M} is a metric structure, or even just a $[0, 1]$ -valued structure, with vocabulary *V*, the *theory of* \mathcal{M} is the set $\text{Th}(\mathcal{M})$ of all *V*-sentences true in \mathcal{M} .

Let \mathcal{N} be a metric structure with vocabulary $V \cup W$. The *V-part* of \mathcal{N} is the metric structure $\mathcal{M} = \mathcal{N} \upharpoonright V$ with vocabulary *V* that agrees with \mathcal{N} on all symbols of *V*, and \mathcal{N} is called an *expansion* of \mathcal{M} to $V \cup W$. Note that if *T* is a *V-theory*, then *T* is also a $V \cup W$ -theory, and that $\mathcal{N} \models T$ if and only if $\mathcal{N} \upharpoonright V \models T$.

For *V*-sentences φ and θ , $\varphi \leq \theta$ (also written $\varphi \div \theta$) denotes the *V*-sentence $\max(\varphi - \theta, 0)$. The sentence $\varphi \leq \theta$ is useful because in every metric structure \mathcal{M} we have

$$(\varphi \leq \theta)^{\mathcal{M}} = 0 \text{ if and only if } \varphi^{\mathcal{M}} \leq \theta^{\mathcal{M}}.$$

We will use either \leq or \div , whichever seems more natural in each case. Also, $\varphi \geq \theta$ denotes the *V*-sentence $\theta \leq \varphi$, and $\varphi \dagger \theta$ denotes the *V*-sentence $\min(\varphi + \theta, 1)$.

We will repeatedly use the following fact and its corollary.

Fact 2.1. (*Compactness Theorem, by Theorem 5.8 of [BBHU]*) *If every finite subset of a V-theory T is consistent, then T is consistent.*

Corollary 2.2. (*Compactness Corollary*) *$T \cup \{\varphi\}$ is inconsistent if and only if there exists $r \in (0, 1]$ such that $T \models r \leq \varphi$.*

Proof. If there exists $r \in (0, 1]$ such that $T \models r \leq \varphi$, it trivially follows that $T \cup \{\varphi\}$ has no metric models and hence is inconsistent.

Suppose there is no $r \in (0, 1]$ such that $T \models r \leq \varphi$. Then for each $r \in (0, 1]$ there is a metric model \mathcal{M} of *T* such that $(r \leq \varphi)^{\mathcal{M}} > 0$, so $r > \varphi^{\mathcal{M}}$, $\varphi^{\mathcal{M}} \leq r$, and $(\varphi \leq r)^{\mathcal{M}} = 0$. Then every finite subset of the theory

$U = T \cup \{\varphi \leq r : r \in (0, 1]\}$ is consistent. By the Compactness Theorem, U is consistent, and thus has a metric model \mathcal{N} . Then $(\varphi \leq r)^{\mathcal{N}} = 0$ for each $r \in (0, 1]$, so $\varphi^{\mathcal{N}} = 0$ and $\mathcal{N} \models T \cup \{\varphi\}$. Therefore $T \cup \{\varphi\}$ is not inconsistent. \square

Two metric structures \mathcal{M}, \mathcal{N} with vocabulary V are *isomorphic*, in symbols $\mathcal{M} \cong \mathcal{N}$, if there is a bijection from the universe of \mathcal{M} onto the universe of \mathcal{N} that preserves the truth value of all atomic formulas.

Fact 2.3. *If $\mathcal{M} \cong \mathcal{N}$ then $\varphi^{\mathcal{M}} = \varphi^{\mathcal{N}}$ for every V -sentence φ .*

A cardinal κ is *special* if $2^\lambda \leq \kappa$ for all $\lambda \leq \kappa$ (for example, \beth_ω is special). A metric structure \mathcal{M} is κ -*special* if κ is an uncountable special cardinal, $|M| \leq \kappa$, and \mathcal{M} is the union of an elementary chain of metric structures $\langle \mathcal{M}_\lambda : \aleph_0 \leq \lambda < \kappa \rangle$ such that each \mathcal{M}_λ is λ^+ -saturated.

Fact 2.4. *(Facts 2.4.6 and 2.4.8 in [Ke].) Suppose T is a V -theory, κ is special, and $\kappa \geq |V| + \aleph_0$. If T is consistent then T has a κ -special model with vocabulary V . If T is complete then up to isomorphism T has a unique κ -special metric model with vocabulary V .*

Fact 2.5. *(Remark 2.4.4 in [Ke]) The V -part of a κ -special metric model with vocabulary $V \cup W$ is a κ -special metric model with vocabulary V .*

Convention 2.6. As before, V and W will always denote vocabularies, φ will always be a V -sentence, and ψ will always be a W -sentence. We also fix a V -theory T_V and a W -theory T_W .

The following result is a continuous analogue of the Robinson Consistency Theorem.

Theorem 2.7. *Let $T = \text{Th}(\mathcal{P})$ for some metric structure \mathcal{P} with vocabulary $V \cap W$. If $T \cup T_V$ is consistent and $T \cup T_W$ is consistent, then $T \cup T_V \cup T_W$ is consistent.*

Proof. Let κ be a special cardinal with $\kappa > |V \cup W| + \aleph_0$. By Fact 2.4, there is a κ -special metric model \mathcal{M} of $T \cup T_V$ with vocabulary V , and a κ -special metric model \mathcal{N} of $T \cup T_W$ with vocabulary W . By Fact 2.5, the $V \cap W$ -parts of \mathcal{M} and of \mathcal{N} are both metric models of T , and are both κ -special with vocabulary $V \cap W$. Since T is complete, by Fact 2.4 the $V \cap W$ -parts of \mathcal{M} and \mathcal{N} are isomorphic. Therefore we may take \mathcal{M} and \mathcal{N} to have the same $V \cap W$ -parts. Then \mathcal{M} and \mathcal{N} have a common expansion to a metric model of $T \cup T_V \cup T_W$. \square

3. WEAK INTERPOLANTS

Convention 2.6, where we fix the theories T_V and T_W , is still in force. Instead of proving the interpolation theorems in the simple form stated in the Introduction, it will be easier and hence better to prove them in the more general setting where we restrict attention to the class of metric models of $T_V \cup T_W$ instead of the class of all metric structures.

The Weak Interpolant Theorem 1.3 in the Introduction is a special case of Theorem 3.1 below (that special case arises when both T_V and T_W are empty sets of sentences).

Theorem 3.1. *Suppose that*

$$T_V \cup T_W \cup \{\varphi\} \models \psi = 0.$$

Then for each $\varepsilon \in (0, 1]$, there is a $V \cap W$ -sentence θ such that

$$T_V \cup \{\varphi\} \models \theta = 0, \quad T_W \cup \{\theta\} \models \psi \leq \varepsilon.$$

Proof. Let U be the set of all $V \cap W$ -sentences ρ such that $T_V \cup \{\varphi\} \models \rho = 0$. Then every metric model of $T_V \cup \{\varphi\}$ is a metric model of U . We first prove the following Claim.

Claim. $T_W \cup U \models \psi = 0$.

Proof of Claim. Suppose \mathcal{M} is a metric model of $T_W \cup U$. To show that $\psi^{\mathcal{M}} = 0$, we assume that $0 < \psi^{\mathcal{M}}$ and get a contradiction. Since $0 < \psi^{\mathcal{M}}$, $r \leq \psi^{\mathcal{M}}$ for some $r \in (0, 1]$. Let \mathcal{P} be the $V \cap W$ -part of \mathcal{M} and $T = \text{Th}(\mathcal{P})$. Then \mathcal{M} is a metric model of $T \cup T_W \cup \{r \leq \psi\}$, so

$$(1) \quad T \cup T_W \cup \{r \leq \psi\} \text{ is consistent.}$$

We prove

$$(2) \quad T \cup T_V \cup \{\varphi\} \text{ is consistent.}$$

Assume that (2) fails. By the Compactness Theorem, there is a finite subset $T_0 \subseteq T$ such that $T_0 \cup T_V \cup \{\varphi\}$ is inconsistent. Let ρ be the $V \cap W$ -sentence $\max(T_0)$. Then $\{\rho\} \cup T_V \cup \{\varphi\}$ is inconsistent. By the Compactness Corollary, there is an $s \in (0, 1]$ such that

$$T_V \cup \{\varphi\} \models s \leq \rho.$$

Hence $s \leq \rho \in U$. We have $\mathcal{M} \models \rho = 0$ because $\mathcal{M} \models T$, and $\mathcal{M} \models s \leq \rho$ because $\mathcal{M} \models U$. This is a contradiction, so (2) holds after all.

It follows from (1), (2), and Theorem 2.7 that

$$T \cup T_V \cup T_W \cup \{\varphi, r \leq \psi\} \text{ is consistent.}$$

This contradicts the hypothesis that $T_V \cup T_W \cup \{\varphi\} \models \psi = 0$ in the statement of the theorem, so the assumption that $0 < \psi^{\mathcal{M}}$ is false. Thus $\psi^{\mathcal{M}} = 0$ and hence

$$T_W \cup U \models \psi = 0$$

as required. This completes the proof of the Claim. \square

Let $\varepsilon \in (0, 1]$. Then by the Claim,

$$T_W \cup U \cup \{\varepsilon \leq \psi\}$$

is inconsistent. By the Compactness Theorem, there is a finite subset U^ε of U such that

$$T_W \cup U^\varepsilon \cup \{\varepsilon \leq \psi\}$$

is inconsistent. Let $\theta := \max(U^\varepsilon)$. Then $\theta \in U$, so

$$T_V \cup T_W \cup \{\varphi\} \models \theta = 0, \quad T_V \cup T_W \cup \{\theta\} \models \psi \leq \varepsilon.$$

as required. \square

Corollary 3.2. *Suppose that*

$$T_V \cup T_W \cup \{\varphi\} \models \psi = 0.$$

Then for each $\varepsilon \in (0, 1]$, there is a $V \cap W$ -sentence ρ such that

$$T_V \cup \{\varphi\} \models \rho \leq \varepsilon, \quad T_W \cup \{\rho \leq \varepsilon\} \models \psi \leq \varepsilon.$$

Proof. If θ is as in Theorem 3.1 then $\rho := \theta \dot{+} \varepsilon$ has the required properties. \square

Corollary 3.3. *Suppose $T_V \cup T_W \cup \{\varphi, \psi\}$ is inconsistent. Then there is a $V \cap W$ -sentence θ such that*

$$T_V \cup \{\varphi\} \models \theta = 1, \quad T_W \cup \{\psi\} \models \theta = 0.$$

Proof. By the Compactness Corollary, there exists $r \in (0, 1]$ such that

$$T_V \cup T_W \cup \{\varphi\} \models r \leq \psi.$$

Let $\varepsilon \in (0, r]$. By Theorem 3.1 there is a $V \cap W$ -sentence ρ such that

$$T_V \cup \{\varphi\} \models \rho = 0, \quad T_W \cup \{\rho\} \models (r \dot{-} \psi) \leq \varepsilon/2.$$

One can easily check that for any $x \in [0, 1]$ we have $(r \dot{-} x) \leq \varepsilon/2$ if and only if $(r - \varepsilon/2) \leq x$. Therefore

$$0 < r - \varepsilon/2, \quad T_W \cup \{\rho\} \models r - \varepsilon/2 \leq \psi,$$

so $T_W \cup \{\rho\} \cup \{\psi\}$ is inconsistent. By the Compactness Corollary, there exists $s \in (0, 1/2]$ such that

$$T_W \cup \{\psi\} \models s \leq \rho.$$

Then $\theta := 1 \dot{-} (\rho/s)$ has the required properties. \square

Note that Theorem 3.1 is also an easy consequence of Corollary 3.3. To see that, suppose $T_V \cup T_W \cup \{\varphi\} \models \psi = 0$ and $\varepsilon \in (0, 1]$. Then $T_V \cup T_W \cup \{\varphi, \varepsilon \leq \psi\}$ is inconsistent, so Corollary 3.3 gives a $V \cap W$ -sentence θ such that $T_V \cup \{\varphi\} \models \theta = 0$ and $T_W \cup \{\varepsilon \leq \psi\} \models \theta = 1$, and hence $T_W \cup \{\theta\} \models \psi \leq \varepsilon$.

While the Weak Interpolant Theorem shows that a single $V \cap W$ sentence serves as a weak ε -interpolant, the next corollary shows that a countable set of $V \cap W$ -sentences can serve as a weak interpolant (without the ε).

Corollary 3.4. *Suppose $T_V \cup T_W \cup \{\varphi\} \models \psi = 0$. Then for some countable set Θ of $V \cap W$ -sentences, $T_V \cup \{\varphi\} \models \Theta$ and $T_W \cup \Theta \models \psi = 0$.*

Proof. By Theorem 3.1, for each $n \in \mathbb{N}$ there is a $V \cap W$ -sentence θ_n such that $T_V \cup \{\varphi\} \models \rho_n = 0$ and $T_W \cup \{\rho_n\} \models \psi \leq 2^{-n}$. So the result holds with $\Theta = \{\rho_n \mid n \in \mathbb{N}\}$. \square

We say that a sequence of sentences $\langle \theta_n \rangle_{n \in \mathbb{N}}$ is *uniformly convergent* if for every $\varepsilon > 0$ there exists n such that for all $k > n$, $\models |\theta_k - \theta_n| \leq \varepsilon$. Note that if $\langle \theta_n \rangle_{n \in \mathbb{N}}$ is uniformly convergent then $\lim_{n \rightarrow \infty} \theta_n$ exists in every metric structure, and the ‘‘rate of convergence’’ is uniform across all metric structures. Also, if $\models |\theta_{n+1} - \theta_n| \leq 2^{-n}$ for every n then $\langle \theta_n \rangle_{n \in \mathbb{N}}$ is uniformly convergent.

The next corollary shows that a uniform limit of $V \cap W$ sentences can serve as a weak interpolant.

Corollary 3.5. *Suppose $T_V \cup T_W \cup \{\varphi\} \models \psi = 0$. Then there is a uniformly convergent sequence $\langle \theta_n \rangle_{n \in \mathbb{N}}$ of $V \cap W$ -sentences such that*

- (i) *In every model of T_V , if $\varphi = 0$ then $\lim_{n \rightarrow \infty} \theta_n = 0$.*
- (ii) *In every model of T_W , if $\lim_{n \rightarrow \infty} \theta_n = 0$ then $\psi = 0$.*

Proof. Let $\theta_n = \max_{m \leq n} \min(\rho_m, 2^{-m})$ where ρ_n is as in the proof of Corollary 3.4. For each n , θ_n is a $V \cap W$ -sentence. We let \mathcal{M} be a metric structure with vocabulary $V \cap W$. Clearly, $\theta_n^{\mathcal{M}} \leq \theta_{n+1}^{\mathcal{M}}$ for each n . Therefore $\lim_{n \rightarrow \infty} \theta_n^{\mathcal{M}} = 0$ if and only if $(\forall n) \theta_n^{\mathcal{M}} = 0$. We show that

$$(3) \quad \theta_{n+1}^{\mathcal{M}} \leq \theta_n^{\mathcal{M}} + 2^{-(n+1)}$$

for each n . (3) is trivially true if $\theta_{n+1}^{\mathcal{M}} = \theta_n^{\mathcal{M}}$. If $\theta_{n+1}^{\mathcal{M}} > \theta_n^{\mathcal{M}}$, then (3) still holds because

$$\theta_{n+1}^{\mathcal{M}} = \min(\rho_{n+1}^{\mathcal{M}}, 2^{-(n+1)}) \leq 2^{-(n+1)} \leq \theta_n^{\mathcal{M}} + 2^{-(n+1)}.$$

It follows that $\langle \theta_n \rangle_{n \in \mathbb{N}}$ is uniformly convergent, and by Corollary 3.4, (i) and (ii) hold. \square

4. STRONG INTERPOLANTS

Convention 2.6 is still in force. The Strong Interpolant Theorem 1.5 in the Introduction is a special case of Theorem 4.1 below (Theorem 4.1 reduces to Theorem 1.5 when both T_V and T_W are empty sets of sentences).

Theorem 4.1. *Suppose that $T_V \cup T_W \models \varphi \geq \psi$. Then for each $\varepsilon > 0$ there is a $V \cap W$ -sentence θ such that*

- (i) $T_V \models \varphi \geq \theta$,
- (ii) $T_W \models \theta \geq (\psi \dot{-} \varepsilon)$.

Proof. Fix $n \in \mathbb{N}$ and let $\varepsilon = 2^{-n}$. It suffices to find a $V \cap W$ -sentence θ such that

- (i*) $T_V \models \varphi \geq (\theta \dot{-} \varepsilon)$.
- (ii) $T_W \models \theta \geq (\psi \dot{-} \varepsilon)$.

Because if (i*) and (ii) hold with $\varepsilon/2$ in place of ε , then (i) and (ii) hold with $\theta \dot{-} \varepsilon/2$ in place of θ .

Our proof will have two parts. In the first part we will use Theorem 3.1 to find, for each $k < 2^n$, a $V \cap W$ -sentence ρ_k such that

$$(4) \quad T_V \cup \{\varphi \leq k\varepsilon\} \models \rho_k = 0,$$

and

$$(5) \quad T_W \cup \{\psi \geq (k+1)\varepsilon\} \models \rho_k = 1.$$

In the second part of the proof we will find a continuous function $f: [0, 1]^{2^n} \rightarrow [0, 1]$ such that $\theta := f(\langle \rho_k \rangle_{k < 2^n})$ satisfies (i*) and (ii).

The “graph” of $\rho_k^{\mathcal{M}}$ as a function of \mathcal{M} is illustrated by the bold line in Figure 2. As in Figure 1, we put the interval $[0, 1]$ on the vertical axis and the class of all metric models \mathcal{M} of $T_V \cup T_W$ on the horizontal axis.

First part of proof: Let $k < 2^n$. Since $T_V \cup T_W \models \varphi \geq \psi$, we have

$$T_V \cup T_W \cup \{\varphi \leq k\varepsilon\} \models \psi \leq k\varepsilon.$$

Then by Theorem 3.1 (with $\varphi \leq k\varepsilon$, $\psi \leq k\varepsilon$, $\varepsilon/2$ in place of $\varphi, \psi, \varepsilon$), there is a $V \cap W$ -sentence γ_k such that

$$(6) \quad T_V \cup \{\varphi \leq k\varepsilon\} \models \gamma_k = 0$$

and

$$T_W \cup \{\gamma_k\} \models \psi \leq k\varepsilon + \varepsilon/2.$$

Therefore the set of sentences

$$T_W \cup \{\gamma_k\} \cup \{\psi \geq (k+1)\varepsilon\}$$




FIGURE 2. Graphs of $\varphi^{\mathcal{M}}$, $\psi^{\mathcal{M}}$, and $\rho_k^{\mathcal{M}}$

is inconsistent. By Compactness Corollary, there exists $r \in (0, 1]$ such that

$$(7) \quad T_W \cup \{\psi \geq (k+1)\varepsilon\} \models \gamma_k \geq r.$$

Now $\rho_k := \min(\gamma_k/r, 1)$ is a $V \cap W$ -sentence. By the definition of ρ_k , (4) follows at once from (6), and (5) follows at once from (7).

Second part of proof: Let $f: [0, 1]^{2^n} \rightarrow [0, 1]$ be the continuous function

$$f(\vec{x}) := \max_{k < 2^n} \left[(k+1)\varepsilon \prod_{j \leq k} x_j \right].$$

Let

$$\theta := f(\langle \rho_k \rangle_{k < 2^n}).$$

Since f is continuous and each ρ_k is a $V \cap W$ -sentence, θ is a $V \cap W$ -sentence. We show that θ satisfies (i*) and (ii). It is clear that f has the following properties for each $k < 2^n$:

- (a) If $x_k = 0$ then $f(\vec{x}) \leq k\varepsilon$.

(b) If $x_j = 1$ for each $j \leq k$ then $f(\vec{x}) \geq (k+1)\varepsilon$.

To prove (i*), we work in an arbitrary metric model \mathcal{M} of T_V . In \mathcal{M} , if $\varphi \geq 1 - \varepsilon$ then it is trivial that $\varphi \geq (\theta \dot{-} \varepsilon)$. Suppose instead that $\varphi < 1 - \varepsilon$. Then there is a unique $k < 2^n - 1$ such that

$$k\varepsilon \leq \varphi < (k+1)\varepsilon.$$

Then $k+1 < 2^n$ and $\varphi \leq (k+1)\varepsilon$, so by (4) we have $\rho_{k+1} = 0$. Hence by (a), $\theta \leq (k+1)\varepsilon$. Therefore $(\theta \dot{-} \varepsilon) \leq k\varepsilon$ and $k\varepsilon \leq \varphi$. This proves (i*).

To prove (ii) we work in an arbitrary model of T_W . If $\psi \leq \varepsilon$ then it is trivial that $\theta \geq (\psi \dot{-} \varepsilon)$. Suppose instead that $\psi > \varepsilon$. Then there is a unique $0 < \ell < 2^n$ such that

$$\ell\varepsilon < \psi \leq (\ell+1)\varepsilon.$$

Hence for each $j \leq \ell - 1$ we have $(j+1)\varepsilon < \psi$, so $(j+1)\varepsilon \leq \psi$. By (5), $\rho_j = 1$ for each $j \leq \ell - 1$. Since $\ell - 1 \in \mathbb{N}$ we may apply (b) to get $\theta \geq \ell\varepsilon$. Since $\psi \leq (\ell+1)\varepsilon$, we also have $\ell\varepsilon \geq \psi \dot{-} \varepsilon$. Therefore $\theta \geq (\psi \dot{-} \varepsilon)$. This proves (ii). \square

Note that in the above proof, the function f is non-decreasing in each variable. Such functions are sometimes called aggregation functions.

Corollary 4.2. *Suppose that $T_V \cup T_W \models \varphi \geq \psi$. Then for each $\varepsilon > 0$ there is a $V \cap W$ -sentence ρ such that*

- (i) $T_V \models \varphi \geq (\rho \dot{-} \varepsilon)$,
- (ii) $T_W \models (\rho \dot{-} \varepsilon) \geq (\psi \dot{-} \varepsilon)$.

Proof. If θ is as in Theorem 4.1 then $\rho := \theta \dot{+} \varepsilon$ has the required properties. \square

The next corollary shows that a uniformly convergent sequence of $V \cap W$ -sentences can serve as a strong interpolant.

Corollary 4.3. *Suppose $T_V \cup T_W \models \varphi \geq \psi$. Then there is a uniformly convergent sequence $\langle \theta_n \rangle_{n \in \mathbb{N}}$ of $V \cap W$ -sentences such that*

- (i) *In every metric model of T_V , $\varphi \geq \lim_{n \rightarrow \infty} \theta_n$.*
- (ii) *In every metric model of T_W , $\lim_{n \rightarrow \infty} \theta_n \geq \psi$.*

Proof. By Theorem 4.1, for each $n \in \mathbb{N}$ there is a $V \cap W$ -sentence γ_n such that $\varphi \geq \gamma_n$ in any metric model of T_V , and $\gamma_n \geq (\psi \dot{-} 2^{-n})$ in any metric model of T_W . Now let $\theta_0 := 0$, and for each n let

$$\theta_{n+1} := \max(\theta_n, \min(\gamma_n, \theta_n \dot{+} 2^{-n})).$$

It is clear that each θ_n is a $V \cap W$ -sentence, and

$$\models \theta_n \leq \theta_{n+1}, \quad \models |\theta_{n+1} - \theta_n| \leq 2^{-n}.$$

Therefore $\langle \theta_n \rangle_{n \in \mathbb{N}}$ is uniformly convergent. We also have

$$\models \theta_{n+1} \leq \max(\theta_n, \gamma_n), \quad T_V \models \varphi \geq \gamma_n.$$

It follows by induction that $T_V \models \varphi \geq \theta_n$, so (i) holds. And

$$\models \theta_{n+1} \geq \min(\gamma_n, \theta_n + 2^{-n}), \quad T_W \models \gamma_n \geq (\psi \div 2^{-n}),$$

so it follows by induction that $T_W \models \theta_n \geq (\psi \div 2^{-n})$. Hence (ii) holds as well. \square

The arguments in this paper actually prove more general results that apply to arbitrary $[0, 1]$ -valued structures as developed in [Ke] as well as to metric structures.

Hereafter, V and W will denote vocabularies, φ will always be a V -sentence, and ψ will always be a W -sentence. We fix a V -theory T_V and a W -theory T_W . But no metric signature is given. $T \models_g U$ will mean that every general $[0, 1]$ -valued model of T is a model of U .

Here is the analogue of the Weak Interpolant Theorem for general structures.

Theorem 4.4. *Suppose that*

$$T_V \cup T_W \cup \{\varphi\} \models_g \psi = 0.$$

Then for each $\varepsilon \in (0, 1]$, there is a $V \cap W$ -sentence θ such that

$$T_V \cup \{\varphi\} \models_g \theta = 0, \quad T_W \cup \{\theta\} \models_g \psi \leq \varepsilon.$$

Here is the analogue of the Strong Interpolant Theorem for general structures.

Theorem 4.5. *Suppose that $T_V \cup T_W \models_g \varphi \geq \psi$. Then for each $\varepsilon > 0$ there is a $V \cap W$ -sentence θ such that*

- (i) $T_V \models_g \varphi \geq \theta$,
- (ii) $T_W \models_g \theta \geq (\psi \div \varepsilon)$.

To complete the picture, we recall one more result from [BBHU].

Fact 4.6. *(Theorem 3.7 in [BBHU]) If \mathbb{L} is a metric signature with vocabulary V and T_V contains sentences that specify the uniform continuity moduli for \mathbb{L} , then every general $[0, 1]$ -valued model of T_V is elementarily equivalent to a metric structure with signature \mathbb{L} .*

Thus when T_V and T_W contain sentences that specify the uniform continuity moduli for \mathbb{L} , Theorem 3.1 is a special case of Theorem 4.4, and Theorem 4.1 is a special case of Theorem 4.5.

REFERENCES

- [Ba] Seyed-Mohammad Bagheri. Linear model theory for Lipschitz structures. *Arch. Math. Logic* 53 (2014), 897-927.
- [BM] Seyed-Mohammad Bagheri and Mahya Malekghasemi. Consistency and interpolation in linear continuous logic. *Arch. Math. Logic* 62 (2023), 931-939.
- [BY] Itai Ben Yaacov, private communication, June 16, 2022.
- [BYP] Itai Ben Yaacov and Authur Paul Pedersen. A proof of completeness for continuous first-order logic. *J. Symbolic Logic* 75 (2010), 168–190.
- [BBHU] Itai Ben Yaacov, Alexander Berenstein, C. Ward Henson and Alexander Usvyatsov. Model Theory for Metric Structures. In *Model Theory with Applications to Algebra and Analysis*, vol. 2, London Math. Society Lecture Note Series, vol. 350 (2008), 315-427.
- [CK] C.C.Chang and H. Jerome Keisler. *Model Theory*. Dover 2012.
- [Cr] William Craig. Three uses of the Herbrand-Gentzen theorem in relating model theory and proof theory. *J. Symbolic Logic* 22 (1957), 269-285.
- [Fa] Ilijas Farah et.al. Model theory of C^* -algebras. arXiv:1602.08072v4 [math.LO] (2017), *Memoirs of the American Mathematical Society* No. 1324, Vol. 271 (2021).
- [Fe] Solomon Feferman. Harmonious Logic: Craig’s Interpolation Theorem and its Descendants. *Synthese* 164 no. 3 (2008), 341-357.
- [Ke] H. Jerome Keisler. Model theory for real-valued structures. In “Beyond First Order Model Theory, Vol. 2”, ed. by Jose Iovino, CRC Press 2023. Also arXiv:2005.11851, 2020.
- [KK1] H. Jerome Keisler and Jeffrey M. Keisler. Craig Interpolation for Networks of Sentences. *Annals of Pure and Applied Logic* 163 (2012), pp. 1322-1344.
- [KK2] H. Jerome Keisler and Jeffrey M. Keisler. Observing, Reporting, and Deciding in Networks of Sentences. *Annals of Pure and Applied Logic* 165 (2014), pp. 812-836.
- [Ro1] Abraham Robinson. A result on consistency and its application to the theory of definition. *Koninkl. Ned. Akad. Wetensch. Proc. Ser. A* 59 (= *Indag. Math* 18) (1956), 47–58.
- [Ro2] Abraham Robinson. *Introduction to Model Theory and to the Metamathematics of Algebra*. North-Holland 1963.

UNIVERSITY OF WISCONSIN-MADISON, DEPARTMENT OF MATHEMATICS, MADISON, WI 53706-1388

Email address: keisler@math.wisc.edu