

Enumerations of families closed under finite differences*

Noam Greenberg[†], Matthew Harrison-Trainor[‡],
Joseph S. Miller, and Dan Turetsky

March 31, 2021

Abstract

Slaman and Wehner independently built a family of sets with the property that every non-computable degree can compute an enumeration of the family, but there is no computable enumeration of the family. We call such a family a Slaman–Wehner family. Loosely motivated by the problem of whether there is an abelian group with the Slaman–Wehner degree spectrum, we consider families \mathcal{F} that are closed under finite differences: if $A \in \mathcal{F}$ and $B =^* A$, then $B \in \mathcal{F}$. The main question of this paper is whether there is a Slaman–Wehner family closed under finite differences. The Slaman–Wehner construction relies on the fact that all of the sets in the family are finite, and so no similar construction can work for a family closed under finite differences. Nonetheless, we are unable to answer this question, though we obtain a number of interesting partial results which can be interpreted as saying that the question is quite hard.

First of all, no Slaman–Wehner family closed under finite differences can contain a finite set, and the enumeration of the family from a non-computable degree cannot be uniform (whereas, in the Slaman–Wehner construction, it is uniform). On the other hand, we build the following examples of families closed under finite differences which show the impossibility of several natural attempts to show that no Slaman–Wehner family exists: (1) a family that can be enumerated by every non-low degree, but not by any low degree; (2) a family that can be enumerated by any set in a given uniform list of c.e. sets, but which cannot be enumerated computably; and (3) a family that can be enumerated by a given Δ_2^0 set, but which cannot be computably enumerated.

1 Introduction

This paper is concerned with the computational properties of families of subsets of ω . Such families have played an important role in computable structure theory because they can be coded into countable structures. Many important examples have been produced by constructing a family with certain computational properties, and then coding that family into

*We thank Uri Andrews, Steffen Lempp, Noah Schweber, and Mariya Soskova for helpful discussions.

[†]Greenberg and Turetsky were supported by a Marsden Fund grant #17-VUW-090.

[‡]Partially supported by an NSERC Banting postdoctoral fellowship.

a countable graph (often called the bouquet graph of that family, see e.g. [Kho86, AK00, GMS13]). For example, Goncharov constructed a computable structure of effective dimension 2 in this way [Gon80a, Gon80b].

Given a family of subsets of ω , we can ask: how hard is it to build a copy of that family? To build a copy of the family, we must build a copy of each set in the family, though it does not matter what order we present the sets in.

Definition 1.1. An enumeration of a countable family $\mathcal{F} \subseteq \mathcal{P}(\omega)$ is a set $W \subseteq \omega^2$ such that $\mathcal{F} = \{W^{[i]} : i \in \omega\}$, where $W^{[i]}$ is the i th column of W . The order of the columns in W and their multiplicity is not relevant. To avoid issues with multiplicity, we may always assume that in an enumeration of a family each column is replicated infinitely many times. A family \mathcal{F} is *computably enumerable* from a set X if there is an X -c.e. enumeration of \mathcal{F} .

A surprising result of Slaman [Sla98] and Wehner [Weh98] is that there is a family consisting of finite sets that can be enumerated by every non-computable degree, but which has no computable enumeration. Thus the problem of enumerating this family captures exactly the property of being non-computable.

We focus on families that are closed under finite differences.

Definition 1.2. We say that a family \mathcal{F} is closed under finite differences if whenever $A \in \mathcal{F}$ and $B =^* A$, $B \in \mathcal{F}$. (Here, $A =^* B$ means that A and B differ on only finitely many elements.)

There are natural places in which such families arise; for example natural computability-theoretic families such as the family of computable sets are closed under finite differences. The motivating example for us comes from torsion-free abelian groups. If G is a torsion-free abelian group, one can consider the family of sets $\{p \text{ prime} : p \mid g\}$ for $g \in G$. If G has no infinite divisibilities, then this family is closed under finite differences. One might hope to gain some insight into the following difficult open question by studying families closed under finite difference, though there are no formal implications in any direction:

Question 1.3. Is there a torsion-free abelian group that has a presentation computable in every non-computable degree but no computable presentation?

Our main goal in writing this paper was to build a Slaman–Wehner family that is closed under finite differences. Unfortunately, we are unable to solve this question.

Question 1.4. Is there a family closed under finite differences that can be enumerated by every non-computable degree, but which has no computable enumeration?

We call such a family a Slaman–Wehner family closed under finite differences. Standard forcing arguments (see Proposition 3.1) show that if there were such a family, then it would contain only c.e. sets, so we restrict our attention to those. Also, a theorem of Lachlan [Lac65] quickly yields that from an enumeration of a family closed under finite differences allowing repetition, we can compute a Friedberg enumeration, i.e. an enumeration with no repetition (Corollary 2.2). So, unlike families not closed under finite differences, we do not need to make any convention about how many times each set is enumerated in an enumeration.

Though this question remains open, we prove a number of interesting results. We have two negative results which place some restriction on such a family, should it exist. First, we show that any Slaman–Wehner family closed under finite differences must consist only of infinite sets. Second, we show that such a family cannot be enumerated in a uniform way:

Theorem 1.5. *Let \mathcal{F} be a family of sets closed under finite differences. Suppose that there is a c.e. operator Φ such that for every non-computable set X , Φ^X is an enumeration of \mathcal{F} . Then \mathcal{F} can be enumerated computably.*

It is interesting to note that there is a Slaman–Wehner family (not closed under finite differences) that has no uniform enumeration. This is an argument due to Faizrahmanov and Kalimullin [FK15]; the non-uniformity comes from the fact that every non-computable set, viewed as a binary expansion, is either not left-c.e. or not right-c.e. (or neither), but which of these is the case is not effective.

On the other hand, there are no obvious obstacles to such a family existing. We prove the existence of the following families:

Theorem 1.6. *There is a family closed under finite differences that can be enumerated uniformly by every non-low degree, but which cannot be enumerated by any low degree.*

Theorem 1.7. *Given a uniformly c.e. sequence $(D_e)_{e \in \omega}$ of non-computable c.e. sets, there is a family of c.e. sets closed under finite differences that can be enumerated by every D_e , but which cannot be enumerated computably.*

Theorem 1.8. *For any non-computable Δ_2^0 set X , there is a family of c.e. sets closed under finite differences that can be X -computably enumerated, but which cannot be enumerated computably.*

Combining the first and third of these positive results, note that for every non-computable set X , there is a family of c.e. sets closed under finite differences that can be X -enumerated, but which cannot be enumerated computably. So no single set forms a barrier to the existence of a Slaman–Wehner family closed under finite differences. The next natural question is whether there is a minimal pair:

Question 1.9. *Is there a pair of non-computable sets X and Y such that any family of c.e. sets closed under finite differences that can be enumerated by both X and Y can be enumerated computably?*

We know from Theorem 1.7 that no pair of c.e. sets is a minimal pair in this sense. Faizrahmanov has recently announced that there is a family of c.e. sets closed under finite differences which can be enumerated by every non-computable c.e. set, but which cannot be enumerated computably.

Notation

For a c.e. operator Φ , an oracle X , and $i \in \omega$, we let $\Phi^X[i] = \{n : (i, n) \in \Phi^X\}$ be the i^{th} column of Φ^X . For a string ξ and $k < |\xi|$ we let $\xi \upharpoonright k = \xi \upharpoonright (k + 1)$.

2 Friedberg Enumerations

Friedberg [Fri58] showed that there is an effective enumeration of the c.e. sets in which each set appears exactly once. More generally, a *Friedberg enumeration* or an *injective enumeration* of a family \mathcal{F} is an enumeration in which each set appears exactly once. There is a long history of providing sufficient conditions for a family with a computable enumeration to have a computable Friedberg enumeration. We will use the following condition due to Lachlan [Lac65, Lac67].

Given a finite set C , let $\mathcal{W}_C = \{W_x : x \in C\}$. A class \mathcal{F} has the property (E) if there is a binary partial computable function $\eta : [\omega]^{<\omega} \times [\omega]^{<\omega} \rightarrow \omega$ such that if C and D are finite sets and $\mathcal{W}_C \subseteq \mathcal{F}$, then $\eta(C, D)$ is defined if and only if

$$(\mathcal{F} - \mathcal{W}_C) \cap \{X : X \supseteq D\}$$

is non-empty, and in this case $\eta(C, D)$ is an index of a member of this class. Lachlan proved:

Theorem 2.1 (Lachlan [Lac65],[Lac67]). *An infinite c.e. class \mathcal{F} has the property (E) if and only if given a finite subclass \mathcal{G} of \mathcal{F} we can effectively enumerate $\mathcal{F} - \mathcal{G}$ without repetition.*

It is not hard to show that a family of infinite sets closed under finite differences satisfies condition (E), yielding:

Corollary 2.2. *Let \mathcal{F} be a family of sets closed under finite differences. From an enumeration of \mathcal{F} allowing repetition, we can compute an enumeration of \mathcal{F} with no repetition.*

Proof. If \mathcal{F} contains a finite set, then it contains every finite set. The standard Friedberg argument goes through. (See e.g. [Mal65].)

Now suppose that every set in \mathcal{F} is infinite. We relativise Lachlan's theorem to a given enumeration of \mathcal{F} . Fix a set $X \in \mathcal{F}$. Given finite sets C and D , for each $x \in C$, look for $y_x \in W_x - D$. Once we find such a y_x for each $x \in C$, which we eventually will if $\mathcal{W}_C \subseteq \mathcal{F}$, then define $\eta(C, D)$ to be an index for $X \cup D - \{y_x : x \in C\}$. This is a finite difference of the set X , and hence it is in \mathcal{F} . So \mathcal{F} has condition (E) relative to every enumeration of \mathcal{F} . \square

3 Negative Results

In this section we present our two negative results, i.e. results which say that families of a certain kind have computable enumerations, and hence cannot have Slaman–Wehner spectrum. These results are proved using the forcing relation; in all cases, when we say that a real is sufficiently generic, we mean with respect to Cohen forcing. We present the proof of the following well-known fact as a warm-up.

Proposition 3.1. *Let \mathcal{F} be a family of sets that can be enumerated by every degree $\mathbf{d} > 0$. Each set in \mathcal{F} is c.e.*

Proof. Let X be sufficiently generic relative to \mathcal{F} . Let Φ be a Turing functional such that Φ^X is an enumeration of \mathcal{F} . Fix i . Let $Y = \Phi^X[i]$. There is $\sigma \leq X$ such that $\sigma \Vdash Y = \Phi^X[i]$. Then

$$Y = \{n \mid \exists \rho \geq \sigma \text{ such that } \rho \Vdash n \in \Phi^G[i]\}.$$

The right-hand-side is c.e. \square

Our first negative result is that a family that is closed under finite differences and contains a (and hence every) finite set cannot have Slaman–Wehner spectrum: if it can be enumerated by every non-computable degree, then it has a computable enumeration.

Theorem 3.2. *Let \mathcal{F} be a family of sets, closed under finite differences, that can be enumerated by every degree $\mathbf{d} > 0$. If \mathcal{F} contains a finite set, then \mathcal{F} can be enumerated computably.*

Proof. The idea of the proof is as follows: \mathcal{F} can be enumerated by a generic set X via a function Φ , and there is an initial segment η of X that forces the fact that every set enumerated by Φ is a set in \mathcal{F} . Now for a given i , some of the extensions of η force that $\Phi^G[i]$ is a particular set W_e , and others do not; if an extension σ of η does not force the identity of $\Phi^G[i]$, then it has extensions σ_1 and σ_2 that force that $\Phi^G[i] = W_{e_1}$ and that $\Phi^G[i] = W_{e_2}$ respectively, with $W_{e_1} \neq W_{e_2}$. For each σ and i , we will enumerate a set $Q_{i,\sigma}$. By guessing at whether or not σ forces that $\Phi_e^G[i] = W_e$ for some e , we will be able to make $Q_{i,\sigma}$ equal to W_e if σ does force this fact, and finite otherwise. Since \mathcal{F} contains the finite sets, it will not be a problem when $Q_{i,\sigma}$ is finite.

Let X be sufficiently generic, and let Φ be a Turing functional such that Φ^X is an enumeration of \mathcal{F} . There is an initial segment of X that forces this fact. We will assume for convenience that this initial segment is the empty string; otherwise, we just work among extensions of this string.

For each σ and i , we will define a set $Q_{i,\sigma}$ such that either $Q_{i,\sigma}$ is finite, or $\sigma \Vdash \Phi^G[i] = W_e$ and $Q_{i,\sigma} = W_e$. Define $Q_{i,\sigma}$ as follows. Put $n \in Q_{i,\sigma}$ if:

1. for all $m < n$, if $m \in \Phi^\tau[i]$ for some $\tau \geq \sigma$ of length $|\sigma| + n$, then for every $\tau \geq \sigma$ of length $|\sigma| + n$ there is $\rho \geq \tau$ with $m \in \Phi^\rho[i]$, and
2. there is a string $\rho \geq \sigma$ with $n \in \Phi^\rho[i]$.

Suppose that $\sigma \Vdash \Phi^G[i] = W_e$ for some e . It is not hard to see that $Q_{i,\sigma} \subseteq W_e$. On the other hand, suppose that $n \in W_e$. We show that $n \in Q_{i,\sigma}$, so that $Q_{i,\sigma} = W_e$. For (1): Suppose that $m < n$ is such that $m \in \Phi^\tau[i]$ for some τ of length $|\sigma| + n$; then $m \in W_e$. Given any τ of length $|\sigma| + n$, there is a sufficiently generic $Y \geq \tau$. Then $\Phi^Y[i] = W_e$ and so $m \in \Phi^Y[i]$. Thus there is $\rho \geq \tau$ with $m \in \Phi^\rho[i]$. For (2): There is $Y \geq \sigma$ sufficiently generic, and so $\Phi^Y[i] = W_e$. So there is $\rho \geq \sigma$ with $n \in \Phi^\rho[i]$.

On the other hand, suppose that there is no e for which $\sigma \Vdash \Phi^G[i] = W_e$. Then there are $Y, Z \geq \sigma$ sufficiently generic with $\Phi^Y[i] \neq \Phi^Z[i]$. Suppose without loss of generality that there is $m \in \Phi^Y[i] - \Phi^Z[i]$. Let $\tau_Y \leq Y$ be such that $m \in \Phi^{\tau_Y}[i]$, and let $\tau_Z \leq Z$ be such that $\tau_Z \Vdash m \notin \Phi^G[i]$; so there is no extension ρ of τ_Z with $m \in \Phi^\rho[i]$. Let $N = \max(m, |\tau_Y|, |\tau_Z|)$. Then no $n \geq N$ satisfies (1), and so no element of $Q_{i,\sigma}$ is larger than N . Thus $Q_{i,\sigma}$ is finite.

In either case, $Q_{i,\sigma}$ is in \mathcal{F} . It remains to show that each element of \mathcal{F} is one of the $Q_{i,\sigma}$. Fix W_e in \mathcal{F} . Let X be sufficiently generic, and let i be such that $\Phi^X[i] = W_e$. Let $\sigma \leq X$ be such that $\sigma \Vdash \Phi^G[i] = W_e$. Then $Q_{i,\sigma} = W_e$. \square

Now we will show that there is no Slaman–Wehner family closed under finite differences with this witnessed in a uniform way. The families constructed by Slaman and Wehner were uniform, so if there is a Slaman–Wehner family closed under finite differences it must be constructed in a different way. By “uniform”, we mean that there is an operator Φ such that Φ^X enumerates the family whenever X is non-computable.

Theorem 3.3. *Let \mathcal{F} be a family of sets, closed under finite differences, that can be enumerated uniformly by every degree $\mathbf{d} > 0$. Then \mathcal{F} can be enumerated computably.*

Proof. Let Φ be a Turing functional which witnesses that \mathcal{F} can be uniformly enumerated by every non-computable degree. Uniformly in each σ and i , we will enumerate a set $Q_{i,\sigma}$. For some 1-generic $X \geq \sigma$, we will have $Q_{i,\sigma} =^* \Phi^X[i] \in \mathcal{F}$. Before constructing the sets $Q_{i,\sigma}$, we will show how they suffice to prove the theorem.

For each $W \in \mathcal{F}$, we argue that there are i and σ such that $W =^* Q_{i,\sigma}$, so that if we take the family $\{Q_{i,\sigma} \mid i \in \omega, \sigma \in 2^{<\omega}\}$ and close it under finite differences we get an enumeration of \mathcal{F} . Let Y be sufficiently generic. There is i such that $W = \Phi^Y[i]$. So there is $\sigma \leq Y$ such that $\sigma \Vdash W = \Phi^G[i]$. Let $X \geq \sigma$ be a 1-generic such that $Q_{i,\sigma} =^* \Phi^X[i]$. We claim that $W = \Phi^X[i]$. Indeed, suppose that $n \in \Phi^X[i]$. Then there is $\rho \geq \sigma$ such that $\rho \Vdash n \in \Phi^G[i]$, and so there is a sufficiently generic $Z \geq \rho$ with $n \in \Phi^Z[i] = W$. A similar argument works when $n \notin \Phi^X[i]$.

We now construct $Q_{i,\sigma}$ and an approximation to a 1-generic $X \geq \sigma$ with $Q_{i,\sigma} =^* \Phi^X[i]$. Let $(W_e)_{e \in \omega}$ be an enumeration of the c.e. sets. For simplicity, we will write $\Psi^X = \Phi^X[i]$.

We can think of the argument as a failed diagonalization argument. We will attempt to construct a set Y to meet the following requirements:

$$R_e : \Psi^Y \neq W_e.$$

Of course, we will not be able to meet all of these requirements, as no matter what Y is, Ψ^Y is c.e. If Y is computable, Φ^Y is c.e., and if Y is non-computable, then Φ^Y is in the family \mathcal{F} and hence is c.e. (We use Y here because the 1-generic set X we want will be obtained by combining the approximation to Y with that of a 1-generic Z .)

The possible outcomes of a requirement R_e are the infinitary outcome ∞ , and finitary outcomes $\boxed{n \notin W_e, \tau}$ and $\boxed{n \in W_e, \tau}$. An outcome $\boxed{n \notin W_e, \tau}$ represents that $n \notin W_e$ and $n \in \Psi^\tau$. An outcome $\boxed{n \in W_e, \tau}$ represents that $n \in W_e$ and for all $\rho \geq \tau$, $n \notin \Psi^\rho$. These two types of outcomes are finitary because we will leave $\boxed{n \notin W_e, \tau}$ when n enters W_e and we will leave $\boxed{n \in W_e, \tau}$ when we find a $\rho \geq \tau$ with $n \in \Psi^\rho$. If either of the finitary outcomes is the true outcome, then R_e will be satisfied; but in the infinitary outcome, R_e will not be satisfied.

For a strategy ξ on the tree, we also define the (finite) binary string $y(\xi)$ which is the initial segment of Y determined by ξ . The empty sequence $\langle \rangle$ is a strategy and $y(\langle \rangle) = \sigma$. Suppose that we have determined that ξ , of length e , is a strategy. If the last entry of ξ is ∞ , then ξ is terminal — no proper extensions of ξ are strategies. Otherwise, the possible outcomes of ξ on the tree of strategies are ∞ , and the outcomes $\boxed{n \in W_e, \tau}$ and $\boxed{n \notin W_e, \tau}$ with $\tau \geq y(\xi)$. We let $y(\xi \hat{\ } \infty) = y(\xi)$. For all n and $\tau \geq y(\xi)$, we let $y(\xi \hat{\ } \boxed{n \in W_e, \tau}) = y(\xi \hat{\ } \boxed{n \notin W_e, \tau}) = \tau$.

At each stage, we will define a strategy ξ_s and let $y_s = y(\xi_s)$. We will let ξ , the true path, be the path of strategies leftmost visited infinitely often (where the outcome ∞ lies to the left of the others). We will argue that ξ is in fact finite (it is terminal, with last entry ∞); otherwise, we show that $Y = y(\xi)$ meets all requirements, which is impossible. The last entry of ξ indicates the least e for which the requirement R_e is not met. Each R_i , $i < e$, will be

satisfied with one of the finitary outcomes, and so once these have stabilized, say at stage s^* , we will have $y(\xi) \leq y_s$ for all $s \geq s^*$, and $y_s = y(\xi)$ whenever $s \geq s^*$ is a true stage. (This will all be verified after the construction.)

Letting Z be a Δ_2^0 1-generic with computable approximation $(z_s)_{s \in \omega}$, our 1-generic $X \geq \sigma$ will be $X = y(\xi) \hat{\ } Z$. We have an approximation to X given by $x_s = y_s z_s$. (Note that this is not a Δ_2^0 approximation; we cannot specify $y(\xi)$ as a parameter, as the construction has to be uniform in i and σ). Using the fact that we failed to satisfy R_e , we will show in Claim 3 that defining $Q_{i,\sigma} = \{n : \exists s \ n \in \Psi_s^{x_s}\}$, we get $Q_{i,\sigma} =^* \Psi^X = W_e$.

Construction.

Stage 0: Begin with $\xi_0 = \langle \infty \rangle$.

Stage $s + 1$: a strategy $\zeta < \xi_s$ is *discovered to be incorrect* at stage $s + 1$ if, letting $e = |\zeta|$, we have:

- (1) $\xi_s(e) = \boxed{n \notin W_{e,\tau}}$ and $n \in W_{e,s}$;
- (2) $\xi_s(e) = \boxed{n \in W_{e,\tau}}$ and $n \in \Psi_s^\rho$ for some $\rho \geq \tau$; or
- (3) $\xi_s(e) = \infty$, and there are some n and $\tau \geq y(\zeta)$ (with $n, |\tau| \leq s$) such that either
 - (i) $n \notin W_{e,s}$ and $n \in \Psi_s^\tau$; or
 - (ii) $n \in W_{e,s}$ and $n \notin \Psi_s^\rho$ for all $\rho \geq \tau$ (of length $\leq s$).

If no $\zeta < \xi_s$ is discovered to be incorrect at stage $s + 1$, then we let $\xi_{s+1} = \xi_s$. Otherwise, let ζ be the shortest initial segment of ξ_s discovered to be wrong at stage $s + 1$. We define ξ_{s+1} as follows:

- In cases (1) and (2), i.e., when $\xi_s(e) \neq \infty$, we let $\xi_{s+1} = \zeta \hat{\ } \infty$.
- In sub-case (3i), choosing the least τ and n , we let $\xi_{s+1} = \zeta \hat{\ } \boxed{n \notin W_{e,\tau}} \hat{\ } \infty$.
- In sub-case (3ii), again choosing the least τ and n , we let $\xi_{s+1} = \zeta \hat{\ } \boxed{n \in W_{e,\tau}} \hat{\ } \infty$.

End construction.

Verification.

Let ξ be the leftmost path visited infinitely often. Then either ξ is a finite, terminal strategy, or it is infinite (and so does not contain an entry ∞). We will shortly see that it is the former.

We note that if $\zeta < \xi$ is non-terminal, then $\zeta < \xi_s$ for all but finitely many stage s ; for every $e < |\zeta|$, once $\zeta \upharpoonright e$ has stabilised, once the outcome $\zeta(e)$ is chosen, it will never be un-chosen, as we cannot return to a discarded outcome.

If ξ is infinite, let $y(\xi) = \bigcup_{\zeta < \xi} y(\zeta)$; if ξ is finite, then $y(\xi)$ is already defined. We are in the slightly notationally awkward situation where we do not yet know whether ξ and $y(\xi)$ are finite or infinite strings, so in the next lemma we will have to talk about strings $Y \in 2^\omega$ with $Y \geq y$ where it might be that $Y = y(\xi)$ if y is infinite.

Claim 1. *If $e < |\xi|$ and $\xi(e) \neq \infty$ then $\Psi^Y \neq W_e$ for any $Y \geq y(\xi)$.*

Proof. As discussed, there is some stage after which $\xi_t \uparrow e = \xi \uparrow e$.

First suppose that $\xi(e) = \boxed{n \notin W_e, \tau}$. Then $n \notin W_e$, since if we found that $n \in W_e$, we would never again have this outcome. Also, when we first had this outcome, we had $n \in \Psi^\tau$. Since $Y \geq \tau$, $\Psi^Y \neq W_e$.

Now if $\xi(e) = \boxed{n \in W_e, \tau}$, then $n \in W_e$, but for all $\rho \geq \tau$, $n \notin \Psi^\rho$. Since $Y \geq \tau$, $n \notin \Psi^Y$ and so $\Psi^Y \neq W_e$. \square

If $Y \geq y(\xi)$, then Ψ^Y is a c.e. set W_e (either for obvious reasons because Y is computable, or if Y is non-computable, then Ψ^Y is in the family \mathcal{F} and hence a c.e. set). So as a consequence of this lemma, ξ must be finite, with last entry ∞ , and $y := y(\xi)$ is finite as well. Let s_0 be a stage after which $\xi^- < \xi_s$ (where ξ^- is the result of removing the last entry ∞ from ξ). So $y(\xi) \leq y_s$ for all $s \geq s_0$. We say that a stage $s \geq s_0$ is *true* if $\xi_s = \xi$. There are infinitely many true stages.

Claim 2. *Let $e = |\xi| - 1$ (so $\xi(e) = \infty$). Let $Y \geq y$ be a 1-generic extending y . Then $\Psi^Y = W_e$.*

Proof. If there is $n \in \Psi^Y - W_e$, then there is some $\tau \leq Y$ with $n \in \Psi^\tau$. Then the outcome $\xi(e)$ would be $\boxed{n \notin W_e, \tau}$ for some such pair n, τ .

If there is $n \in W_e - \Psi^Y$, then since Y is 1-generic, there is a τ such that $y \leq \tau < Y$ and for all $\rho \geq \tau$, $n \notin \Psi^\rho$. Then the outcome $\xi(e)$ would be $\boxed{n \in W_e, \tau}$ for some such pair n, τ . \square

Let $(z_s)_{s \in \omega}$ be a Δ_2^0 approximation of a 1-generic Z . Let $X = y \hat{\ } Z$ and $x_t = y_t z_t$; X is a 1-generic extending y . For each m , there are infinitely many stages $t \geq s_0$ such that $x_t = y_t z_t \geq X \upharpoonright_m$, indeed this holds for almost all true stages.

Claim 3. *Let $Q_{i,\sigma} = \{n : \exists s \ n \in \Psi_s^{x_s}\}$. Then $Q_{i,\sigma} =^* \Psi^X = W_e$.*

Proof. Let e be as in the previous claim. Let $Q_{i,\sigma}[t] = \{n : \exists s \leq t \ n \in \Psi_s^{x_s}\}$. Each of these sets is finite, and $Q_{i,\sigma} = \bigcup_t Q_{i,\sigma}[t]$.

For every stage $t \geq s_0$, if $n \in Q_{i,\sigma}[t] - Q_{i,\sigma}[t-1]$, then $n \in \Psi_t^{x_t}$. Since $t \geq s_0$, $x_t \geq y$. This shows that $n \in \Psi_t^\tau$ for some $\tau \geq y$. There is some 1-generic $G \geq \tau$; so $n \in \Psi^G$; by the previous claim, $\Psi^G = W_e = \Psi^X$. So $Q_{i,\sigma} \subseteq^* \Psi^X$.

On the other hand, if $n \in \Psi^X$, there is some t such that $n \in \Psi_t^{x_t}$. As $X = y \hat{\ } Z$ is a 1-generic extending y , $\Psi^X = W_e$. So $\Psi^X = W_e \subseteq Q_{i,\sigma}$. \square

We have shown that $Q_{i,\sigma} =^* \Psi^X = \Phi^X[i]$ for some 1-generic X extending σ . By the arguments given at the beginning of the proof, if we take the family $\{Q_{i,\sigma} \mid i \in \omega, \sigma \in 2^{<\omega}\}$ and close it under finite differences, we get an enumeration of \mathcal{F} . \square

The next lemma gives another restriction on what a Slaman-Wehner family closed under finite differences would have to look like. Let A be a low set that enumerates such a family. Each set in the family has a computable enumeration, but also an enumeration relative to A as part of the enumeration of the family by A . We show that for at least one set in the family, the enumeration relative to A must be faster than any computable enumeration.

Lemma 3.4. *Let A be a low set, and \mathcal{F} a family of c.e. sets closed under finite differences that can be enumerated by A using Φ . Suppose that for each i , for some e with $\Phi^A[i] = W_e$ we have $\Phi_s^A[i] \subseteq W_{e,s}$ for all s . Then \mathcal{F} can be computably enumerated.*

Proof. Let $\pi \in 2^\omega$ be such that $\pi(e, i) = 1$ if and only if $(\forall n)(\forall s) n \in \Phi_s^A[i] \Rightarrow n \in W_{e,s}$. Since A is low, $\pi \leq \emptyset'$. Let σ_s be a computable sequence of finite binary strings that approximate π , and such that there are infinitely many true stages s with $\sigma_s < \pi$.

We will uniformly compute sets $U_i =^* \Phi^A[i]$. Let $U_{i,s+1} = U_{i,s} \cup [\{0, \dots, s\} \cap_{\sigma_s(e,i)=1} W_{e,s}]$ and let $U_i = \bigcup U_{i,s}$. Fix a particular index e' such that $\Phi^A[i] = W_{e'}$ and for all s we have $\Phi_s^A[i] \subseteq W_{e',s}$. Let s' be a stage such that for all $s \geq s'$, $\sigma_s(e', i) = 1$. Then for all $s \geq s'$, $U_{i,s+1} - U_{i,s} \subseteq W_{e'}$ and so $U_i \subseteq^* W_{e'}$. Now suppose that $n \in \Phi_s^A[i]$, $s \geq s'$, and s is a true stage of the approximation σ_s of π , so that $\sigma_s < \pi$. Then $n \in \bigcap_{\sigma_s(e,i)=1} W_{e,s}$ and so $n \in U_i$. Thus $U_i =^* W_{e'} = \Phi^A[i]$. \square

4 Examples of Families

In this section, we produce the three examples of families of c.e. sets, closed under finite difference, that were promised in the introduction: (1) A family that can be enumerated exactly by the non-low degrees; (2) For any effective list of non-computable c.e. sets, a family that can be enumerated by those sets but not computably enumerated; and (3) For any non-computable Δ_2^0 set, a family that can be enumerated by that set but which cannot be computably enumerated. Recall that in all of these cases, our family consists of c.e. sets, so that in (3) for example we cannot simply take the family to consist solely of the given non-computable Δ_2^0 set.

We begin by adapting Wehner's construction to produce a family of sets closed under finite differences that can be enumerated exactly by the non-low degrees. We can think of this as essentially a jump inversion of Wehner's construction relativized to $\mathbf{0}'$.

Theorem 1.6. *There is a family closed under finite differences that can be enumerated uniformly by every non-low degree, but which cannot be enumerated by any low degree.*

Proof. Following [GMS13], relativizing Wehner's construction of a family that can be enumerated by exactly the non-computable degrees, we get a family

$$\mathcal{F}_{\emptyset'} = \{\{e\} \oplus F : e < \omega, F \subseteq \omega \text{ finite, and } F \neq W_e^{\emptyset'}\}.$$

Then X can enumerate \mathcal{F} if and only if $X \not\leq_T \emptyset'$. (See Propositions 2.3 and 2.4 of [GMS13].)

Now, given a set S , define

$$U_S = \{\langle s, n \rangle : s \notin S\}.$$

We have the following jump inversion result using U_S :

Lemma 4.2. *For finite sets S , we can uniformly pass between an X -index i for an enumeration of a set $W_i^X =^* U_S$ and an X' -index j for $W_j^{X'} = S$.*

Proof. Given $W_i^X =^* U_S$, X' can enumerate S by, for each $s \in \omega$, searching for n_0 such that for all $n \geq n_0$, $\langle s, n \rangle \notin W_e^X$, and enumerating s if such an n_0 is found.

On the other hand, if $W_j^{X'} = S$, then S is Σ_2^0 relative to X . So we can find an X -computable function f such that if $s \in S$, $W_{f(s)}^X$ is finite, and if $s \notin S$, $W_{f(s)}^X = \omega$. Let

$$V = \{\langle s, n \rangle : n \in W_{f(s)}^X\}$$

Since S is finite, $V =^* U_S$. □

Define

$$\mathcal{G} = \{V : \exists S \in \mathcal{F}_{\emptyset'} [V =^* U_S]\}.$$

We claim that \mathcal{G} is uniformly enumerable by all non-low degrees, and is not enumerable from any low degree.

If X is non-low, then $X' >_T \emptyset'$. So X' uniformly enumerates $\mathcal{F}_{\emptyset'}$, and by Lemma 4.2, X uniformly enumerates a family which, when we close under $=^*$, will give us \mathcal{G} .

If X is low, then any enumeration of \mathcal{G} would give an X' -listing of $\mathcal{F}_{\emptyset'}$, by Lemma 4.2, which is impossible. □

Next we consider an effective sequence of non-computable c.e. sets.

Theorem 1.7. *Given a uniformly c.e. sequence $(D_e)_{e \in \omega}$ of non-computable c.e. sets, there is a family of c.e. sets closed under finite differences that can be enumerated by every D_e , but which cannot be enumerated computably.*

Proof. For each i , let Q_i be a computable set of indices such that $\{W_k : k \in Q_i\}$ is the i th computable enumeration of a family of sets. When we construct our family \mathcal{F} , for each set in \mathcal{F} there will be some i such that the set has infinitely many elements of the form $\langle i, \cdot \rangle$ and only finitely many elements of the form $\langle j, \cdot \rangle$ for $j \neq i$. We will diagonalize against the family $\{W_k : k \in Q_i\}$ using the sets containing only elements of the form $\langle i, \cdot \rangle$.

Let P_i be a Π_1^0 set of indices for the c.e. sets $\{n : \langle i, n \rangle \in W_k\}$ where $k \in Q_i$ and W_k contains only elements of the form $\langle i, n \rangle$. The following lemma contains the heart of the argument: that we can diagonalize against this family.

Lemma 4.4. *Given a Π_1^0 set $P \subseteq \omega$, there is a family of c.e. sets \mathcal{F} closed under finite differences such that*

$$\mathcal{F} \neq \{W_e : e \in P\}$$

and \mathcal{F} can be enumerated by every D_e . Moreover, we can build \mathcal{F} uniformly from an index for P , and the enumeration of \mathcal{F} from the D_e is uniform as well.

Before proving the lemma, we show how to use it to finish the theorem. For each i , let \mathcal{F}_i be the family of c.e. sets obtained by this lemma applied to P_i . The families \mathcal{F}_i can be enumerated uniformly by every D_e , and $\mathcal{F}_i \neq \{W_k : k \in P_i\}$. Now we need to combine each of these families into a single family \mathcal{F} ; let \mathcal{F} be the closure under taking finite differences of the family

$$\{\{\langle i, n \rangle : n \in W\} : W \in \mathcal{F}_i\}.$$

Then \mathcal{F} can be enumerated by each D_e , because D_e can enumerate each \mathcal{F}_i uniformly. Also, we argue that \mathcal{F} is not the same as the i th computable enumeration $\{W_k : k \in Q_i\}$. Note

that \mathcal{F}_i is the family of all sets $\{n : \langle i, n \rangle \in W\}$ for $W \in \mathcal{F}$ containing only elements of the form $\langle i, \cdot \rangle$. Since $\mathcal{F}_i \neq \{W_k : k \in P_i\}$, and P_i was obtained from Q_i by the same process that \mathcal{F}_i is obtained from \mathcal{F} , $\mathcal{F} \neq \{W_k : k \in Q_i\}$. \square

Now we prove Lemma 4.4. As one might expect, this is a c.e. permitting argument.

Proof of Lemma 4.4. For each index e , we build a D_e -enumeration of a family \mathcal{F}_e of c.e. sets. We will ensure that the families \mathcal{F}_e and $\mathcal{F}_{e'}$ enumerated by D_e and $D_{e'}$ are the same; we call this family \mathcal{F} . We will also make sure that \mathcal{F} is different from $\{W_e : e \in P\}$. Though they will be the same family, the correspondence between the sets in \mathcal{F}_e and the sets in $\mathcal{F}_{e'}$ may not be computable, and this is why we give them different names.

First it will be helpful to think of what the construction does in the case of a single non-computable c.e. set D_e . We will describe a restriction of the general strategy to this simpler case. The family \mathcal{F}_e will consist of a single set U_e and its finite differences. We want to ensure that either U_e is not in $\{W_k : k \in P\}$, or that there is some set in $\{W_k : k \in P\}$ that is different from U_e and all of its finite differences. What we will do is to consider each W_k in turn, and try to make U_e different from W_k . We use the elements of the form $\langle k, \cdot \rangle$ for the sake of W_k .

For each k in turn, we do the following, beginning with $k = 0$. We call this process the *module for W_k* . Suppose that the module for W_k begins at stage s . If $k \notin P_s$, then we do not have to make U_e different from W_k , and we can just go on to W_{k+1} . Otherwise, if $k \in P_s$, begin by putting $\langle k, n \rangle$ into U_e for each n , with use $D_{e,s} \upharpoonright n$ where s is the current stage. At each later stage t , do one of the following:

- If $k \notin P_t$, then this means that we no longer have to make U_e different from W_k . We put $\langle k, n \rangle$ into U_e with no use, and go on to the module for W_{k+1} .
- If there is some n that entered D_e at stage t , and there is $m \geq n$ with $\langle k, m \rangle \in W_{k,t}$, then we have received permission from D_e to diagonalize against W_k . For $n' < n$, put $\langle k, n' \rangle$ into U_e with no use. On the other hand, because n has entered D_e , we are able to remove $\langle k, m \rangle$ from U_e . So $U_e \neq W_k$. We go on to the module for W_{k+1} .
- Otherwise, it might be that D_e has changed, but we have not been given an opportunity to diagonalize. Put $\langle k, n \rangle$ into U_e with use $D_{e,t} \upharpoonright n$ and continue the module for W_k at stage $t + 1$.

If we end the module for W_k for any reason, then we have succeeded against W_k , either because $k \notin P$ and we do not need to do anything, or because we have ensured that $U_e \neq W_k$. It is also possible that for some k , the module for W_k never ends. In this case, we will argue that W_k is not a finite difference of U_e . Suppose that it was; then U_e contains every element $\langle k, n \rangle$, and so W_k must contain all but finitely many of those elements. But then by a standard permission argument, there must be some n that enters D_e at a stage t at which $\langle k, m \rangle \in W_{k,t}$ for some $m \geq n$; otherwise, we would be able to compute D_e . This gives us a contradiction, because if this ever happened then the module for W_k would have been given permission.

We also need to make sure that the set U_e is c.e. (and in fact computable). For each k , U_e contains either finitely many elements $\langle k, n \rangle$, or every element $\langle k, n \rangle$. If the module for

W_k never ends, then U_e is actually computable using the knowledge of how the modules for W_0, \dots, W_k ended. Otherwise, each module ends; to decide which elements $\langle k, n \rangle$ to put into U_e , we can wait until the module for W_e ends, at which point the construction fixes how many of those elements to put into U_e .

Before returning to the general case, it will be helpful to show how we may assume that we can delete sets from our enumerations of the families \mathcal{F}_e . When we delete a set, we will delete its entire equivalence class under finite differences. We do this as follows. Reserve infinitely many elements $\langle 0, n \rangle$ as garbage labels. Whenever we want to delete a set from the family \mathcal{F}_e , we put every label, including all of the garbage labels, into it; we also add all of these labels onto finite differences of the set, and add new sets to \mathcal{F}_e that are the finite differences of these sets. Since there are infinitely many garbage labels, each of these sets will still have infinitely many garbage labels. Instead of working with $\{W_e : e \in P\}$, we work with


$$\{W_e : e \in P \text{ and } W_e \text{ does not contain any garbage labels}\}.$$

This is still a Π_1^0 set of indices. If we can make the family of sets in \mathcal{F} not including a garbage label different from this family, then \mathcal{F} together with the deleted sets will be different from $\{W_e : e \in P\}$. So for the rest of the construction, we will simply allow ourselves to delete sets from the families \mathcal{F}_e (without talking about garbage labels).

Now we return to the full sequence $(D_e)_{e \in \omega}$. Now the main issue that arises is that the different sets D_e might give us permission at different times, which would mean e.g. that D_e would give us permission to remove $\langle k, n \rangle$ from U_e , but $D_{e'}$ would not give us permission to remove $\langle k, n \rangle$ from $U_{e'}$; and then later $D_{e'}$ would give us permission to remove some other $\langle k, m \rangle$ from $U_{e'}$, but D_e would not give us permission to remove it from U_e .

The solution to this is to take advantage of the fact that U_e and $U_{e'}$ do not have to be the same set, as long as there is some other set in \mathcal{F}_e that is equal to $U_{e'}$, and some other set in $\mathcal{F}_{e'}$ that is equal to U_e . Moreover, which set this is does not have to be fixed. Just before the start of the module for W_k , each \mathcal{F}_e will consist only of the set U_e (and of course, its finite differences), and we will have $U_e = U_{e'}$ for each e, e' . At the start of the module for

W_k , we will add new sets $V_e^{e'}$ to \mathcal{F}_e with the intent of copying $U_{e'}$.



Note that the subscripts of the sets denote which family they belong to, and the superscripts denote the set they are copying. We use the elements $\langle k, e, n \rangle$ for U_e in the module for W_k . When D_e gives permission to make $U_e \neq W_k$ by removing an element $\langle k, e, n \rangle$ from U_e , this element was never put in $U_{e'}$ for $e' \neq e$; so we can delete all of the sets V and make $U_e = U_{e'}$ again for each e, e' . So if each module ends, then we end up with $U_e = U_{e'}$; but if the module for W_k never ends, then we have $U_e \neq U_{e'}$ for each e, e' , but $V_e^e = U_e$.

For a particular value of k , we act against W_k using the following module. Assume that at each stage only a single element enters exactly one of D_0, D_1, \dots . At the start of the module, each \mathcal{F}_e will consist of only a single set U_e (plus its finite differences); the sets U_e will contain exactly the same elements, and will contain only elements of the form $\langle k', \cdot, \cdot \rangle$ for $k' < k$.

We will describe below the module for W_k as it builds each \mathcal{F}_e . The events that trigger the beginning and ending of each module will be computable. The modules will build the families \mathcal{F}_e as c.e. operators, the e th one with oracle D_e , describing elements enumerated into sets in the families with various uses.

Module for W_k :

Suppose that the module for W_k begins at stage s . As described, at stage s , each \mathcal{F}_e consists of a set U_e (and its finite differences). We begin the module by adding to each family \mathcal{F}_e infinitely many sets $V_e^{e'}$, all equal to the U_e 's, again with no use.

Now let $t \geq s$ be a stage, and suppose that the module for W_k is still running at stage t . We do the following:

- If $k \notin P_t$:
 - delete all the sets $V_e^{e'}$ from all the families \mathcal{F}_e ;
 - with no use, put $\langle k, e', n \rangle$ into U_e for each n and e' (including $e' = e$);

- end the module.
- Otherwise, suppose that some n^* enters a set D_{e^*} at stage t (recall there will be only one such pair (n^*, e^*) at each stage), and there is $m \geq n^*$ with $\langle k, e^*, m \rangle \in W_{k,t}$:
 - delete all the sets $V_e^{e'}$ from all the families \mathcal{F}_e ;
 - for every $e' \neq e^*$, every e and every n , put $\langle k, e', n \rangle$ into U_e with no use;
 - for every $n < n^*$ and every e , put $\langle k, e^*, n \rangle$ into each U_e with no use;
 - end the module.
- Otherwise, for each e and n , put $\langle k, e, n \rangle$ into U_e with use $D_{e,t} \Vdash n$. Continue the module.

The complete construction piecing together the modules is as follows:

Construction: Run the module for W_0 ; if it returns, run the module for W_1 , and when that returns run the module for W_2 , and so on.

Now we must check that the construction works. We make a few remarks about the state of the construction whenever a module finishes. At the end of any module:

- each \mathcal{F}_e consists of only the set U_e (and finite differences);
- every element in U_e is in U_e with no use;
- $U_e = U_{e'}$ for each e, e' .

Claim 1. *For each e, e' , \mathcal{F}_e and $\mathcal{F}_{e'}$ are the same family \mathcal{F} .*

Proof. Suppose first that each module ends. Then by the remark above, each \mathcal{F}_e consists of only of the finite differences of a single set U_e , and $U_e = U_{e'}$ for each e, e' .

So now suppose that the module for W_k does not end. Then each \mathcal{F}_e will consist of a set U_e and sets $V_e^{e'}$ for $e' \neq e$, and all the finite differences. Moreover, after the beginning of the module no elements are ever removed from $U_{e'}$ (since, if an element would be removed by a change in $D_{e'}$, it is added back into $U_{e'}$ with the new use). So we have $V_e^{e'} = U_{e'}$. Thus the families \mathcal{F}_e are all the same. \square

Claim 2. $\mathcal{F} \neq \{W_k : k \in P\}$.

Proof. We have two cases. First suppose that each module ends. Then \mathcal{F} will consist of the equivalence class of a single set U (with $U = U_e$ for each e). We claim that $U \neq W_k$ for any $k \in P$, so that $U \notin \{W_k : k \in P\}$. Indeed, for each k , the module for W_k ends, either because we find that $k \notin P$, or because there is n that enters D_e at a stage t , and there is $m \geq n$ with $\langle k, e, m \rangle \in W_{k,t}$. In the latter case, $\langle k, e, m \rangle \in W_k$ but we remove $\langle k, e, m \rangle$ from U ; so $W_k \neq U$.

Now suppose that the module for W_k does not end. Then we claim that $W_k \notin \mathcal{F}$. The family \mathcal{F} consists of the sets U_e and their finite differences; each U_e contains the elements $\langle k, e, n \rangle$ for all n while not containing $\langle k, e', n \rangle$ for any $e' \neq e$. Suppose to the contrary that $W_k \in \mathcal{F}$, so that for some e , $W_k =^* U_e$. Then choose N such that for all $n \geq N$, $\langle k, e, n \rangle \in W_k$.

Since the module for W_k does not end, whenever n enters D_e at a stage t , there is no $m \geq n$ with $\langle k, e, m \rangle \in W_{k,t}$. Thus $D_e \upharpoonright_n = D_{e,t} \upharpoonright_n$ if there is $m \geq n$ with $\langle k, e, m \rangle \in W_{k,t}$. This allows us to compute the non-computable set D_e . From this contradiction we can conclude that $W_k \notin \mathcal{F}$ \square

This completes the proof of the lemma. \square

It would be natural to try to extend this argument to all of the non-computable c.e. sets by showing that for any list $(D_e)_{e \in \omega}$ of c.e. sets, some of which might be computable, there is a family \mathcal{F} of c.e. sets closed under finite differences which has no computable enumeration, but has an enumeration from any non-computable D_e . The problem we run into is that during the module for W_k , W_k could copy U_e for some e with D_e non-computable, so that we are never given permission. One would then have to add a guessing argument to guess at when D_e is non-computable, but we could not make this work with the sets $V_{e'}$. We leave this question open:

Question 4.5. If \mathcal{F} is a family of c.e. sets closed under finite differences, and it can be enumerated by every non-computable c.e. set, must it have a computable enumeration?

One way to show that there is no Slaman–Wehner family closed under finite differences would be to show that there is a degree \mathbf{d} such that any family of c.e. sets closed under finite differences that can be enumerated by \mathbf{d} has a computable enumeration. We show that this is not the case. We already know that such a family exists for any non-low degree, and now we show that such a family exists for any Δ_2^0 degree. Note that if we do not require the family to consist of c.e. sets, then it is not hard to construct such a family; so this is the main difficulty.

Theorem 1.8. *For every Δ_2^0 set D , there is a family \mathcal{F} of c.e. sets closed under finite differences that can be enumerated by D , but cannot be enumerated computably.*

Proof. Let D be Δ_2^0 but not computable. The following lemma will give us a strategy for diagonalizing against a single set:

Lemma 4.7. *There is, uniformly in e , a set U_e^D that is c.e. with $U_e^D \neq^* W_e$.*

Before proving the lemma, we show how to use it to prove the theorem. Let $\{W_{f(n,i)}\}_{i \in \omega}$ be an enumeration of the n th computably enumerable family of sets. Let $W_{g(n,i)} = \{x : \langle 1, i, x \rangle \in W_{f(n,i)}\}$. Let \mathcal{F} consist of, for each n , the set A_n defined as follows, together with all finite differences. Let i_0, i_1, \dots be an enumeration of the indices i for sets $W_{f(n,i)}$ that contain an element of the form $\langle 0, n, m \rangle$ for some m . The set A_n contains:

1. $\langle 0, n, m \rangle$ for each $m \in \omega$;
2. $\langle 1, i_\ell, x \rangle$ for $x \in U_{g(n,i_\ell)}^D$, if $W_{f(n,i_0)}, \dots, W_{f(n,i_{\ell-1})}$ all contain an element of the form $\langle 0, n', m \rangle$, $n' \neq n$.
3. $\langle 1, i, x \rangle$ for each $x \in \omega$ if $W_{f(n,i)}$ contains an element of the form $\langle 0, n', m \rangle$, $n' \neq n$.

Think of the elements $\langle 0, n, m \rangle$ as coding into A_n the value of n . Thus the indices i_0, i_1, \dots are an enumeration of the sets $W_{f(n, \cdot)}$ that might be equal to A_n . The set A_n will diagonalize against the n th computably enumerable family of sets $\{W_{f(n, i)} : i \in \omega\}$ using the elements $\langle 1, \cdot, \cdot \rangle$, and in particular it will diagonalize against $W_{f(n, i)}$ using the elements $\langle 1, i, \cdot \rangle$. If $W_{f(n, i)}$ contains an element $\langle 0, n', m \rangle$ for some m , and $n' \neq n$, then it cannot be A_n ; so as in (3) we do not need to diagonalize against it. In (2), we find the least $W_{f(n, i)}$ that might be A_n , and we use Lemma 4.7 to diagonalize against it. The function g strips off this first two entries of the elements $\langle 1, i, \cdot \rangle$ that we use to diagonalize. It is important to note that everything put into A_n for (2) is also put into A_n for (3); this is important as it is possible for the conditions for both (2) and (3) to be true.

It is clear that D can enumerate \mathcal{F} . We will show that the n th computably enumerable family is different from \mathcal{F} . Suppose that the n th computably enumerable family were equal to \mathcal{F} . Then there is some least ℓ such that $W_{f(n, i_\ell)} =^* A_n$ and such that $W_{f(n, i_\ell)}$ does not contain an element of the form $\langle 0, n', m \rangle$ for $n' \neq n$. For each $k < \ell$ such that $W_{f(n, i_k)} \neq^* A_n$, $W_{f(n, i_k)} =^* A_{n'}$ for some $n' \neq n$, and so it contains an element of the form $\langle 0, n', m \rangle$. Thus for every $k < \ell$, $W_{f(n, i_k)}$ contains an element of the form $\langle 0, n', m \rangle$. Then

$$W_{g(n, i_\ell)} = \{x : \langle 1, i_\ell, x \rangle \in W_{f(n, i_\ell)}\} =^* \{x : \langle 1, i_\ell, x \rangle \in A_n\} = U_{g(n, i_\ell)}^D.$$

But $W_{g(n, i)} \neq^* U_{g(n, i)}^D$, so this is a contradiction. Thus we have shown that \mathcal{F} cannot be computably enumerated.

We must also check that each A_n is c.e. This is because we can non-uniformly know the least index i_ℓ , if it exists, such that $W_{f(n, i_0)}, \dots, W_{f(n, i_{\ell-1})}$ all contain elements of the form $\langle 0, n', m \rangle$ for $n' \neq n$. For each $k' < \ell$, any element enumerated by (2) is also enumerated by (3), and so A_n consists of:

1. $\langle 0, n, m \rangle$ for each $m \in \omega$;
2. $\langle 1, i_\ell, x \rangle$ for $x \in U_{g(n, i_\ell)}^D$;
3. $\langle 1, i, x \rangle$ for each $x \in \omega$ if $W_{f(n, i)}$ contains an element of the form $\langle 0, n', m \rangle$, $n' \neq n$.

This is c.e. as $U_{g(n, i_\ell)}^D$ is c.e. □

We now return to the proof of Lemma 4.7.

Proof of Lemma 4.7. Uniformly in e , we need to define $U_e^D \neq^* W_e$. One can think of the construction as trying to define

$$U_e^D = \{\langle k, D(k), n \rangle : k, n \in \omega\}$$

where $D(k) = 0$ or $D(k) = 1$. This can easily be enumerated by D , and it is not equal to W_e because any enumeration of U_e^D can compute D . But U_e^D is not c.e. We need to make U_e^D c.e., but of course it cannot be c.e. uniformly in e .

It will be helpful to think of the c.e. set W_e trying to copy the set $\{\langle k, D(k), n \rangle : k, n \in \omega\}$ using the computable approximation D_s , while we as the builders of U_e^D are trying to come up with infinitely many differences between W_e and U_e^D . The two most extreme strategies that W_e might take (neither of which will work of course) can be thought of as the greedy strategy

and the cautious strategy. The greedy strategy computes, at each stage s , $D_s(k)$, and puts $\langle k, D_s(k), t \rangle$, $t < s$, into W . Then W_e will include all of the elements $\langle k, D(k), n \rangle$, but it will also contain some elements $\langle k, 1 - D(k), n \rangle$. The cautious strategy never enumerates any elements into W_e , because it can never be sure that $D_s(k)$ has stabilized.

Think of W_e as choosing a different one of these strategy for each k ; so for example it might be greedy for $k = 0$, cautious for $k = 1$, etc. (Of course W_e might take some other strategy, but in some sense the cautious and greedy strategies are prototypical and we can consider those other strategies later.)

If W_e chooses the cautious strategy for some least k , then we already have infinitely many differences between W_e and U_e^D using only the elements $\langle k, \cdot, \cdot \rangle$, because W_e contains none of these elements and U_e^D will contain all of the elements $\langle k, D(k), \cdot \rangle$. So we do not need to add to U_e^D any elements $\langle k', \cdot, \cdot \rangle$ for $k' > k$, and in fact we will keep all such elements out of U_e^D . Then, non-uniformly knowing the values of $D(0), \dots, D(k)$, we will be able to enumerate U_e^D .

Otherwise, suppose that W_e takes the greedy strategy for every k . Then we need to make sure that there are infinitely many elements $\langle k, i, n \rangle \in W_e - U_e^D$. In this case, we will have

$$U_e^D = \{\langle k, D(k), n \rangle : k, n \in \omega\} \cup V$$

where V is a c.e. set that also follows the greedy strategy for every k , but it will do so slower than W_e does. What we mean by this is that, for example, $\langle k, 0, 0 \rangle$ will not be enumerated into V until $\langle k, 0, 1 \rangle$ is enumerated into W_e , and only if $D_s(k)$ is still equal to 0, and $\langle k, 0, 1 \rangle$ will not be enumerated into V until $\langle k, 0, 2 \rangle$ is enumerated into W_e , and so on; thus if in fact $D(k) = 1$, then W_e will still have made a mistake by containing at least one element $\langle k, 0, n \rangle$ that is not in V . (Since V follows the greedy strategy, in fact it will contain $\{\langle k, D(k), n \rangle : k, n \in \omega\}$; and it will contain only finitely many elements $\langle k, 1 - D(k), n \rangle$.)

Of course W_e we have to be able to combine all of this, as well as defeating any other strategy W_e might take. But this will be the guiding idea behind our construction of U_e^D .

Let D_s be a Δ_2^0 approximation to D . For each k , define $U_{e,k}^D$ as follows:

1. Enumerate $\langle k, 0, n \rangle$ into $U_{e,k}^D$ if $D(k) = 0$, and enumerate $\langle k, 1, n \rangle$ into $U_{e,k}^D$ if $D(k) = 1$.
2. If $\langle k, 0, n \rangle$ is in $W_{e,s}$ and $D_s(k) = 0$, enumerate $\langle k, 0, 0 \rangle, \dots, \langle k, 0, n - 1 \rangle$ into $U_{e,k}^D$.
3. If $\langle k, 1, n \rangle$ is in $W_{e,s}$ and $D_s(k) = 1$, enumerate $\langle k, 1, 0 \rangle, \dots, \langle k, 1, n - 1 \rangle$ into $U_{e,k}^D$.

This process is not uniformly computable in k because of (1), which requires the oracle D ; but (2) and (3) enumerate elements computably. (2) and (3) are following the greedy strategy described above, but slower than W_e . If $D(k) = 0$, then $U_{e,k}^D$ contains all of the elements $\langle k, 0, n \rangle$ and only finitely many elements $\langle k, 1, n \rangle$; and if $D(k) = 1$, then $U_{e,k}^D$ contains all of the elements $\langle k, 1, n \rangle$ and only finitely many elements $\langle k, 0, n \rangle$.

Now we need to put the sets $U_{e,k}^D$ together into a single set U_e^D . Recall that if W_e follows the cautious strategy for k , then we do not want to put any elements $\langle k', \cdot, \cdot \rangle$ into U_e^D for $k' > k$; so we do not want to put $U_{e,k'}^D$ into U_e^D . Define $\text{agreement}(k, s)$ to be ∞ if $k = 0$, and otherwise it is the greatest $\ell \leq s$ such that for $k' < k$, if $D_s(k') = 0$ then $W_{e,s}$ contains

ℓ elements of the form $\langle k, 0, n \rangle$; and if $D_s(k') = 1$ then $W_{e,s}$ contains ℓ elements of the form $\langle k, 1, n \rangle$. Let

$$U_e^D = \bigcup_{k,s} U_{e,k}^D \upharpoonright_{\text{agreement}(k,s)-k}.$$

(If $k \geq \text{agreement}(k, s)$ then take $U_{e,k}^D \upharpoonright_{\text{agreement}(k,s)-k}$ to be empty.) Note that for a fixed s , $\text{agreement}(k, s)$ is decreasing in k , and that for each fixed k it will converge to a limit (which might be ∞) as $s \rightarrow \infty$. So if $\lim_{s \rightarrow \infty} \text{agreement}(k^*, s) < \infty$ for some k^* , then

$$\bigcup_{k \geq k^*, s} U_{e,k}^D \upharpoonright_{\text{agreement}(k,s)-k}$$

will be finite. One should think of the agreement function as measuring the extent to which W_e is following a greedy strategy by actually enumerating elements. The agreement function is computable.

Claim 1. U_e^D is c.e.

Proof. Let A_k be the c.e. set defined by (2) and (3), namely:

1. Whenever $\langle k, 0, n \rangle$ enters $W_{e,s}$ and $D_s(k) = 0$, enumerate $\langle k, 0, 0 \rangle, \dots, \langle k, 0, n-1 \rangle$ in A_k .
2. Whenever $\langle k, 1, n \rangle$ enters $W_{e,s}$ and $D_s(k) = 1$, enumerate $\langle k, 1, 0 \rangle, \dots, \langle k, 1, n-1 \rangle$ in A_k .

Let

$$A = \bigcup_{k,s} A_k \upharpoonright_{\text{agreement}(k,s)-k}.$$

One possibility is that $U_e^D = A$, so that U_e^D is c.e. since A is c.e. (Think of this as being when W_e follows a greedy strategy for every k .)

Now suppose otherwise. It is clear that $A \subseteq U_e^D$. So suppose that there is an element $\langle k, i, n \rangle \in U_e^D$ with $\langle k, i, n \rangle \notin A$; thus $\langle k, i, n \rangle \notin A_k$. We will show that U_e^D is c.e. in this case as well. Choose such a $\langle k^*, i, n \rangle$ with k^* minimal. It must be that $\langle k^*, i, n \rangle \in U_e^D$ due to (1), namely that $D(k^*) = i$. For sufficiently large stages s , $D_s(k^*) = i$. Since $\langle k^*, i, n \rangle \notin A_k$, there is no $\langle k^*, i, m \rangle \in W_e$ with $m > n$. (Think of this as meaning that W_e was following a cautious strategy for k^* .) So for every $k > k^*$, $\text{agreement}(k, s) \leq n$. Thus

$$U_e^D = \bigcup_{k,s} U_{e,k}^D \upharpoonright_{\text{agreement}(k,s)-k} =^* \bigcup_{k < k^*, s} U_{e,k}^D \upharpoonright_{\text{agreement}(k,s)-k}.$$

Each $U_{e,k}^D$ is c.e., but not uniformly over k ; one must know the value of $D(k)$. So U_e^D is a finite difference of a finite union of c.e. sets, hence c.e. \square

Claim 2. If D is non-computable, then $U_e^D \neq^* W_e$.

Proof. Suppose to the contrary that $U_e^D =^* W_e$. We argue by induction on k that for each k , $\lim_{s \rightarrow \infty} \text{agreement}(k, s) = \infty$. For $k = 0$, $\text{agreement}(k, s) = \infty$ by definition. Now suppose that $\lim_{s \rightarrow \infty} \text{agreement}(k', s) = \infty$ for $k' < k$. Then for $k' < k$, $U_{e,k'}^D \subseteq U_e^D$, and so $U_{e,k'}^D \subseteq^* W_e$. If $D(k') = 0$, then $U_{e,k'}^D$ contains $\langle k', 0, n \rangle$ for each n , and so W_e contains infinitely many such elements; and from some stage $s_{k'}$ on, $D_s(k') = 0$. Similarly, if $D(k') = 1$, then $U_{e,k'}^D$ contains

$\langle k', 1, n \rangle$ for each n , and so W_e contains infinitely many such elements; and from some stage $s_{k'}$ on, $D_s(k') = 1$. It follows that $\lim_{s \rightarrow \infty} \text{agreement}(k, s) = \infty$. So

$$U_e^D = \bigcup_k U_{e,k}^D.$$

Note that every element in $U_{e,k}^D$ has the form $\langle k, i, n \rangle$, so it is easy to break U_e^D up into the union $U_{e,k}^D$.

Now for each k , wait for some element $\langle k, i, n \rangle$ to enter W_e . Define $C(k) = i$ where $\langle k, i, n \rangle$ is the first such element to enter W_e . Note that C is computable and defined on every input. We will argue that $C =^* D$, showing that D is computable. In particular, we will argue that if $C(k) \neq D(k)$, then there is an element $\langle k, i, n \rangle \in W_e - U_{e,k}^D$. Since $W_e =^* U_e^D$ there can only be finitely many such elements.

If $C(k) \neq D(k)$, then there is $\langle k, i, n \rangle \in W_e$ with $i \neq D(k)$. Now as $i \neq D(k)$, $\langle k, i, n \rangle$ cannot have been enumerated into $U_{e,k}^D$ by (1). So it must have been enumerated by (2) or (3). There are at most finitely many stages s at which $D_s(k) = i$, and at every such stage s there is some element $\langle k, i, m \rangle$ in $W_{e,s}$ (namely, the largest such element) that is not enumerated into $U_{e,k}^D$. So there is some such element $\langle k, i, m \rangle \in W_e - U_{e,k}^D$. \square

These claims complete the proof of the lemma. \square

References

- [AK00] C. J. Ash and J. Knight. *Computable structures and the hyperarithmetical hierarchy*, volume 144 of *Studies in Logic and the Foundations of Mathematics*. North-Holland Publishing Co., Amsterdam, 2000.
- [FK15] Marat Faizrahmanov and Iskander Kalimullin. Limitwise monotonic sets of reals. *MLQ Math. Log. Q.*, 61(3):224–229, 2015.
- [Fri58] Richard M. Friedberg. Three theorems on recursive enumeration. I. Decomposition. II. Maximal set. III. Enumeration without duplication. *J. Symb. Logic*, 23:309–316, 1958.
- [GMS13] Noam Greenberg, Antonio Montalbán, and Theodore A. Slaman. Relative to any non-hyperarithmetical set. *J. Math. Log.*, 13(1):1250007, 26, 2013.
- [Gon80a] S. S. Gončarov. Computable univalent numerations. *Algebra i Logika*, 19(5):507–551, 617, 1980.
- [Gon80b] S. S. Gončarov. The problem of the number of nonautoequivalent constructivizations. *Algebra i Logika*, 19(6):621–639, 745, 1980.
- [Kho86] B. M. Khossainov. Strongly effective unars and nonautoequivalent constructivizations. In *Some problems in differential equations and discrete mathematics (Russian)*, pages 33–44. Novosibirsk. Gos. Univ., Novosibirsk, 1986.

- [Lac65] A. H. Lachlan. On recursive enumeration without repetition. *Z. Math. Logik Grundlagen Math.*, 11:209–220, 1965.
- [Lac67] A. H. Lachlan. A correction to: “On recursive enumeration without repetition”. *Z. Math. Logik Grundlagen Math.*, 13:99–100, 1967.
- [Mal65] A. I. Mal’cev. Positive and negative enumerations. *Dokl. Akad. Nauk SSSR*, 160:278–280, 1965.
- [Sla98] Theodore A. Slaman. Relative to any nonrecursive set. *Proc. Amer. Math. Soc.*, 126(7):2117–2122, 1998.
- [Weh98] Stephan Wehner. Enumerations, countable structures and Turing degrees. *Proc. Amer. Math. Soc.*, 126(7):2131–2139, 1998.