

MAXIMAL TOWERS AND ULTRAFILTER BASES IN COMPUTABILITY THEORY

STEFFEN LEMPP, JOSEPH S. MILLER,
ANDRÉ NIES, AND MARIYA I. SOSKOVA

ABSTRACT. The tower number \mathfrak{t} and the ultrafilter number \mathfrak{u} are cardinal characteristics from set theory, They are based on combinatorial properties of classes of subsets of ω and the almost inclusion relation \subseteq^* between such subsets. We consider analogs of these cardinal characteristics in computability theory.

We say that a sequence $\langle G_n \rangle_{n \in \omega}$ of computable sets is a *tower* if $G_0 = \omega$, $G_{n+1} \subseteq^* G_n$, and $G_n \setminus G_{n+1}$ is infinite for each n . A tower is *maximal* if there is no infinite computable set contained in all G_n . A tower $\langle G_n \rangle_{n \in \omega}$ is an *ultrafilter base* if for each computable R , there is n such that $G_n \subseteq^* R$ or $G_n \subseteq^* \bar{R}$; this property implies maximality of the tower. A sequence $\langle G_n \rangle_{n \in \omega}$ of sets can be encoded as the “columns” of a set $G \subseteq \omega$. Our analogs of \mathfrak{t} and \mathfrak{u} are the mass problems of sets encoding maximal towers, and of sets encoding towers that are ultrafilter bases, respectively. The relative position of a cardinal characteristic broadly corresponds to the relative computational complexity of the mass problem. We use Medvedev reducibility to formalize relative computational complexity, and thus to compare such mass problems to known ones.

We show that the mass problem of ultrafilter bases is equivalent to the mass problem of computing a function that dominates all computable functions, and hence, by Martin’s characterization, it captures highness. On the other hand, the mass problem for maximal towers is below the mass problem of computing a non-low set. We also show that some, but not all, noncomputable low sets compute maximal towers: Every noncomputable (low) c.e. set computes a maximal tower but no 1-generic Δ_2^0 -set does so.

We finally consider the mass problems of maximal almost disjoint, and of maximal independent families. We show that they are Medvedev equivalent to maximal towers, and to ultrafilter bases, respectively.

CONTENTS

1. Introduction	2
2. Basics of the mass problems $\mathcal{T}_{\mathbb{B}}$	5
3. Complexity of \mathcal{T} and of \mathcal{U}	7

Date: May 29, 2021.

2020 Mathematics Subject Classification. Primary 03D30.

Key words and phrases. sequences of computable sets, mass problems, Medvedev reducibility, cardinal characteristics, highness, ultrafilters.

Lempp was partially supported by Simons Collaboration Grant for Mathematicians #626304. Miller was partially supported by grant #358043 from the Simons Foundation. Nies was partially supported by the Marsden fund of New Zealand, grant 19-UOA-346. Soskova was partially supported by NSF Grant DMS-1762648. The authors thank Jörg Brendle and Noam Greenberg for helpful discussions during a workshop at the Casa Matemática Oaxaca in August 2019, where this research received its initial impetus.

4. Maximal independent families in computability	11
5. The co-computably enumerable case	13
6. Ultrafilter bases for other Boolean algebras	17
References	19

1. INTRODUCTION

Cardinal characteristics measure how far the set-theoretic universe deviates from satisfying the continuum hypothesis. They are natural cardinals greater than \aleph_0 and at most 2^{\aleph_0} . For instance, the *bounding number* \mathfrak{b} is the least size of a collection of functions $f: \omega \rightarrow \omega$ such that no single function dominates the entire collection.¹ Related is the *dominating number* \mathfrak{d} , the least size of a collection of functions $f: \omega \rightarrow \omega$ such that every function is dominated by some function in the collection. Here, for functions $f, g: \omega \rightarrow \omega$, we say that g *dominates* f if $g(n) \geq f(n)$ for sufficiently large n . An important program in set theory is to prove less than or equal-relations between characteristics in ZFC, and to separate them in suitable forcing extensions.

Analogues of cardinal characteristics in computability theory were first studied by Rupperecht [14, 15] and further investigated by Brendle, Brooke-Taylor, Ng, and Nies [2]. An article by Greenberg, Kuyper, and Turetsky [7], in part based on Rupperecht's work, provides a systematic approach to the two connected settings of set theory and computability, at least for certain types of cardinal characteristics. The relevant characteristics are given by binary relations, such as the domination relation \leq^* between functions; their computability-theoretic analogs are ordered by reducibilities that measure relative computability. A well understood example of this is how the relation \leq^* gives rise to the bounding number $\mathfrak{b}(\leq^*)$ and the dominating number $\mathfrak{d}(\leq^*)$, and their analogs in computability, which are highness and having hyperimmune degree. A general reference in set theory is the survey paper by Blass [1]. The recent brief survey by Soukup [18] contains a diagram displaying the ZFC inequalities between the most important characteristics in this setting, along with $\mathfrak{b}(\leq^*)$ and $\mathfrak{d}(\leq^*)$.

In this paper, we consider cardinal characteristics that do not fit into the framework of Rupperecht, and Greenberg et al. [7]. In particular, we initiate the study of the computability-theoretic analogs of the ultrafilter, tower, and independence numbers. These characteristics are defined in the setting of subsets of ω up to almost inclusion \subseteq^* ; we give definitions below.

The *ultrafilter number* \mathfrak{u} is the least size of a subset of $[\omega]^\omega$ with upward closure a nonprincipal ultrafilter on ω . We note that one cannot in general require here that the subset is linearly ordered by \subseteq^* . Recall that an ultrafilter F on ω is a P -point if for each partition $\langle C_n \rangle$ of ω such that $C_n \notin F$ for each n , there is $A \in F$ such that $C_n \cap A$ is finite for each n . An ultrafilter with a linear base is a P -point. Shelah (see Wimmers [19]) has shown that the non-existence of P -points is consistent with ZFC, so it is consistent that a version of \mathfrak{u} relying on linear bases would be undefined.

The *tower number* \mathfrak{t} is the minimum size of a subset of $[\omega]^\omega$ that is linearly ordered by \subseteq^* and cannot be extended by adding a new element below all given

¹This is less commonly, but perhaps more sensibly, called the *unbounding number*.

elements. To define the *pseudointersection number* \mathfrak{p} , the requirement in the definition of towers that the sets in the class be linearly ordered under \subseteq^* is weakened to requiring that every finite subset of the class has an infinite intersection. So, trivially, $\mathfrak{p} \leq \mathfrak{t}$. In celebrated work, Malliaris and Shelah [12] showed (in ZFC) that $\mathfrak{p} = \mathfrak{t}$ (see also [18]). It is not hard to see that ZFC proves $\mathfrak{t} \leq \mathfrak{u}$. It is consistent that $\mathfrak{t} < \mathfrak{u}$ (see [1] for both statements).

A class \mathcal{C} of subsets of ω is *independent* if any intersection of finitely many sets in \mathcal{C} or their complements is infinite. The *independence number* \mathfrak{i} is the least cardinal of a maximal independent family. There has been much work recently on \mathfrak{i} in set theory, in particular, the descriptive complexity of maximal independent families, such as in Brendle, Fischer, and Khomskii [3].

1.1. Comparing the complexity of the analogs in computability. The main setting for our analogy is given by the Boolean algebra of computable sets modulo finite differences. We consider maximal towers, the closely related maximal almost disjoint sets, and thereafter ultrafilter bases and maximal independent sets. As already demonstrated in the above-mentioned papers [14, 15, 2, 7], the relative position of a cardinal characteristic tends to correspond to the relative computational complexity of the associated class of objects.

Note that the usual formal definitions of computation relative to an oracle only directly apply to functions $f: \omega \rightarrow \omega$, and to subsets of ω (simply called sets from now on, and identified with their characteristic functions). The complexity of other objects is studied indirectly, via names that are functions on ω giving discrete representations of the object in question. A particular choice of names has to be made. For instance, real numbers can be named by rapidly converging Cauchy sequences of rational numbers.

The witnesses for cardinal characteristics are always uncountable. In contrast, in our setting, the analogous objects are countable. They will be considered as sequences of sets rather than unordered collections. For, a single set X can be used as a name for such a sequence of sets: Let $X^{[n]}$ denote the ‘‘column’’ $\{u: \langle u, n \rangle \in X\}$.² To every set X , we can associate a sequence $\langle X_n \rangle_{n \in \omega}$ in a canonical way by setting $X_n = X^{[n]}$. (When introducing terminology, we will sometimes ignore the difference between $\langle X_n \rangle_{n \in \omega}$ and X .) An alternate viewpoint is that a set X is a name for the unordered collection of sets in its coded sequence. Although such a name includes more information than is in the unordered family, this information is suppressed when we quantify over all names; our results can be read in this context.

With this naming system, one can now use sequences as oracles in computations. We view the combinatorial classes of sequences as mass problems. To measure their relative complexity, we compare them via *Medvedev reducibility* \leq_s : Let \mathcal{C} and \mathcal{D} be sets of functions on ω , also known as *mass problems*. We say that \mathcal{C} is Medvedev reducible to \mathcal{D} and write $\mathcal{C} \leq_s \mathcal{D}$ if there is a Turing functional Θ such that $\Theta^g \in \mathcal{C}$ for each $g \in \mathcal{D}$. Less formally, we say that functions in \mathcal{D} uniformly compute functions in \mathcal{C} . We will also refer to the weaker *Muchnik reducibility*: $\mathcal{C} \leq_w \mathcal{D}$ if each function in \mathcal{D} computes a function in \mathcal{C} .

With subsequent research in mind, we will set up our framework to apply to general countable Boolean algebras rather than merely the Boolean algebra of the

²For definiteness, we employ the usual computable Cantor pairing function $\langle x, n \rangle$. Note that $\langle x, n \rangle \geq x, n$. This property is useful in simplifying notation in some of the constructions below.

computable sets. Throughout, we fix a countable Boolean algebra \mathbb{B} of subsets of ω closed under finite differences. Our basic objects will be sequences of sets in \mathbb{B} . We will obtain meaningful results already when we fix a countable Turing ideal \mathcal{I} and let \mathbb{B} be the sets with degree in \mathcal{I} . While we mainly study the case when \mathbb{B} consists of the computable sets, in Section 6, we briefly consider two other cases: the K -trivial sets and the primitive recursive sets.

1.2. The mass problem $\mathcal{T}_{\mathbb{B}}$ of maximal towers.

Definition 1.1. We say that a sequence $\langle G_n \rangle_{n \in \omega}$ of sets in \mathbb{B} is a \mathbb{B} -tower if $G_0 = \omega$, $G_{n+1} \subseteq^* G_n$, and $G_n \setminus G_{n+1}$ is infinite for each n . If \mathbb{B} consists of the computable sets, we use the term *tower of computable sets*.

Definition 1.2. We say that a function p is *associated with* a tower G if p is increasing and $p(n) \in \bigcap_{i \leq n} G_i$ for each n .

The following fact is elementary.

Fact 1.3. *A tower G uniformly computes a function p associated with it.*

Proof. Let Φ be the Turing functional such that $\Phi^G(0) = \min(G_0)$, and $\Phi^G(n+1)$ is the least number in $\bigcap_{i \leq n+1} G_i$ greater than $\Phi^G(n)$. This Φ establishes the required uniform reduction. \square

Definition 1.4. Given a countable Boolean algebra \mathbb{B} of sets, the mass problem $\mathcal{T}_{\mathbb{B}}$ is the class of sets G such that $\langle G_n \rangle_{n \in \omega}$ is a \mathbb{B} -tower that is *maximal*, i.e., such that for each infinite set $R \in \mathbb{B}$, there is n such that $R \setminus G_n$ is infinite.

Clearly, being maximal implies that no associated function is computable. In particular, a maximal tower is never computable. (Note that our notion of maximality only requires that the tower cannot be extended from below, in keeping with our set-theoretic analogy.)

1.3. The mass problem $\mathcal{U}_{\mathbb{B}}$ of ultrafilter bases. We now define the mass problem $\mathcal{U}_{\mathbb{B}}$ corresponding to the ultrafilter number. Since all filters of our Boolean algebras are countable, any base will compute a linearly ordered base by taking finite intersection. So we can restrict ourselves to linearly ordered bases. This is not always possible in the set theory setting, as discussed in the introduction.

Definition 1.5. Given a countable Boolean algebra \mathbb{B} of sets, let $\mathcal{U}_{\mathbb{B}}$ be the class of sets F such that F is a \mathbb{B} -tower as in Definition 1.1 and for each set $R \in \mathbb{B}$, there is n such that $F_n \subseteq^* \overline{R}$ or $F_n \subseteq^* R$. We will call a set F in $\mathcal{U}_{\mathbb{B}}$ a \mathbb{B} -ultrafilter base.

Each ultrafilter base is a maximal tower. In the cardinal setting, one has $\mathfrak{t} \leq \mathfrak{u}$. Correspondingly, since $\mathcal{U}_{\mathbb{B}} \subseteq \mathcal{T}_{\mathbb{B}}$, we trivially have $\mathcal{T}_{\mathbb{B}} \leq_s \mathcal{U}_{\mathbb{B}}$ via the identity reduction. The following indicates that for many natural Boolean algebras, ultrafilter bases necessarily have computational strength.

Proposition 1.6. *Suppose that the degrees of sets in \mathbb{B} form a Turing ideal \mathcal{K} . Then for each \mathbb{B} -ultrafilter base F and associated function p in the sense of Definition 1.2, the function p is not dominated by a function with Turing degree in \mathcal{K} .*

Proof. Assume that there is a function $f \geq p$ in \mathcal{K} . The conditions $n_0 = 0$ and $n_{k+1} = f(n_k) + 1$ define a sequence that is computable from some oracle in \mathcal{K} , and

for every k we have that $[n_k, n_{k+1})$ contains an element of $\bigcap_{i \leq k} F_i$. So the set

$$E = \bigcup_{i \in \omega} [n_{2i}, n_{2i+1})$$

is in \mathcal{K} , and clearly $F_n \not\subseteq^* E$ and $F_n \not\subseteq^* \bar{E}$ for each n . Therefore, F is not a \mathbb{B} -ultrafilter base. \square

1.4. The Boolean algebra of computable sets. We finish the introduction by summarizing our results in the case that \mathbb{B} is the Boolean algebra of all computable sets. By Theorem 3.1, every non-low set computes a set in $\mathcal{T}_{\mathbb{B}}$, and this is uniform. This is not a characterization, however, because by Corollary 5.3, every noncomputable c.e. set computes a maximal tower. On the other hand, we know that there are noncomputable (necessarily low) sets that do not compute maximal towers; in particular, no 1-generic Δ_2^0 -set does so. This is because 1-generic Δ_2^0 -sets are index guessable by Theorem 3.4, and by Proposition 2.4, no index guessable set can compute a maximal tower. Here, an oracle G is *index guessable* if \emptyset' can find a computable index for φ_e^G uniformly in e , provided that φ_e^G is computable. We do not know whether index guessability characterizes the oracles that are unable to compute a maximal tower. It seems unlikely; index guessability appears to be stronger than necessary.

As already mentioned, in the setting of cardinal characteristics, $\mathfrak{t} < \mathfrak{u}$ is consistent with ZFC. Since non-low oracles can be computably dominated, it follows from Proposition 1.6 that there is a member of $\mathcal{T}_{\mathbb{B}}$ that does not compute any member of $\mathcal{U}_{\mathbb{B}}$. In other words, $\mathcal{U}_{\mathbb{B}} \not\leq_w \mathcal{T}_{\mathbb{B}}$ in the case that \mathbb{B} consists of the computable sets.

The separation above only uses the fact that members of $\mathcal{U}_{\mathbb{B}}$ are not computably dominated; in fact, they are *high*. As we show in Theorems 3.6 and 3.8, $\mathcal{U}_{\mathbb{B}}$ is Medvedev equivalent to the mass problem of dominating functions. In Section 4, we prove that the mass problem $\mathcal{I}_{\mathbb{B}}$ of maximal independent families is also Medvedev equivalent to the mass problem of dominating functions. Thus, in the case that \mathbb{B} is the Boolean algebra of computable sets, we have $\mathcal{U}_{\mathbb{B}} \equiv_s \mathcal{I}_{\mathbb{B}}$. Interestingly, we do not have a direct proof. Contrast this with the equivalence of $\mathcal{T}_{\mathbb{B}}$ and $\mathcal{A}_{\mathbb{B}}$, the mass problem of maximal almost disjoint families; this equivalence is direct and holds for an arbitrary Boolean algebra, as we will see presently.

2. BASICS OF THE MASS PROBLEMS $\mathcal{T}_{\mathbb{B}}$

2.1. The equivalent mass problems $\mathcal{T}_{\mathbb{B}}$ and $\mathcal{A}_{\mathbb{B}}$. Recall that in set theory, the almost disjointness number \mathfrak{a} is the least possible size of a maximal almost disjoint (MAD) family of subsets of ω . In our analogous setting, we call a sequence $\langle F_n \rangle_{n \in \omega}$ of sets in \mathbb{B} *almost disjoint* (AD) if each F_n is infinite and $F_n \cap F_k$ is finite for distinct n and k .

Definition 2.1. In the context of a Boolean algebra \mathbb{B} of sets, the mass problem $\mathcal{A}_{\mathbb{B}}$ is the class of sets F such that $\langle F_n \rangle_{n \in \omega}$ is a *maximal almost disjoint* (MAD) family in \mathbb{B} . Namely, the sequence is AD, and for each infinite set $R \in \mathbb{B}$, there is n such that $R \cap F_n$ is infinite.

Fact 2.2. $\mathcal{A}_{\mathbb{B}} \leq_s \mathcal{T}_{\mathbb{B}} \leq_s \mathcal{A}_{\mathbb{B}}$.

Proof. We suppress the subscript \mathbb{B} . To check that $\mathcal{A} \leq_s \mathcal{T}$, given a set G , let $\text{Diff}(G)$ be the set F such that $F_n = G_n \setminus G_{n+1}$ for each n . Clearly, the operator

Diff can be seen as a Turing functional. If G is a maximal \mathbb{B} -tower, then $F = \text{Diff}(G)$ is MAD. For, if $R \in \mathbb{B}$ is infinite, then $R \setminus G_n$ is infinite for some n , and hence $R \cap F_i$ is infinite for some $i < n$.

For $\mathcal{T} \leq_s \mathcal{A}$, given a set F , let $G = \text{Cp}(F)$ be the set such that

$$x \in G_n \leftrightarrow \forall i < n [x \notin F_i].$$

Again, Cp is a Turing functional. If F is an almost disjoint family of sets from \mathbb{B} , then G is a \mathbb{B} -tower, and if F is MAD, then G is a maximal tower. \square

Recall that a maximal tower is not computable. Hence no MAD family is computable. (This corresponds to the cardinal characteristics being uncountable.)

2.2. Descriptive complexity and index complexity for maximal towers.

For the rest of this section, as well as the subsequent three sections, we will mainly be interested in the case that \mathbb{B} is the Boolean algebra of all computable sets. We will omit the parameter \mathbb{B} when we name the mass problems. In the final section, we will consider other Boolean algebras.

Besides looking at the relative complexity of mass problems such as \mathcal{T} and \mathcal{U} , one can also look at the individual complexity of their members (as sets encoding sequences). Recall that a characteristic index for a set M is a number e such that $\chi_M = \varphi_e$. The following two questions arise:

- (1) How low in the arithmetical hierarchy can the set be located?
- (2) How hard is it to find characteristic indices for the sequence members?

Arithmetical complexity.

Fact 2.3. *No maximal tower G is c.e., and no MAD set is co-c.e.*

Proof. For the first statement, note that otherwise there is a computable function p associated with G . The range of p would extend the tower G .

For the second statement, note that the reduction Cp, introduced above to show that $\mathcal{T} \leq_s \mathcal{A}$, turns a co-c.e. set F into a c.e. set G . \square

We will return to Question (1) in Section 5, where we show that c.e. MAD sets exist in every nonzero c.e. degree, and that some ultrafilter base is co-c.e.

Complexity of finding characteristic indices for the sequence members. In several constructions of towers $\langle G_n \rangle_{n \in \omega}$ below, such as in Corollary 5.3 and Theorem 5.4, the oracle \emptyset'' is able to compute, given n , a characteristic index for G_n . The oracle \emptyset' does not suffice by the following result.

Proposition 2.4. *Suppose that G is a maximal tower. The oracle \emptyset' is not able to compute, from input n , a characteristic index for G_n .*

Proof. Assume the contrary. Then there is a computable function f such that $\varphi_{\lim_s f(n,s)}$ is the characteristic function of G_n . Let \widehat{G} be defined as follows. Given n and x , compute the least $s > x$ such that $\varphi_{f(n,s),s}(x) \downarrow$. If the output is not 0, put x into \widehat{G}_n . Clearly \widehat{G} is computable. Since $G_n =^* \widehat{G}_n$ for each n , \widehat{G} is a maximal tower, contrary to Fact 2.3, or to the earlier observation that maximal towers cannot be computable. \square

3. COMPLEXITY OF \mathcal{T} AND OF \mathcal{U}

In this section, we compare our two principal mass problems, maximal towers and ultrafilter bases, to well-known benchmark mass problems: non-lowness and highness. We also define index guessability. No index guessable oracle computes a maximal tower. We show that every 1-generic Δ_2^0 -set is index guessable.

As we said above, we restrict ourselves to the case that \mathbb{B} is the Boolean algebra of computable sets.

3.1. Maximal towers, non-lowness, and index guessability. We now show that each non-low oracle computes a set in \mathcal{T} . The result is uniform in the sense of mass problems. Let NonLow denote the class of oracles Z such that $Z' \not\leq_T \emptyset'$.

Theorem 3.1. $\mathcal{T} \leq_s \text{NonLow}$.

Proof. In the following, x, y , and z denote binary strings; we identify such a string x with the number whose binary expansion is $1x$. For example, the string 000 is identified with 8 , the number with binary representation 1000 . Define a Turing functional Θ for the Medvedev reduction as follows: Set $\Theta^Z = G$, where for each n ,

$$G_n = \{x: n \leq s := |x| \wedge Z'_s \upharpoonright n = x \upharpoonright n\}.$$

Here Z' denotes the jump of Z , which is computably enumerated relative to Z in a standard way. Note that, for each n , for sufficiently large s , the string $Z'_s \upharpoonright n$ settles. So it is clear that for each n , we have $G_{n+1} \subseteq^* G_n$ and $G_n \setminus G_{n+1}$ is infinite. Also G_n is computable.

Suppose now that R is an infinite set such that $R \subseteq^* G_n$ for each n . Then for each k ,

$$Z'(k) = \lim_{x \in G_k, |x| > k} x(k) = \lim_{x \in R, |x| > k} x(k),$$

and hence $Z' \leq_T R'$. So if $Z \in \text{NonLow}$, then R cannot be computable, and hence $\Theta^Z \in \mathcal{T}$. \square

Remark 3.2. The proof above yields a more general result. Suppose that \mathcal{K} is a countable Turing ideal and \mathbb{B} is the Boolean algebra of sets with degree in \mathcal{K} . Then $\mathcal{T}_{\mathbb{B}} \leq_s \text{NonLow}_{\mathcal{K}}$, where $\text{NonLow}_{\mathcal{K}} := \{Z: \forall R \in \mathcal{K} [Z' \not\leq_T R']\}$.

We next introduce a property of oracles that we call *index guessability*; it implies that an oracle does not compute a maximal tower. As usual, let $\langle \Phi_e \rangle_{e \in \omega}$ be an effective list of the Turing functionals with one input, and write φ_e for Φ_e^\emptyset . Note that if L is a Δ_2^0 -oracle, then \emptyset'' can compute from e a characteristic index for Φ_e^L in case that the function Φ_e^L is computable. To be index guessable means that \emptyset' suffices.

Definition 3.3. We call an oracle L *index guessable* if \emptyset' can compute from e an index for Φ_e^L whenever Φ_e^L is a computable function. In other words, there is a functional Γ such that

$$\Phi_e^L \text{ is computable} \Rightarrow \Phi_e^L = \varphi_{\Gamma(\emptyset'; e)}.$$

No assumption is made on the convergence of $\Gamma(\emptyset'; e)$ in case Φ_e^L is not a computable function.

Clearly, being index guessable is closed downward under \leq_T . A total function is computable if and only if its graph is computable, in a uniform way. So for index

guessability of L , it suffices that there is a Turing functional Γ such that $\Gamma(\emptyset'; e)$ provides an index for Φ_e^L in case it is a computable $\{0, 1\}$ -valued function.

Every index guessable oracle D is low. To see this, for $i \in \omega$, let $B_i = \{t : i \in D'_t\}$. If $i \in D'$ then B_i is cofinite, otherwise $B_i = \emptyset$. There is a computable function g such that $\Phi_{g(i)}^D$ is the characteristic function of B_i . To show that $D' \leq_T \emptyset'$, on input i , let \emptyset' compute a computable index $r(i)$ for B_i . Now use \emptyset' again to determine $\lim_k \varphi_{r(i)}(k)$, which equals $D'(i)$.

By Proposition 2.4, an index guessable oracle D does not compute a maximal tower. The following provides examples of such oracles.

Theorem 3.4. *If L is Δ_2^0 and 1-generic, then L is index guessable.*

Proof. Suppose that $F = \Phi_e^L$ and F is a computable set. Let S_e be the c.e. set of strings σ above which there is a Φ_e -splitting in the sense that

$$S_e = \{\sigma : (\exists p)(\exists \tau_1 \succ \sigma)(\exists \tau_2 \succ \sigma) \Phi_e^{\tau_1}(p) \neq \Phi_e^{\tau_2}(p)\}.$$

Suppose that S_e is dense along L . Then we claim that the set

$$C_e = \{\tau : (\exists p) \Phi_e^\tau(p) \neq F(p)\}$$

is also dense along L , i.e., for every k , there is some $\tau \succeq L \upharpoonright k$ such that $\tau \in C_e$. Indeed, let $\sigma \succeq L \upharpoonright k$ be a member of S_e and let p , τ_1 , and τ_2 witness this. Let τ_i for $i = 1$ or 2 be such that $\Phi_e^{\tau_i}(p) \neq F(p)$. Then $\tau_i \succeq L \upharpoonright k$ is in C_e . The set C_e is c.e. and hence L meets C_e , contradicting our assumption that $F = \Phi_e^L$.

It follows that S_e is not dense along L . In other words, there is some least k_e such that there is no splitting of Φ_e above $L \upharpoonright k_e$. On input e , the oracle \emptyset' can compute k_e and $L \upharpoonright k_e$. This allows \emptyset' to find an index for F , given by the following procedure: To compute $F(p)$, find the least $\tau \succeq L \upharpoonright k_e$ such that $\Phi_e^\tau(p) \downarrow$ (in $|\tau|$ many steps). Such a τ exists because $\Phi_e^L(p) \downarrow$. By our choice of k_e , it follows that $\Phi_e^\tau(p) = \Phi_e^L(p) = F(p)$. \square

We summarize the known implications:

$$1\text{-generic } \Delta_2^0 \Rightarrow \text{index guessable} \Rightarrow \text{computes no maximal tower} \Rightarrow \text{low}.$$

The last arrow does not reverse by Theorem 5.1 below; the others might. In particular, we ask whether any oracle that computes no maximal tower is index guessable. This would strengthen Theorem 3.1. Note that the following potential weakening of index guessability of L still implies that the oracle computes no maximal tower: For each $S \leq_T L$ such that each S_n is computable, there is binary computable function f such that $S_n = \varphi_{\lim_s f(n,s)}$ for each n .

Aside. We pause briefly to mention a potential connection of our topic to computational learning theory. One says that a class S of computable functions is *EX-learnable* if there is a total Turing machine M such that $\lim_s M(f \upharpoonright s)$ exists for each $f \in S$ and is an index for f . For an oracle A , one says that S is *EX[A]-learnable* if there is an oracle machine M that is total for each oracle and such that $\lim_s M^A(f \upharpoonright s)$ exists for each $f \in S$ and is an index for f . One calls an oracle A *EX-trivial* if $\text{EX} = \text{EX}[A]$. Slaman and Solovay [16] showed that A is EX-trivial if and only if A is Δ_2^0 and has 1-generic degree. This used an earlier result of Haight that the Turing degrees of the 1-generic Δ_2^0 -sets are closed downward.

3.2. Ultrafilter bases and highness. Let $\text{Tot} = \{e: \varphi_e \text{ is total}\}$. Let DomFcn denote the mass problem of functions h that dominate every computable function and also satisfy $h(s) \geq s$ for all s . Note that a set F is high if and only if $\text{Tot} \leq_T F'$. To represent highness by a mass problem in the Medvedev degrees, one can equivalently choose the set of functions dominating each computable function, or the set of approximations to Tot , i.e., the $\{0, 1\}$ -valued binary functions f such that $\lim_s f(e, s) = \text{Tot}(e)$. This follows from the next fact; we omit the standard proof.

Fact 3.5. *DomFcn is Medvedev equivalent to the mass problem of approximations to $\text{Tot} = \{e: \varphi_e \text{ is total}\}$.*

We show that exactly the high oracles compute ultrafilter bases, and that the reductions are uniform. By Fact 3.5, it suffices to show that $\mathcal{U} \equiv_s \text{DomFcn}$. We will obtain the two Medvedev reductions through separate theorems, with proofs that are unrelated.

Theorem 3.6. *Every ultrafilter base uniformly computes a dominating function. In other words, $\mathcal{U} \geq_s \text{DomFcn}$.*

Our proof is directly inspired by a proof of Jockusch [8, Theorem 1, (iv) \implies (i)], who showed that any family of sets containing exactly the computable sets must have high degree.

Lemma 3.7. *There is a uniformly computable sequence P_0, P_1, \dots of nonempty Π_1^0 -classes such that for every e ,*

- if φ_e is total, then P_e contains a single element, and
- if φ_e is not total, then P_e contains only bi-immune elements.

Proof. Note that each Martin-Löf (or even Kurtz) random set is bi-immune: For an infinite computable set R , the class of sets containing R is a Π_1^0 -null class and hence determines a Kurtz test. A similar fact holds for the class of sets disjoint from R .

For each s , let n_s be the largest number such that $\varphi_{e,s}$ converges on $[0, n_s)$. We build the Π_1^0 -class P_e in stages, where $P_{e,s}$ is the nonempty clopen set we have before stage s of the construction. Let $P_{e,0} = 2^\omega$.

Stage 0. Start constructing P_e as a nonempty Π_1^0 -class containing only Martin-Löf random elements.

Stage s . If $n_s = n_{s-1}$, continue the construction that is currently underway, which will produce a nonempty Π_1^0 -class of random elements.

On the other hand, if $n_s > n_{s-1}$, fix a string σ such that $[\sigma] \subseteq P_{e,s}$ and $|\sigma| > s$. Let $P_{e,s+1} = [\sigma]$. End the construction that we have been following and start a new construction for P_e , starting at stage $s+1$, as a nonempty Π_1^0 -subclass of $[\sigma]$ containing only Martin-Löf random elements.

It is clear that if φ_e is total, then P_e will be a singleton. Otherwise, there will be a final construction of a nonempty Π_1^0 -class of randoms which will run without further interruption. \square

Of course, when P_e is a singleton, its lone element must be computable.

Proof of Theorem 3.6. For any set C , let $S_C = \{X \in 2^\omega: C \subseteq X\}$. Note that if C is computable (or even merely c.e.), then S_C is a Π_1^0 -class. Let $Q_e = \{X: \bar{X} \in P_e\}$ be the Π_1^0 -class of complements of elements of P_e .

Now let F be an ultrafilter base. We have that

$$\begin{aligned} \varphi_e \text{ is total} &\iff (\exists i)(\exists n) [F_i \setminus [0, n] \text{ is a subset of some} \\ &\quad X \in P_e \text{ or its complement}] \\ &\iff (\exists i)(\exists n) [P_e \cap S_{F_i \setminus [0, n]} \neq \emptyset \text{ or } Q_e \cap S_{F_i \setminus [0, n]} \neq \emptyset]. \end{aligned}$$

Even though $S_{F_i \setminus [0, n]}$ is a Π_1^0 -class, we cannot hope to compute an index using F . However, $S_{F_i \setminus [0, n]}$ is a $\Pi_1^0[F]$ -class uniformly in i, n . Using the fact that the nonemptiness of a $\Pi_1^0[F]$ -class is a $\Pi_1^0[F]$ -property, we see that $\text{Tot} = \{e: \varphi_e \text{ is total}\}$ is $\Sigma_2^0[F]$. Note that the Σ_2^0 -index does not depend on F . Since Tot is also Π_2^0 , it is $\Delta_2^0[F]$ via a fixed pair of indices, and hence Turing reducible to F' via a fixed reduction. One direction of the usual proof of the (relativized) Limit Lemma now shows that we can uniformly compute an approximation to Tot from F . Hence, from F we can uniformly compute a dominating function by Fact 3.5. \square

Theorem 3.8. *Every dominating function uniformly computes an ultrafilter base. In other words, $\mathcal{U} \leq_s \text{DomFcn}$.*

Proof. Let $\langle \psi_e \rangle_{e \in \omega}$ be an effective listing of the $\{0, 1\}$ -valued partial computable functions defined on an initial segment of ω . Let $V_{e,k} = \{x: \psi_e(x) = k\}$ so that $\langle (V_{e,0}, V_{e,1}) \rangle$ is an effective listing that contains all pairs of computable sets and their complements.

Let $T = \{0, 1, 2\}^{<\omega}$. Uniformly in $\alpha \in T$, we will define a set S_α . We first explain the basic idea and then modify it to make it work. The basic idea is to start with $S_\emptyset = \omega$ and build $S_{\alpha \frown k} = S_\alpha \cap V_{e,k}$ for $k = 0, 1$ and $e = |\alpha|$, that is, we split S_α according to the listing above. We then consider the leftmost path g so that $S_{g \upharpoonright e}$ is infinite for each e . A dominating function h can eventually discover each initial segment of this path and use this to compute a set F such that $F_e =^* S_{g \upharpoonright e}$ for each e .

The problem is that both $S_\alpha \cap V_{e,0}$ and $S_\alpha \cap V_{e,1}$ could be finite (because e is not a proper index of a computable set). In this case we still need to make sure that $F_e \setminus F_{e+1}$ is infinite. So the rightmost option at level n is a set $S_{\alpha \frown 2} = \tilde{S}_\alpha$ that simply removes every other element from S_α . The sets $S_{\alpha \frown k}$ for $k \leq 1$ will be subsets of \tilde{S}_α .

We now provide the details. The set S_α is enumerated in increasing fashion, and possibly finite. So each S_α is computable, but not uniformly in α . All the sets and functions defined below can be interpreted at stages.

Let $S_{\emptyset, s} = [0, s)$. If we have defined (at stage s) the set $S_\alpha = \{r_0 < \dots < r_k\}$, let \tilde{S}_α contain the numbers of the form r_{2i} . Let $S_{\alpha \frown 2} = \tilde{S}_\alpha$. Let $S_{\alpha \frown k} = \tilde{S}_\alpha \cap V_{e,k}$ for $k = 0, 1$, $e = |\alpha|$. We define a uniform list of Turing functionals Γ_e so that the sequence $\langle \Gamma_e^h(t) \rangle_{t \in \omega}$ is nondecreasing and unbounded, for each e and each oracle function h such that $h(s) \geq s$ for each s . We will let $F_e = \{\Gamma_e^h(t): t \in \omega\}$.

Definition of Γ_e . Given an oracle function h , we will write a_s for $\Gamma_e^h(s)$. Let $a_0 = 0$. Suppose $s > 0$ and a_{s-1} has been defined. Check if there is $\alpha \in T$ of length e such that $|S_{\alpha, h(s)}| \geq s$. If there is no such α , let $a_s = a_{s-1}$. Otherwise, let α be leftmost such. If $\max S_{\alpha, h(s)} > a_{s-1}$, let $a_s = \max S_{\alpha, h(s)}$. Otherwise, again let $a_s = a_{s-1}$.

Note that the sequence $\{a_s\}_{s < \omega}$ is unbounded because for the rightmost string $\alpha \in T$ of length e , the set $S_{\alpha, t}$ consists of the numbers in $[0, t)$ divisible by 2^e . We

may combine the functionals Γ_e to obtain a functional Ψ such that $(\Psi^h)_e = F_e$ for each h with $h(s) \geq s$ for each s .

Claim 3.9. *If $h \in \text{DomFcn}$, then $F = \Psi^h \in \mathcal{U}$.*

To verify this, let $g \in 2^\omega$ denote the leftmost path in $\{0, 1, 2\}^\omega$ such that the set $S_{g \upharpoonright e}$ is infinite for every e . Note that g is an infinite path, because for every α , if the set S_α is infinite then so is $S_{\alpha \smallfrown 2}$.

Fix e and let $\alpha = g \upharpoonright e$. Let $p(s)$ be the least stage t such that $S_{\alpha, t}$ has at least s elements. Since h dominates the computable function p , we will eventually always pick α in the definition of $a_s = \Gamma_e^h(s)$. Hence $F_e =^* S_\alpha$. This implies that F_e is computable and $F_{e+1} \subseteq^* F_e$. Clearly, if S_α is infinite, then $S_\beta \subseteq_\infty S_\alpha$ for every $\beta \succ \alpha$. Thus $F_{e+1} \subseteq_\infty^* F_e$.³

Now let R be a computable set. Pick e such that $R = V_{e,0}$ and $\bar{R} = V_{e,1}$. If $g(e) = 0$, then $S_{g \upharpoonright e+1} \subseteq V_{e,0}$ and hence $F_{e+1} \subseteq^* R$. Otherwise, $S_{g \upharpoonright e+1} \subseteq V_{e,1}$ and hence $F_{e+1} \subseteq^* \bar{R}$. \square

4. MAXIMAL INDEPENDENT FAMILIES IN COMPUTABILITY

In this short section, we determine the complexity of the computability-theoretic analog of the independence number i for the Boolean algebra of computable sets. It turns out that in the context of the computable sets, *maximal independent families* behave in a way similar to ultrafilter bases.

Given a sequence $\langle F_n \rangle_{n \in \omega}$, for each binary string σ we write

$$(1) \quad F_\sigma = \bigcap_{\sigma(i)=1} F_i \cap \bigcap_{\sigma(i)=0} \bar{F}_i.$$

We call (a set F encoding) such a sequence *independent* if each set F_σ is infinite.

Definition 4.1. Given a Boolean algebra of sets \mathbb{B} , the mass problem $\mathcal{I}_\mathbb{B}$ is the class of sets F such that $\langle F_n \rangle_{n \in \omega}$ is a family that is *maximal independent*, namely, it is independent, and for each set $R \in \mathbb{B}$, there is σ such that $F_\sigma \subseteq^* R$ or $F_\sigma \subseteq^* \bar{R}$.

In the following, we let \mathbb{B} be the Boolean algebra of computable sets, and we drop the parameter \mathbb{B} as usual. An easy modification of the proof of Theorem 3.6 yields the following

Theorem 4.2. *Every maximal independent family F uniformly computes a dominating function. In other words, $\mathcal{I} \geq_s \text{DomFcn}$.*

Proof. Define the Π_1^0 -classes P_e as in Lemma 3.7. As before let $Q_e = \{X : \bar{X} \in P_e\}$ be the Π_1^0 -class of complements of elements of P_e . Recall that for any set C , we let $S_C = \{X \in 2^\omega : C \subseteq X\}$. Now we have that

$$\begin{aligned} \varphi_e \text{ is total} &\iff (\exists \sigma)(\exists n) [F_\sigma \setminus [0, n] \text{ is a subset of some} \\ &\quad X \in P_e \text{ or its complement}] \\ &\iff (\exists \sigma)(\exists n) [P_e \cap S_{F_\sigma \setminus [0, n]} \neq \emptyset \text{ or } Q_e \cap S_{F_\sigma \setminus [0, n]} \neq \emptyset] \end{aligned}$$

As before, this shows that from F one can uniformly compute a dominating function. \square

³Here, $A \subseteq_\infty B$ means that $A \subseteq B$ and $B \setminus A$ is infinite. We define \subseteq_∞^* similarly.

Theorem 4.3. *Every dominating function h uniformly computes a maximal independent family. In other words, $\mathcal{I} \leq_s \text{DomFcn}$.*

In fact, we will prove that a dominating function h uniformly computes a set F such that the $=^*$ -equivalence classes of the sets F_e freely generate the Boolean algebra of computable sets modulo finite sets.

Proof. As in the proof of Theorem 3.8, let $\langle \psi_e \rangle_{e \in \omega}$ be an effective listing of the $\{0, 1\}$ -valued partial computable functions defined on an initial segment of ω , and let $V_{e,k} = \{x : \psi_e(x) = k\}$ for $k = 0, 1$.

In Phase e of the construction, we will define a computable set F_e such that $F_e = \Theta_e^h$ for a Turing functional Θ_e determined uniformly in e . Suppose we have defined Θ_i for $i < e$, and thereby the sets F_σ defined in (1), where σ is a string of length e .

The idea for building F_e is to try to follow $V_{e,0}$ while maintaining independence from the previous sets. We apply this strategy separately on each F_σ . Using h as an oracle we compute recursively an increasing sequence $\langle r_n^e \rangle_{n \in \omega}$. We carry out the attempts on intervals $[r_n^e, r_{n+1}^e)$. If $V_{e,0}$ appears to split F_σ on the current interval, then we follow it; otherwise, we merely make sure that F_e remains independent from F_σ on the interval by putting one number in and leaving another one out. To decide which case holds, we consult the dominating function.

We now provide the details for Phase e . As e is fixed, we drop the superscripts e . By σ we will always denote a string of length e . Let $r_0 = 0$. If r_n has been defined, let $r_{n+1} > r_n$ be the least number r such that for each σ

- (a) $|[r_n, r) \cap F_\sigma| \geq 2$, and
- (b) if there are $u, w \in \text{dom}(\psi_{e,h(r_n)}) \cap F_\sigma$ with $r_n \leq u < w$ such that $\psi_e(u) = 1$ and $\psi_e(w) = 0$, then $r > w$ for the least such w .

We define $F_e(x) = \Theta_e^h(x)$ for $x \in [r_n, r_{n+1})$ as follows. For each σ ,

- if condition (b) holds and ψ_e is defined on $[r_n, r_{n+1})$, then let $F_e(x) = \psi_e(x)$;
- otherwise, if $x = \min([r_n, r_{n+1}) \cap F_\sigma)$, let $F_e(x) = 1$, else let $F_e(x) = 0$.

Verification. By induction on e , one verifies that F_σ is infinite for each σ with $|\sigma| = e$, and that the sequence $\langle r_n^e \rangle_{n \in \omega}$ defined in Phase e of the construction is infinite. Thus Θ_e^h is total for each function h . So $F \leq_T h$ where $F_e = \Theta_e^h$, and F is an independent family.

Claim 4.4. *Each set F_e is computable.*

We verify this by induction on e . Suppose it holds for each $i < e$. So F_σ is computable for $|\sigma| = e$.

First assume that ψ_e is partial. Then for sufficiently large n , condition (b) does not apply, and so the sequence $\langle r_n \rangle_{n \in \omega}$ and hence F_e are computable.

Now assume that ψ_e is total. Let

$$D_e = \{\sigma : |\sigma| = e \wedge |F_\sigma \cap V_{e,0}| = |F_\sigma \cap V_{e,1}| = \infty\}.$$

Define a function p by letting $p(m)$ be the least stage s such that for each $\sigma \notin D_e$, condition (a) holds with $r_n = m$ and $r = s$, and for each $\sigma \in D_e$, there are $u, w \in \text{dom}(\psi_{e,s})$ such that $m \leq u < w$ as in condition (b). (Let $p(m) = 0$ if m is not of the form r_n .) Since F_σ is computable for each σ of length e , the function p is computable. Since h dominates p , for sufficiently large n , we will define r_{n+1} by checking ψ_e at a stage $h(r_n) \geq p(r_n)$; since we chose the witnesses minimal, r_{n+1}

is determined by stage $p(r_n)$. So we might as well check ψ_e at that stage and do not need h . Hence the sequence $\langle r_n \rangle_{n \in \omega}$ and therefore F_e are computable.

Claim 4.5. *Suppose that ψ_e is total. Then for each string $\tau = \sigma \hat{\ } a$ of length $e + 1$, $F_\tau \subseteq^* V_{e,0}$ or $F_\tau \cap V_{e,0} =^* \emptyset$.*

If $\sigma \notin D_e$, then this is immediate since $F_\sigma \subseteq^* V_{e,i}$ for some i . Otherwise, Phase e of the construction ensures that $F_{\sigma \hat{\ } 0} =^* F_\sigma \cap V_{e,0}$.

By the last claim, the $=^*$ -equivalence classes of the F_e freely generate the Boolean algebra of the computable sets modulo finite sets. In particular, F is a maximal independent family. \square

As mentioned in the introduction, we do not know at present whether there is a “natural” Medvedev equivalence between the two mass problems \mathcal{U} and \mathcal{I} as is the case for \mathcal{A} and \mathcal{T} . This would require direct proofs avoiding the detour via the mass problem of dominating functions. For what it is worth, the cardinal characteristics \mathfrak{u} and \mathfrak{i} are incomparable (i.e., ZFC cannot determine their order).

5. THE CO-COMPUTABLY ENUMERABLE CASE

Recall from Fact 2.3 that no maximal tower, and in particular no ultrafilter base, can be computably enumerable. In contrast, in this section we will see that even ultrafilter bases can have computably enumerable complement. As in the previous sections, we are restricting our attention to the Boolean algebra of all computable sets.

Recall that a coinfinite c.e. set A is called *simple* if it meets every infinite c.e. (or, equivalently, computable) set; A is called *r -maximal* if $\overline{A} \subseteq^* \overline{R}$ or $\overline{A} \subseteq^* R$ for each computable set R . Each r -maximal set is simple. For more background, see, e.g., Soare [17].

5.1. Computably enumerable MAD sets, and co-c.e. towers. We will show that if A is a noncomputable c.e. set, then there is a co-c.e. maximal tower $G \leq_T A$. Given that it is more standard to build c.e. rather than co-c.e. sets, it will be convenient to first build a c.e. MAD set $F \leq_T A$ and then use the Medvedev reduction in Fact 2.2 to get a co-c.e. maximal tower. We employ a priority construction with requirements that act only finitely often.

Theorem 5.1. *For each noncomputable c.e. set A , there is a MAD c.e. set $F \leq_T A$.*

Proof. The construction is akin to Post’s construction of a simple set. In particular, it is compatible with permitting.

Let $\langle M_e \rangle_{e \in \omega}$ be a uniformly c.e. sequence of sets such that $M_{2e} = W_e$ and $M_{2e+1} = \omega$ for each e . We will build an auxiliary c.e. set $H \leq_T A$ and let the c.e. set $F \leq_T A$ be defined by $F^{[e]} = H^{[2e]} \cup H^{[2e+1]}$. The purpose of the sets M_{2e+1} is to make the sets $H^{[2e+1]}$, and hence the sets $F^{[e]}$, infinite. The construction also ensures that H , and hence F , is AD, and that $\bigcup_n H^{[n]}$ is coinfinite.

As usual, we will write H_e for $H^{[e]}$. We provide a stage-by-stage construction to meet the requirements

$$P_n: M_e \setminus \bigcup_{i < n} H_i \text{ infinite} \Rightarrow |H_e \cap M_e| \geq k, \text{ where } n = \langle e, k \rangle.$$

(Note that the union is over all i such that $i < n$, not $i < e$.) At stage s , we say that P_n is *permanently satisfied* if $|H_{e,s} \cap M_{e,s}| \geq k$.

Construction.

Stage $s > 0$. See if there is $n < s$ such that P_n is not permanently satisfied, and, where $n = \langle e, k \rangle$, there is $x \in M_{e,s} \setminus \bigcup_{i < n} H_{i,s}$ such that

$$x > \max(H_{e,s-1}), x \geq 2n, \text{ and } A_s \upharpoonright x \neq A_{s-1} \upharpoonright x.$$

If so, choose n least, and put $\langle x, e \rangle$ into H (i.e., put x into H_e).

Verification. Each H_e is enumerated in increasing fashion and hence computable.

Each P_n is active at most once. This ensures that $\bigcup_e H_e$ is coinfinite: For each N , if $x < 2N$ enters this union, then this is due to the action of a requirement P_n with $n < N$, so there are at most N many such x .

To see that a requirement P_n for $n = \langle e, k \rangle$ is met, suppose that its hypothesis holds. Then there are potentially infinitely many candidates x that can go into H_e . Since A is noncomputable, one of them will be permitted.

Now, by the choice of M_{2e+1} and the fact that $\bigcup_e H_e$ is coinfinite, each H_{2e+1} , and hence each F_e , is infinite. By construction, for $e < m$, we have $|H_e \cap H_m| \leq m$. So the family described by H , and therefore also the one described by F , is almost disjoint.

To show that F is MAD, it suffices to verify that if M_e is infinite then $M_e \cap F_p$ is infinite for some p . If all the $P_{e,k}$ are permanently satisfied during the construction, we let $p = e$. Otherwise, we let k be least such that P_n is never permanently satisfied where $n = \langle e, k \rangle$. Then its hypothesis fails, so $M_e \subseteq^* \bigcup_{i < n} H_i$. \square

Since an index guessable set computes no MAD set by Proposition 2.4, we obtain the following

Corollary 5.2. *No noncomputable, c.e. set L is index guessable.*

Downey and Nies have given a direct proof of this fact (see [6]).

Corollary 5.3. *For each noncomputable c.e. set A , there is a co-c.e. set $G \leq_T A$ such that $G \in \mathcal{T}$, i.e., $\langle G_n \rangle_{n \in \omega}$ is a maximal tower.*

Proof. Let F be the MAD set obtained above. Recall the Turing reduction Cp showing that $\mathcal{T} \leq_s \mathcal{A}$ in Fact 2.2. The set $G = \text{Cp}(F)$, given by

$$x \in G_n \leftrightarrow \forall i < n [x \notin F_n]$$

is as required. \square

5.2. Co-c.e. ultrafilter bases.

Theorem 5.4. *There is a co-c.e. ultrafilter base F .*

Proof. We adapt the construction from the proof of the main result in [11], which states that there is an r -maximal set A such that the index set $\text{Cof}_A = \{e: W_e \cup A =^* \omega\}$ is Σ_3^0 -complete. Both the original and the adapted version make use of the fact that we are given a c.e. index for a computable set and also one for its complement (see the pairs $(V_{e,0}, V_{e,1})$ below). Our proof can also be viewed as a variation on the proof of Theorem 3.8 in the setting of co-c.e. sets. We remark that by standard methods, one can extend the present construction to include permitting below a given high c.e. set.

We build a co-c.e. tower F by providing uniformly co-c.e. sets F_e for $e \in \omega$ that form a descending sequence with $F_e \supseteq F_{e+1}$. We achieve the latter condition by

agreeing that whenever we remove x from F_e at a stage s , we also remove it from all F_i for $i > e$. Furthermore, no element is ever removed from F_0 , so $F_0 = \omega$.

Let $\langle (V_{e,0}, V_{e,1}) \rangle_{e \in \omega}$ be an effective listing of all pairs of disjoint c.e. sets as defined in the proof of Theorem 3.8. The construction will ensure that the following requirements are met:

$$\begin{aligned} M_e &: F_e \setminus F_{e+1} \text{ is infinite,} \\ P_e &: V_{e,0} \cup V_{e,1} = \omega \Rightarrow F_{e+1} \subseteq^* V_{e,0} \vee F_{e+1} \subseteq^* V_{e,1}. \end{aligned}$$

This suffices to establish that F is an ultrafilter base.

The tree of strategies is $T = \{0, 1, 2\}^{<\omega}$. Each string $\alpha \in T$ of length e is tied to M_e and also to P_e . We write $\alpha: M_e$ and $\alpha: P_e$ to indicate that we view α as a strategy of the respective type.

Streaming. For each string $\alpha \in T$ with $|\alpha| = e$, at each stage of the construction, we have a set S_α , thought of as a stream of numbers used by α . Each time α is initialized, S_α is removed from F_{e+1} , and S_α is reset to be empty. Also, S_α is enlarged only at stages at which α appears to be on the true path. We will verify the following properties:

- (1) $S_\emptyset = \omega$;
- (2) if α is not the empty node, then S_α is a subset of S_{α^-} (where α^- is the immediate predecessor of α);
- (3) at every stage, $S_\gamma \cap S_\beta = \emptyset$ for incomparable strings γ and β ;
- (4) x is in F_{e+1} at the time a number x first enters S_α ;
- (5) if α is along the true path of the construction, then S_α is an infinite computable set.

Note that S_α is d.c.e. uniformly in α . The set S_α is finite if α is to the left of the true path of the construction; S_α is an infinite computable set if α is along the true path; and S_α is empty if α is to the right of the true path.

The *intuitive strategy* $\alpha: P_e$: Only strategies associated with a string of length $\leq e$ can remove numbers from F_{e+1} . A strategy $\alpha: P_e$ thins out S_α by removing some of its elements from F_{e+1} . It regards the set of remaining numbers as its private version of F_{e+1} . It has to make sure that no strategies β to its right remove numbers from F_{e+1} that it wants to keep. On the other hand, it can only process a number x once it knows whether x is in $V_{e,0}$ or $V_{e,1}$. The solution to this conflict is that α *reserves* a number x from the stream S_α , which by initialization withholds it from any action of such a β . It then waits until all numbers $\leq x$ are in $V_{e,0} \cup V_{e,1}$. If that never happens for some reserved x , then α is satisfied finitarily with eventual outcome 2. Otherwise, it will eventually process x : If $x \in V_{e,0}$, it continues its attempt to build F_{e+1} inside $V_{e,0}$; else it continues to build F_{e+1} inside $V_{e,1}$. It takes outcome 0 or 1, respectively, according to which case applies. Each time the apparent outcome is 0, the content of the output stream based on the assumption that the true outcome is 1 is removed from F_{e+1} . So if 0 is the true outcome, then indeed $F_{e+1} \subseteq^* V_{e,0}$.

The *intuitive strategy* $\alpha: M_e$ simply removes every other element of S_α from F_{e+1} . Then $\alpha: P_e$ actually only works with the stream of remaining numbers. There is no further interaction between the two types of strategies. (Note here that making F_{e+1} smaller is to the advantage of P_e .) Recall that if α is initialized, S_α is removed from F_{e+1} , and S_α is reset to be empty.

Construction.

Stage 0. Let δ_0 be the empty string. Let $F_e = \omega$ for each e . Initialize all strategies.

Stage $s > 0$. Let $S_{\emptyset, s} = [0, s)$. Stage s consists of substages $e = 0, \dots, s-1$, during which we inductively define δ_s , a string of length s .

Substage e . We suppose that $\alpha = \delta_s \upharpoonright e$ and S_α have been defined.

The strategy $\alpha: M_e$ acts as follows. If at the current stage $S_\alpha = \{r_0 < \dots < r_k\}$ and r_k is new in S_α , it puts r_k into \tilde{S}_α if and only if k is even; otherwise, r_k is removed from F_{e+1} .

The strategy $\alpha: P_e$ picks the first applicable case below.

Case 1: Each reserved number of α has been processed: If there is a number x from \tilde{S}_α greater than α 's last reserved number (if any) and greater than the last stage α was initialized, pick x least and *reserve* it. Note that $x < s$ since by definition $S_{\emptyset, s} = [0, s)$. Initialize $\alpha \hat{=} 2$, and let $\alpha \hat{=} 2$ be eligible to act next.

Note that if Case 1 does not apply then α has a unique reserved, but unprocessed number x .

Case 2: $[0, x] \subseteq V_{e,0} \cup V_{e,1}$ and $x \in V_{e,0}$: Let t be the greatest stage $< s$ at which α was initialized. Add x to $S_{\alpha \hat{=} 0}$ and remove from F_{e+1} all numbers in the interval (t, x) that are not in $S_{\alpha \hat{=} 0}$. Declare that α has *processed* x . Let $\alpha \hat{=} 0$ be eligible to act next.

Case 3: $[0, x] \subseteq V_{e,0} \cup V_{e,1}$ and $x \in V_{e,1}$: Let t be the greatest stage $< s$ at which α was initialized or $\alpha \hat{=} 0$ was eligible to act. Add x to $S_{\alpha \hat{=} 1}$ and remove from F_{e+1} all numbers in the interval (t, x) that are not in $S_{\alpha \hat{=} 1}$. Declare that α has *processed* x . Let $\alpha \hat{=} 1$ be eligible to act next.

Case 4: Otherwise, that is, $[0, x] \not\subseteq V_{e,0} \cup V_{e,1}$: Let t be the greatest stage $< s$ at which α was initialized, or $\alpha \hat{=} 0$ or $\alpha \hat{=} 1$ was eligible to act. Let $S_{\alpha \hat{=} 2} = \tilde{S}_\alpha \cap (t, s)$. Let $\alpha \hat{=} 2$ be eligible to act.

We define $\delta_s(e) = i$ where $\alpha \hat{=} i$, $0 \leq i \leq 2$, has been declared eligible to act next. If $e+1 < s$, then carry out the next substage. Else initialize all the strategies β such that $\delta_s <_L \beta$ and end stage s .

Verification. By construction and our convention above, F_e is co-c.e., and $F_e \supseteq F_{e+1}$ for each e .

Let $g \in 2^\omega$ denote the true path, namely, the leftmost path in $\{0, 1, 2\}^\omega$ such that $\forall e \exists^\infty s [g \upharpoonright e \preceq \delta_s]$. In the following, given e , let $\alpha = g \upharpoonright e$, and let s_α be the largest stage s such that α is initialized at stage s . Note that s_α exists: The only concern would be that α has the form $\tilde{\alpha} \hat{=} 2$ and we are at substage $|\tilde{\alpha}|$ in Case 1; however, α can only be initialized once in that way unless $\delta_r <_L \alpha$ at a later stage r , namely, when the number $\tilde{\alpha}$ has reserved is processed.

We verify a number of claims.

Claim 5.5. *The “streaming properties” (1)–(5) hold.*

(1) and (2) hold by construction.

(3) Assume this fails for incomparable γ and β , so $x \in S_\gamma \cap S_\beta$ at stage s . We may as well assume that $\gamma = \alpha \hat{=} i$ and $\beta = \alpha \hat{=} k$ where $i < k$. By construction, $k \leq 1$ is not possible, so $k = 2$. Since $x \in S_{\alpha \hat{=} i}$ and $i \leq 1$, x was reserved by α at some stage $t \leq s$. So x can never enter $S_{\alpha \hat{=} 2}$ by the initialization of $\alpha \hat{=} 2$ when x was reserved.

(4) is true by construction.

(5) holds inductively, by the definition of the true path and because S_α is enumerated in increasing fashion at stages $\geq s_\alpha$.

Claim 5.6. $F_e =^* S_\alpha$.

The claim is verified by induction on e . We show that for all $x > s_\alpha$ we have $x \in F_e$ if and only if $x \in S_\alpha$. It holds for $e = 0$ because $F_0 = S_0 = \omega$. For the inductive step, let $\gamma = g \upharpoonright (e+1)$.

First, we verify that $F_{e+1} \cap (s_\gamma, \infty) \subseteq S_\gamma$. Suppose that $x > s_\gamma$ and $x \in F_{e+1}$. Then $x \in F_e$ and $x > s_\alpha$, so by the inductive hypothesis $x \in S_\alpha$. By construction, any element x that is not promoted to S_γ is also removed from F_{e+1} unless x is the last element α reserves. However, if x is the last reserved element, then necessarily $\gamma = \widehat{\alpha}2$ and this strategy is initialized when this element is reserved, so $x < s_\gamma$ contrary to our assumption.

Next, we verify that $S_\gamma \cap (s_\gamma, \infty) \subseteq F_{e+1}$. Suppose that $x \in S_\gamma$ and $x > s_\gamma$. Then $x \in S_\alpha$, so by the inductive hypothesis $x \in F_e$. At a stage $s \geq s_\gamma$, an element x of S_α cannot be removed from F_{e+1} by a strategy $\beta >_L \alpha$ because $S_\beta \cap S_\alpha = \emptyset$ by (3) as verified above and since β can only remove elements from S_β . So x can only be removed by $\alpha: M_e$ or $\alpha: P_e$.

If $\alpha: M_e$ removes x from F_{e+1} , then $x \notin \widetilde{S}_\alpha$, which contradicts that $x \in S_\gamma$. So, by construction, the only way x can be removed from F_{e+1} is by the strategy $\alpha: P_e$, but since $x > s_\gamma$ this would mean that x is not promoted to S_γ either contrary to our assumption.

Claim 5.7. *Each requirement M_e is met, namely, $F_e \setminus F_{e+1}$ is infinite.*

To see this, recall that $\alpha = g \upharpoonright e$. By the foregoing claim, the action of $\alpha: M_e$ removes infinitely many elements of $S_\alpha \subseteq F_e$ from F_{e+1} .

Claim 5.8. *Each requirement P_e is met.*

Suppose the hypothesis of P_e holds. Then every number that α reserves is eventually processed. So either $g(e) = 0$, in which case $F_{e+1} \subseteq^* V_{e,0}$ by Claim 5.6, or $g(e) = 1$, in which case $F_{e+1} \subseteq^* V_{e,1}$, also by Claim 5.6. \square

6. ULTRAFILTER BASES FOR OTHER BOOLEAN ALGEBRAS

As mentioned, we have set up our framework to apply to general countable Boolean algebras rather than merely the Boolean algebra of the computable sets mainly with subsequent research in mind. In this last section of our paper, we provide two results in the setting of other Boolean algebras of sets.

Recall that $K(x)$ denotes the prefix-free complexity of a string x , and that a set $A \subseteq \omega$ is *K-trivial* if $\exists c \forall n K(A \upharpoonright n) \leq K(0^n) + c$. For more background on *K-trivial* sets, see, e.g., Nies [13, Ch. 5] or Downey and Hirschfeldt [5, Ch. 11]. Note that by combining results of various authors, the *K-trivial* degrees form a Turing ideal in the Δ_2^0 -degrees (see, e.g., Nies [13, Sections 5.2, 5.4]). Thus the *K-trivial* sets form a Boolean algebra.

Theorem 6.1. *There is a Δ_2^0 -ultrafilter base for the Boolean algebra of the *K-trivial* sets.*

Proof. Kučera and Slaman [10] noted that there is a Δ_2^0 -function h that dominates all functions that are partial computable in some K -trivial set. We use h in a variation of the proof of Theorem 3.8.

Let $\langle V_{e,0}, V_{e,1} \rangle_{e \in \omega}$ be a uniform listing of the K -trivials and their complements given by wtt-reductions to \emptyset' ; such a listing exists by Downey, Hirschfeldt, Nies, and Stephan [4] (see also [13, Theorem 5.3.28]). Let $T = \{0, 1\}^{<\omega}$.

For each $\alpha \in T$, we define a (possibly finite) K -trivial set S_α . Let $S_\emptyset = \omega$. Suppose we have defined the set $S_\alpha = \{r_0 < r_1 < \dots\}$. Let \tilde{S}_α contain the numbers of the form r_{2i} . Let $S_{\alpha \hat{\ } k} = \tilde{S}_\alpha \cap V_{e,k}$ for $e = |\alpha|$ and $k = 0, 1$. Note that all these sets are K -trivial.

Uniformly recursively in \emptyset' , we build sets F_e , given by nondecreasing unbounded sequences of numbers $a_0^e \leq a_1^e \leq \dots$. Suppose we have defined a_{k-1}^e . Let $\alpha \in T$ be the leftmost string of length e such that S_α has at least $k+1$ elements less than $h(k)$. If α exists, let a_k^e be the k -th element of S_α , unless this is less than a_{k-1}^e , in which case we let $a_k^e = a_{k-1}^e$.

Let $g \in 2^\omega$ denote the leftmost path in $\{0, 1\}^\omega$ such that for every e , the set $S_{g \upharpoonright e}$ is infinite. Fix e and let $\alpha = g \upharpoonright e$. Let $p(k)$ be the $(k+1)$ -st element of S_α . Since h dominates the function p , eventually in the definition of F_e we will always pick α . Hence $F_e = {}^* S_\alpha$. In particular, F_e is K -trivial. Also, the sequences $\langle a_k^e \rangle_{k \in \omega}$ are unbounded for each e , so F is Δ_2^0 . Clearly, if S_α is infinite then $S_\alpha \supset_\infty S_\beta$ for $\alpha \prec \beta$. So $F_{e+1} \subseteq_\infty^* F_e$.

To verify that F is an ultrafilter base for the K -trivials, let R be a K -trivial set. Pick e such that $R = V_{e,0}$ and $\bar{R} = V_{e,1}$. If $g(e) = 0$ then $S_{g \upharpoonright e+1} \subseteq V_{e,0}$, and hence $F_{e+1} \subseteq^* R$. Otherwise $S_{g \upharpoonright e+1} \subseteq V_{e,1}$, and hence $F_{e+1} \subseteq^* \bar{R}$. \square

Remark 6.2. Any ultrafilter base for the K -trivials must have high degree. We can see this by modifying the proof of Theorem 3.6: Every Martin-Löf random set X is Martin-Löf random relative to every K -trivial (i.e., K -trivial sets are *low for ML-randomness*). Hence neither X nor \bar{X} contains an infinite K -trivial subset.

Finally, we consider the Boolean algebra of primitive recursive sets. One says that an oracle L is of PA degree if it computes a completion of Peano arithmetic. Recall that L is of PA degree if and only if it computes a separating set for each disjoint pair of c.e. sets.

Theorem 6.3. *An oracle C computes an ultrafilter base for the primitive recursive sets if and only if C' is of PA degree relative to \emptyset' .*

Proof. We modify the proof of Jockusch and Stephan [9, Theorem 2.1]. They call a set $S \subseteq \omega$ *p-cohesive* if S is cohesive for the primitive recursive sets. Their theorem states that S is *p-cohesive* if and only if S' is of PA degree relative to \emptyset' .

\Rightarrow : Suppose that C computes an ultrafilter base F for the primitive recursive sets. Let $g \leq_T F$ be a function associated with F as in Definition 1.2. Then the range S of g is *p-cohesive*. Hence S' and therefore C' is of PA relative to \emptyset' by one implication of [9, Theorem 2.1].

\Leftarrow : We modify the proof of the other implication of [9, Theorem 2.1]. Let $\langle A_i \rangle_{i \in \omega}$ be a uniformly recursive list of all the primitive recursive sets. We call i a *primitive recursive index* for A_i (or *index*, for short). By our hypothesis on C , there is a

function $g \leq_T C'$ such that

$$\begin{aligned} |A_i \cap A_n| < |A_i \cap \bar{A}_n| &\Rightarrow g(i, n) = 0 \\ |A_i \cap \bar{A}_n| < |A_i \cap A_n| &\Rightarrow g(i, n) = 1 \end{aligned}$$

(because the conditions on the left are both Σ_2^0 , and so C' computes a separating set for them).

We inductively define a C' -computable sequence of indices $\langle e_n \rangle_{n \in \omega}$. Let e_0 be an index for ω . If e_n has been defined and $A_{e_n} = \{r_0 < r_1 < \dots\}$ (possibly finite), let e'_n be an index, uniformly obtained from e_n , such that $A_{e'_n} = \{r_0, r_2, \dots\}$. Now let

$$\begin{aligned} A_{e_{n+1}} &= A_{e'_n} \cap \bar{A}_n \text{ if } g(e'_n, n) = 0, \text{ and} \\ A_{e_{n+1}} &= A_{e'_n} \cap A_n \text{ if } g(e'_n, n) = 1. \end{aligned}$$

By induction on n , one verifies that A_{e_n} is infinite and $A_{e_{n+1}} \subset_\infty A_{e_n}$. Since $g \leq_T C'$, the numbers e_n have a uniformly C -computable approximation $\langle e_{n,x} \rangle_{x \in \omega}$.

Let the ultrafilter base $F \leq_T C$ be given by $F_n(x) = A_{e_{n,x}}(x)$. Then $F_n =^* A_{e_n}$ is primitive recursive. Since $F_{n+1} \subseteq^* \bar{A}_n$ or $F_{n+1} \subseteq^* A_n$ for each n , the set F is an ultrafilter base for the primitive recursive sets. \square

REFERENCES

- [1] Andreas R. Blass. Combinatorial cardinal characteristics of the continuum. In: Matthew Foreman and Akihiro Kanamori, editors, *Handbook of Set Theory*, volume 1, pages 395–489. Springer, Dordrecht Heidelberg London New York, 2010.
- [2] Jörg Brendle, Andrew D. Brooke-Taylor, Keng Meng Ng, and André O. Nies. An analogy between cardinal characteristics and highness properties of oracles. In: *Proceedings of the 13th Asian Logic Conference: Guangzhou, China*, pages 1–28. World Scientific, 2013. <http://arxiv.org/abs/1404.2839>.
- [3] Jörg Brendle, Vera V. Fischer, and Yurii D. Khomskii. Definable maximal independent families. *Proceedings of the American Mathematical Society*, 147(8):3547–3557, 2019.
- [4] Rodney G. Downey and Denis R. Hirschfeldt and André O. Nies, and Frank Stephan. Trivial reals. In *Proceedings of the 7th and 8th Asian Logic Conferences*, pages 103–131, Singapore, 2003. Singapore University Press.
- [5] Rodney G. Downey and Denis R. Hirschfeldt. *Algorithmic randomness and complexity*. Springer-Verlag, Berlin, 2010. 855 pages.
- [6] André O. Nies (editor). Logic Blog 2019. Available at <http://arxiv.org/abs/2003.03361>, 2019.
- [7] Noam Greenberg, Rutger Kuyper, and Daniel D. Turetsky. Cardinal invariants, non-lowness classes, and Weihrauch reducibility. *Computability*, 8(3-4):305–346, 2019.
- [8] Carl G. Jockusch, Jr. Degrees in which the recursive sets are uniformly recursive. *Canad. J. Math.*, 24:1092–1099, 1972.
- [9] Carl G. Jockusch, Jr. and Frank Stephan. A cohesive set which is not high. *Math. Log. Quart.*, 39:515–530, 1993.
- [10] Antonín Kučera and Theodore A. Slaman. Low upper bounds of ideals. *J. Symbolic Logic*, 74:517–534, 2009.
- [11] Steffen Lempp, André O. Nies, and D. Reed Solomon. On the filter of computably enumerable supersets of an r -maximal set. *Archive for Mathematical Logic*, 40(6):415–423, 2001.
- [12] Maryanthe Malliaris and Saharon Shelah. General topology meets model theory, on \mathfrak{p} and \mathfrak{t} . *Proceedings of the National Academy of Sciences*, 110(33):13300–13305, 2013.
- [13] André O. Nies. *Computability and Randomness*, volume 51 of *Oxford Logic Guides*. Oxford University Press, Oxford, 2009. 444 pages. Paperback version 2011.
- [14] Nicholas A. Rupprecht. *Effective correspondents to cardinal characteristics in Cichoń's diagram*. PhD thesis, University of Michigan, 2010.
- [15] Nicholas A. Rupprecht. Relativized Schnorr tests with universal behavior. *Arch. Math. Logic*, 49(5):555–570, 2010.

- [16] Theodore A. Slaman and Robert M. Solovay. When oracles do not help. In *Proceedings of the fourth annual workshop on Computational learning theory*, pages 379–383. Morgan Kaufmann Publishers Inc., 1991.
- [17] Robert I. Soare. *Recursively Enumerable Sets and Degrees*. Perspectives in Mathematical Logic, Omega Series. Springer-Verlag, Heidelberg, 1987.
- [18] David Soukup. Two infinite quantities and their surprising relationship. *arXiv preprint arXiv:1803.04331*, 2018.
- [19] Edward L. Wimmers. The Shelah P-point independence theorem. *Israel Journal of Mathematics*, 43:28–48, 1982.

(Lempp, Miller, Soskova) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN-MADISON, 480 LINCOLN DRIVE, MADISON, WI 53706, USA

Email address: lempp@math.wisc.edu

Email address: jmiller@math.wisc.edu

Email address: msoskova@math.wisc.edu

(Nies) DEPARTMENT OF COMPUTER SCIENCE, UNIVERSITY OF AUCKLAND, PRIVATE BAG 92019, AUCKLAND, NEW ZEALAND

Email address: andre@cs.auckland.ac.nz