

**A BASIS THEOREM FOR
 Π_1^0 CLASSES OF POSITIVE MEASURE
AND JUMP INVERSION FOR RANDOM REALS**

ROD DOWNEY AND JOSEPH S. MILLER

ABSTRACT. We extend the Shoenfield jump inversion theorem to the members of any Π_1^0 class $\mathcal{P} \subseteq 2^\omega$ with nonzero measure; i.e., for every Σ_2^0 set $S \geq_T \emptyset'$, there is a Δ_2^0 real $A \in \mathcal{P}$ such that $A' \equiv_T S$. In particular, we get jump inversion for Δ_2^0 1-random reals.¹

This paper is part of an ongoing program to study the relationship between two fundamental notions of complexity for real numbers. The first is the computational complexity of a real as captured, for example, by its Turing degree. The second is the intrinsic randomness of a real. In particular, we are interested in the 1-random reals, which were introduced by Martin-Löf [13] and represent the most widely studied randomness class. For the purposes of this introduction, we will assume that the reader is somewhat familiar with basic algorithmic randomness, as per Li-Vitányi [12], and with computability theory [18]. A review of notation and terminology will be given in Section 1.

Intuitively, a 1-random real is very complex. This complexity can be captured formally in terms of unpredictability or incompressibility, but is it reflected in the computational complexity of the real? To put this question more precisely: *which Turing degrees contain 1-random reals?* We call such degrees *1-random*. A beautiful result here is the theorem of Kučera [9] and Gács [6] that every set is Turing reducible to a 1-random. Therefore, 1-random reals can have arbitrarily high Turing degree. Moreover, Kučera proved that every degree $\mathbf{a} \geq \mathbf{0}'$ is 1-random. On the other hand, the distribution of 1-random degrees below $\mathbf{0}'$ is only partially understood.

It is well known that there is a nonempty Π_1^0 class which contains only 1-random reals. In particular, consider the complement of one of the Σ_1^0 classes in a universal Martin-Löf test. Hence, by the low basis theorem of Jockusch and Soare [7], there are low 1-random reals. Several other results on the distribution of 1-random degrees are known. For instance, minimal degrees and 1-generic degrees cannot be 1-random [11], so there are lots of Δ_2^0 degrees which do not contain 1-random reals. Furthermore, Kučera [9] proved that the 1-random degrees are not closed upwards. In particular, he constructed a Δ_2^0 PA degree \mathbf{a} which is not 1-random. Recall that a degree

2000 *Mathematics Subject Classification.* 03D28, 68Q30.

¹Kučera claims this result without proof in a remark at the end of [10].

is PA if it computes a member of every nonempty Π_1^0 class.² Applying this to a nonempty Π_1^0 class containing only 1-randoms, \mathbf{a} must bound a 1-random degree. Thus, even below $\mathbf{0}'$, the degrees of 1-randoms are not closed upwards. Frank Stephan [19] recently clarified the situation by proving the following remarkable theorem: *if \mathbf{a} is PA and 1-random, then $\mathbf{0}' \leq_T \mathbf{a}$.*

These results demonstrate that the distribution of the 1-random degrees is quite complicated, even below $\mathbf{0}'$. In the present paper, we approach the problem by considering the jumps of Δ_2^0 1-random reals. First, let us briefly consider the jumps of arbitrary 1-randoms. Jockusch and Soare [7] observed that if \mathcal{P} is a nonempty Π_1^0 class with no computable members and $S \geq_T \emptyset'$, then there is an $A \in \mathcal{P}$ such that $A' \equiv_T A \oplus \emptyset' \equiv_T S$. This is an extension of the Friedberg completeness criterion [5] to the members of Π_1^0 classes and its proof is a straightforward generalization of the low basis theorem. Because, as above, there is a nonempty Π_1^0 class containing only 1-randoms, we conclude that 1-random reals can have all possible jumps.

Our initial hope was to determine the jumps of Δ_2^0 1-random reals in the same way: simply prove that every Π_1^0 class \mathcal{P} with no computable members had Δ_2^0 members of all possible jumps. Unfortunately this plan fails badly. Cenzer [2] proved that there exists such a class which contains only GL_1 (generalized low) elements, i.e. $A \in \mathcal{P}$ implies that $A' \equiv_T A \oplus \emptyset'$. Therefore, all Δ_2^0 elements of \mathcal{P} are low.

We will instead prove a basis theorem for Π_1^0 classes of *positive measure*. Our result can be viewed as a generalization of the Shoenfield jump inversion theorem [17]. It is easy to see that the jump of a Δ_2^0 set is Σ_2^0 . Shoenfield's theorem gives a converse: if S is Σ_2^0 and $S \geq_T \emptyset'$, then there is an $A \in \Delta_2^0$ such that $A' \equiv_T S$. We extend this result by requiring A to be a member of a given Π_1^0 class with nonzero measure.

Theorem 1. *Let \mathcal{P} be a Π_1^0 class such that $\mu(\mathcal{P}) > 0$. For every Σ_2^0 set $S \geq_T \emptyset'$, there is a Δ_2^0 real $A \in \mathcal{P}$ such that $A' \equiv_T S$.*

The theorem implies jump inversion for Δ_2^0 1-randoms because there is a Π_1^0 class with nonzero measure containing only 1-random reals. In fact, any nonempty Π_1^0 class of 1-randoms will suffice, because Kurtz [11] proved that every Π_1^0 class which contains a 1-random has positive measure.

Corollary 2. *For every Σ_2^0 set $S \geq_T \emptyset'$, there is a 1-random real $A \in \Delta_2^0$ such that $A' \equiv_T S$.*

We have already seen the two extreme cases of this corollary: there is a low 1-random real and a 1-random of degree $\mathbf{0}'$. We remark that any 1-random of c.e. degree—the existence of which is guaranteed by the Kreisel basis theorem—must have degree $\mathbf{0}'$. This was noted by Kučera [9], who proved

²The PA degrees were originally defined to be the degrees of complete consistent extension of Peano Arithmetic. The equivalence of these two definitions follows from the work of Scott [16] and Solovay (unpublished).

that 1-random reals have fixed point free degree. Therefore, Arslanov's completeness criterion [1] implies that $\mathbf{0}'$ is the only 1-random c.e. degree.

Before we turn to the proof of our basis theorem, we would like to mention some other results which clarify the connection between computability and randomness, in particular, strong notions of randomness. If we replace the Σ_1^0 classes by Σ_n^0 classes in the definition (given below) of a Martin-Löf test, then the resulting randomness concept is called *n-randomness*. Kurtz [11] proved that it is equivalent to passing every Martin-Löf test relative to $\emptyset^{(n-1)}$. Kautz [8] proved that for these higher levels of randomness the Turing degrees of jumps are very constrained. In particular, if A is an n -random real, then $A^{(n-1)} \equiv_T A \oplus \emptyset^{(n-1)}$. For instance, all 2-random reals are GL_1 . Finally, Miller and Yu [14] recently found the following very interesting connection between n -randomness and Turing reducibility: *every 1-random real Turing below an n -random is also n -random*. These results and similar ones are reported in the forthcoming survey paper [4], and the forthcoming book [3]. We believe that there are a number of extremely interesting connections between computability and randomness still waiting to be found.

1. DEFINITIONS, NOTATION AND TERMINOLOGY

We consider real numbers to be elements of the Cantor space 2^ω and denote the standard measure on 2^ω by μ . For convenience, we do not distinguish between a set of natural numbers and the infinite binary sequence representing that set. For $x, y \in 2^{<\omega}$, we write $x \preceq y$ if x is a prefix of y . Similarly, $x \prec A$ means that x is a prefix of $A \in 2^\omega$. For $x \in 2^{<\omega}$, let $[x] = \{A \in 2^\omega \mid x \prec A\}$; such sets form a clopen basis for the standard topology on Cantor space. To $V \subseteq 2^{<\omega}$ we associate the open set $[V] = \bigcup_{x \in V} [x]$. If V is computably enumerable, then we call $[V]$ a Σ_1^0 class. A Π_1^0 class is the complement of a Σ_1^0 class. Alternately, a Π_1^0 class is the set of infinite paths through a Π_1^0 tree $T \subseteq 2^{<\omega}$, where a *tree* is a subset of $2^{<\omega}$ closed downward under the prefix relation. Note that there is an effective enumeration $\{\mathcal{P}_e\}_{e \in \omega}$ of all Π_1^0 classes.

Martin-Löf [13] defined the random reals to be those which avoid certain effective sets of measure zero, sets representing properties satisfied by almost no real numbers. A *Martin-Löf test* is a computable sequence $\{\mathcal{V}_i\}_{i \in \omega}$ of Σ_1^0 classes such that $\mu(\mathcal{V}_i) \leq 2^{-i}$. A real $A \in 2^\omega$ *passes* a Martin-Löf test $\{\mathcal{V}_i\}_{i \in \omega}$ if $A \notin \bigcap_{i \in \omega} \mathcal{V}_i$. A real which passes all Martin-Löf tests is called *1-random*. Martin-Löf proved that it is sufficient to consider a single *universal* test; i.e., there is a Martin-Löf test $\{\mathcal{U}_i\}_{i \in \omega}$ such that $\bigcap_{i \in \omega} \mathcal{U}_i$ is exactly the class of non-random reals. In particular, $2^\omega \setminus \mathcal{U}_1$ is a nonempty Π_1^0 class containing only 1-random reals.

2. THE PROOF

The proof of Theorem 1 can be viewed as a finite injury construction relative to the halting problem. In that sense, it is similar to Sacks' construction of a minimal degree below $\mathbf{0}'$ [15]. We require two additional ideas from the literature. The first is the method of forcing with Π_1^0 classes, which was introduced by Jockusch and Soare [7] to prove the low basis theorem. This method is used to ensure that $A' \leq_T S$. The second is a version of a lemma of Kučera [9] which allows us to recursively bound the positions of branchings in a Π_1^0 class with nonzero measure. The lemma allows us to code S into A' using a variation of a process known as Kučera coding.

Lemma 3 (Kučera, 1985). *Let \mathcal{P} be a Π_1^0 class such that $\mu(\mathcal{P}) > 0$. Then there a Π_1^0 subclass $\mathcal{Q} \subseteq \mathcal{P}$ and a computable function $g: \omega \rightarrow \omega$ such that $\mu(\mathcal{Q}) > 0$ and*

$$(\forall e) \left[\mathcal{Q} \cap \mathcal{P}_e \neq \emptyset \implies \mu(\mathcal{Q} \cap \mathcal{P}_e) \geq 2^{-g(e)} \right].$$

To see why this lemma is true, let g be any computable function such that $\sum_{e \in \omega} 2^{-g(e)} < \mu(\mathcal{P})$. Let \mathcal{Q} be the Π_1^0 subclass of \mathcal{P} obtained by removing the reals in $\mathcal{P}_e[s]$ (the stage s approximation to \mathcal{P}_e) whenever $\mathcal{P}_e[s] \cap \mathcal{Q}[s]$ has measure less than $2^{-g(e)}$. The choice of g guarantees that $\mu(\mathcal{Q}) > 0$.³

Proof of Theorem 1. We are given a Π_1^0 class $\mathcal{P} \subseteq 2^\omega$ with nonzero measure and a Σ_2^0 set $S \geq_T \emptyset'$. Take the Π_1^0 class $\mathcal{Q} \subseteq \mathcal{P}$ and computable function $g: \omega \rightarrow \omega$ guaranteed by Lemma 3. We will construct a Δ_2^0 real $A \in \mathcal{Q}$ such that $A' \equiv_T S$.

Before describing the construction we must give a few preliminary definitions. For every $\sigma \in 2^{<\omega}$, define a Π_1^0 class

$$\mathcal{F}_\sigma = \{B \in \mathcal{Q} \mid (\forall e < |\sigma|) \sigma(e) = 0 \implies \varphi_e^B(e) \uparrow\}.$$

At each stage $s \in \omega$ of the construction, we will define a string $\sigma_s \in 2^s$ which is intended to approximate $A' \upharpoonright s$. We will tentatively restrict A to the class \mathcal{F}_{σ_s} in order to *force its jump*. It is important to note that this restriction may be injured at a later stage by the enumeration of an $e < s$ into S .

Next we define a computable function $f: \omega \rightarrow \omega$ which grows fast enough to ensure that it (eventually) bounds the branchings between elements of \mathcal{F}_σ , for every $\sigma \in 2^{<\omega}$. We will use f to code elements of S into A (or more precisely, into A'). Let $h: 2^{<\omega} \times 2^{<\omega} \rightarrow \omega$ be a computable function such that $\mathcal{P}_{h(x,\sigma)} = [x] \cap \mathcal{F}_\sigma$, for all $x, \sigma \in 2^{<\omega}$. Set $f(0) = 0$. For $s \in \omega$, inductively define

$$f(s+1) > \max\{g(h(x,\sigma)) \mid x \in 2^{f(s)} \text{ and } \sigma \in 2^s\}.$$

Now take $x \in 2^{f(t)}$ and $\sigma \in 2^s$, for $t \geq s$, such that $[x] \cap \mathcal{F}_\sigma \neq \emptyset$. We claim that x has distinct finite extensions $y_0, y_1 \in 2^{f(t+1)}$ which extend to reals in \mathcal{F}_σ . Assume not. Let $\hat{\sigma} = \sigma 1^{t-s}$ and note that $\mathcal{F}_{\hat{\sigma}} = \mathcal{F}_\sigma$. Then

³We thank the referee for suggesting this simple proof of Lemma 3.

$\mu(\mathcal{Q} \cap \mathcal{P}_{h(x, \hat{\sigma})}) = \mu([x] \cap \mathcal{F}_\sigma) \leq 2^{-f(t+1)} < 2^{-g(h(x, \hat{\sigma}))}$, which contradicts the lemma.

Kučera used the fact that we can bound branchings in a Π_1^0 class with nonzero measure to code information into members of such a class. The most basic form of Kučera coding constructs a real by extensions, choosing the leftmost or rightmost permissible extension to encode the next bit. For our construction, we only distinguish between the rightmost extension and any other permissible extension. Let \mathcal{R} be a Π_1^0 class and let $x \in 2^{f(s+1)}$, for some $s \in \omega$. Define $H_f(\mathcal{R}; x)$ to be true iff

$$(\exists n) \left[\begin{array}{l} \text{if } y \in 2^{f(s+1)} \text{ extends } x \upharpoonright f(s) \text{ and is} \\ \text{to the right of } x, \text{ then } \mathcal{R}[n] \cap [y] = \emptyset \end{array} \right],$$

where $\mathcal{R}[n]$ is the approximation to \mathcal{R} at stage n . Note that $H_f(\mathcal{R}; x)$ is a Σ_1^0 condition. By compactness, if $\mathcal{R} \cap [y] = \emptyset$, then there is a $n \in \omega$ such that $\mathcal{R}[n] \cap [y] = \emptyset$. This implies that if $\mathcal{R} \cap [x] \neq \emptyset$, then $H_f(\mathcal{R}; x)$ is true iff x is the rightmost length $f(s+1)$ extension of $x \upharpoonright f(s)$ which extends to an element of \mathcal{R} .

It will be useful to understand the interaction between f and H_f . Assume that we have $x \in 2^{f(t)}$, for some $t \geq s$, and $\sigma \in 2^s$ such that $[x] \cap \mathcal{F}_\sigma \neq \emptyset$. Let $\hat{x} \in 2^{f(t+1)}$ be the leftmost extension of x such that $[\hat{x}] \cap \mathcal{F}_\sigma \neq \emptyset$. By the definition of f , there are multiple extensions to choose from, so $H_f(\mathcal{F}_\sigma, \hat{x})$ is false. In fact, if $\tau \preceq \sigma$, then $\mathcal{F}_\sigma \subseteq \mathcal{F}_\tau$. Therefore, $H_f(\mathcal{F}_\tau, \hat{x})$ is also false.

We are now ready to describe the construction. Let $\{S_s\}_{s \in \omega}$ be a \emptyset' -computable enumeration of the Σ_2^0 set S . We may assume that $S_0 = \emptyset$ and $|S_{s+1} \setminus S_s| = 1$ for all $s \in \omega$. We construct A by initial segments using a \emptyset' oracle. At each stage $s \in \omega$, we find a string $x_s \in 2^{f(t)}$, for some $t \geq s$. Define $A = \bigcup_s x_s$. Each stage also produces a string $\sigma_s \in 2^s$, which is an approximation to A' , although not necessarily an initial segment of it. For each $s \in \omega$, we require that

- (1) $[x_s] \cap \mathcal{F}_{\sigma_s} \neq \emptyset$.
- (2) If $e < s$ and $B \in [x_s] \cap \mathcal{F}_{\sigma_s \upharpoonright e+1}$, then $B'(e) = \sigma_s(e)$.

The Construction.

Stage 0. Let x_0 and σ_0 both be the empty string. Then $[x_0] \cap \mathcal{F}_{\sigma_0} = \mathcal{Q} \neq \emptyset$, so (1) is satisfied. Note that (2) is vacuous.

Stage $s+1$. Assume that we have already constructed $x_s \in 2^{f(t)}$, for some $t \geq s$, and $\sigma_s \in 2^s$ satisfying (1) and (2). Let $e \in S_{s+1} \setminus S_s$ (this element is unique).

Case 1. If $e > s$, then let $x_{s+1} \in 2^{f(t+1)}$ be the leftmost extension of x_s such that $[x_{s+1}] \cap \mathcal{F}_{\sigma_s} \neq \emptyset$. Note that \emptyset' can determine if $[y] \cap \mathcal{F}_{\sigma_s} = \emptyset$, for each $y \in 2^{<\omega}$, so \emptyset' can find x_{s+1} . If $[x_{s+1}] \cap \mathcal{F}_{\sigma_s 0} \neq \emptyset$, then let $\sigma_{s+1} = \sigma_s 0$. Otherwise, let $\sigma_{s+1} = \sigma_s 1$. Again, this can be determined using the \emptyset' oracle. Note that (1) and (2) are satisfied by our choices of x_{s+1} and σ_{s+1} .

Case 2. If $e \leq s$, then let $\tau = \sigma_s \upharpoonright e$. Consider the least number $m \in \omega$ such that $f(\langle e, m \rangle) \geq |x_s|$. First define $\hat{x}_s \in 2^{f(\langle e, m \rangle)}$ to be the leftmost

extension of x_s such that $[\widehat{x}_s] \cap \mathcal{F}_\tau \neq \emptyset$. Next let $x_{s+1} \in 2^{f(\langle e, m \rangle + 1)}$ be the rightmost extension of \widehat{x}_s such that $[x_{s+1}] \cap \mathcal{F}_\tau \neq \emptyset$. Let σ_{s+1} be the lexicographically least string of length $s + 1$ which extends τ and satisfies $[x_{s+1}] \cap \mathcal{F}_{\sigma_{s+1}} \neq \emptyset$. Again, the construction is computable relative to \emptyset' and we have ensured that (1) and (2) continue to hold.

End Construction.

We turn to the verification. The construction is computable from a \emptyset' oracle, so A is Δ_2^0 . Furthermore, (1) tells us that $[x_s] \cap \mathcal{F}_{\sigma_s} \neq \emptyset$, for each $s \in \omega$. Because $\mathcal{F}_{\sigma_s} \subseteq \mathcal{Q} \subseteq \mathcal{P}$, this implies that every x_s can be extended to an element of \mathcal{P} . But \mathcal{P} is closed, so $A = \bigcup_s x_s \in \mathcal{P}$. All that remains to verify is that $A' \equiv_T S$.

First we prove that $A' \leq_T S$. To determine whether $e \in A'$, use S and \emptyset' to find a stage $s > e$ such that $S_s \upharpoonright e + 1 = S \upharpoonright e + 1$. Let $\tau = \sigma_s \upharpoonright e + 1$. We claim that $\sigma_t \upharpoonright e + 1 = \tau$, for all $t \geq s$. This is because the only way that $\sigma_s \upharpoonright e + 1$ can be injured during the construction is in Case 2, when an element $i \leq e$ is enumerated into S . But this will never happen after stage s . Therefore, for all $t \geq s$, we have $\tau \preceq \sigma_t$ and hence $\mathcal{F}_{\sigma_t} \subseteq \mathcal{F}_\tau$. So $[x_t] \cap \mathcal{F}_\tau \neq \emptyset$, for all $t \geq s$, which implies that $A \in \mathcal{F}_\tau$. By (2), we have $A'(e) = \tau(e)$. This proves that we can uniformly decide if $e \in A'$ using only $S \oplus \emptyset' \equiv_T S$. Therefore, $A' \leq_T S$.

Now we show that $S \leq_T A'$. Assume by induction that we have determined $S \upharpoonright e$, for some $e \in \omega$. Find the least $s \geq e$ such that $S_s \upharpoonright e = S \upharpoonright e$. Let $\tau = \sigma_s \upharpoonright e$ and note, as above, that $\tau \preceq \sigma_t$, for all $t \geq s$. Find the least $m \in \omega$ such that $f(\langle e, m \rangle) \geq |x_s|$. Of course, both s and m can be found by \emptyset' . We claim that $e \in S$ iff either $e \in S_s$ or $(\exists n \geq m) H_f(\mathcal{F}_\tau; A \upharpoonright f(\langle e, n \rangle + 1))$. If $e \in S \setminus S_s$, then Case 2 ensures that $H_f(\mathcal{F}_\tau; A \upharpoonright f(\langle e, n \rangle + 1))$ holds for some $n \geq m$. So, assume that $e \notin S$. Then for every $n \geq m$, the construction chooses the leftmost extension of $A \upharpoonright f(\langle e, n \rangle)$ which is extendible in the appropriate Π_1^0 class. This class is of the form $\mathcal{F}_{\widehat{\tau}}$, where $\widehat{\tau} \preceq \sigma_t$ for some $t \geq s$ and $|\widehat{\tau}| \geq e$. This implies that $\tau \preceq \widehat{\tau}$, so $\mathcal{F}_{\widehat{\tau}} \subseteq \mathcal{F}_\tau$. The definition of f ensures that there are distinct length $f(\langle e, n \rangle + 1)$ extensions of $A \upharpoonright f(\langle e, n \rangle)$ which can be extended to elements of $\mathcal{F}_{\widehat{\tau}}$. Therefore, the leftmost choice consistent with $\mathcal{F}_{\widehat{\tau}}$ must be left of the rightmost choice consistent with \mathcal{F}_τ . Hence $H_f(\mathcal{F}_\tau; A \upharpoonright f(\langle e, n \rangle + 1))$ is false. Finally, note that A' can decide if $(\exists n \geq m) H_f(\mathcal{F}_\tau; A \upharpoonright f(\langle e, n \rangle + 1))$, because H_f is Σ_1^0 . Therefore A' can determine if $e \in S$, proving that $S \leq_T A'$. \square

REFERENCES

- [1] M. M. Arslanov, R. F. Nadyrov, and V. D. Solov'ev. A criterion for the completeness of recursively enumerable sets, and some generalizations of a fixed point theorem. *Izv. Vysš. Učebn. Zaved. Matematika*, 4 (179):3–7, 1977.
- [2] Douglas Cenzer. Π_1^0 classes in computability theory. In *Handbook of computability theory*, volume 140 of *Stud. Logic Found. Math.*, pages 37–85. North-Holland, Amsterdam, 1999.
- [3] R. Downey and D. Hirschfeldt. *Algorithmic randomness and complexity*. Springer-Verlag, Berlin. to appear.

- [4] Rodney G. Downey, Denis R. Hirschfeldt, André Nies, and Sebastiaan A. Terwijn. Calibrating randomness. *Bulletin of Symbolic Logic*. to appear.
- [5] Richard Friedberg. A criterion for completeness of degrees of unsolvability. *J. Symb. Logic*, 22:159–160, 1957.
- [6] Péter Gács. Every sequence is reducible to a random one. *Inform. and Control*, 70(2-3):186–192, 1986.
- [7] Carl G. Jockusch, Jr. and Robert I. Soare. Π_1^0 classes and degrees of theories. *Trans. Amer. Math. Soc.*, 173:33–56, 1972.
- [8] S. Kautz. *Degrees of random sets*. Ph.D. Dissertation, Cornell University, 1991.
- [9] Antonín Kučera. Measure, Π_1^0 -classes and complete extensions of PA. In *Recursion theory week (Oberwolfach, 1984)*, volume 1141 of *Lecture Notes in Math.*, pages 245–259. Springer, Berlin, 1985.
- [10] Antonín Kučera. On the use of diagonally nonrecursive functions. In *Logic Colloquium '87 (Granada, 1987)*, volume 129 of *Stud. Logic Found. Math.*, pages 219–239. North-Holland, Amsterdam, 1989.
- [11] S. Kurtz. *Randomness and genericity in the degrees of unsolvability*. Ph.D. Dissertation, University of Illinois, Urbana, 1981.
- [12] M. Li and P. Vitányi. *An introduction to Kolmogorov complexity and its applications*. Texts and Monographs in Computer Science. Springer-Verlag, New York, 1993.
- [13] Per Martin-Löf. The definition of random sequences. *Information and Control*, 9:602–619, 1966.
- [14] Joseph S. Miller and Liang Yu. On initial segment complexity and degrees of randomness. to appear.
- [15] Gerald E. Sacks. A minimal degree less than $0'$. *Bull. Amer. Math. Soc.*, 67:416–419, 1961.
- [16] Dana Scott. Algebras of sets binumerable in complete extensions of arithmetic. In *Proc. Sympos. Pure Math., Vol. V*, pages 117–121. American Mathematical Society, Providence, R.I., 1962.
- [17] J. R. Shoenfield. On degrees of unsolvability. *Ann. of Math. (2)*, 69:644–653, 1959.
- [18] Robert I. Soare. *Recursively enumerable sets and degrees*. Springer-Verlag, Berlin, 1987.
- [19] Frank Stephan. Martin-Löf random sets and PA-complete sets. to appear.

SCHOOL OF MATHEMATICAL AND COMPUTING SCIENCES, VICTORIA UNIVERSITY, P.O. BOX 600, WELLINGTON, NEW ZEALAND

E-mail address: Rod.Downey@mcs.vuw.ac.nz

E-mail address: Joe.Miller@mcs.vuw.ac.nz