

HIGHER ORDER DEGREES OF AFFINE PLANE CURVE COMPLEMENTS

EVA ELDUQUE AND LAURENTIU MAXIM

ABSTRACT. We study finiteness (and vanishing) properties of the higher order degrees associated to complements of complex affine plane curves with mild singularities at infinity. Our results impose new obstructions on the class of groups that can be realized as fundamental groups of affine plane curve complements. We also clarify the relationship between the higher order degrees and the multivariable Alexander polynomial of a non-irreducible plane curve.

1. INTRODUCTION

In knot theory, a strategy to address problems that the Alexander polynomial is not strong enough to solve is to consider non-abelian invariants (e.g., see [2]). These are Alexander-type invariants of coverings corresponding to terms of the derived series of a knot group, and share most of the properties of the classical Alexander invariants. Despite the difficulties of working with modules over non-commutative rings, there are applications to estimating knot genus, detecting fibered, prime and alternating knots, and to knot concordance. Higher order Alexander invariants can be associated to any finitely presented group $G = \pi_1(X)$, in terms of coverings of X given by the terms in the rational derived series of G . These in turn have striking applications if one considers the fundamental group of a link complement or that of a closed 3-manifold [6]. For example, they can be used to obtain lower bounds for the Thurston norm, and provide new algebraic obstructions to a 4-manifold of the form $M^3 \times S^1$ admitting a symplectic structure.

Motivated by their success in the classical knot theory and low-dimensional topology, C. Leidy and the second author initiated in [11] the study of higher order Alexander-type invariants for complex affine plane curve complements. The exploration of topology of complex plane curves and of their complements is a subject that goes back to works of Zariski, Enriques, Hirzebruch, Deligne, or Fulton, and which has flourished in more recent research endeavors by Libgober, Dimca, Suciu, Artal-Bartolo, Cogolludo-Agustín, etc. As the fundamental group of a plane curve complement is in general highly non-abelian, one typically considers invariants of the fundamental group that still capture most of the topology of the curve, but which are more manageable, e.g., Alexander-type invariants (see [3], [9], [11], [12], [15], [17], etc.).

To any affine plane curve $C \subset \mathbb{C}^2$, in [11] one associates a sequence $\{\delta_n(C)\}_n$ of (possibly infinite) integers, called the *higher order degrees* of C . Roughly speaking, these integers measure the “sizes” of quotients of successive terms in the rational derived series $\{G_r^{(n)}\}_{n \geq 0}$ of the fundamental group $G := \pi_1(\mathbb{C}^2 \setminus C)$ of the curve complement (see Definition 1). It was also noted in [11] that the higher order degrees of plane curves (at any level n) are sensitive to the “position” of singular points, this being one of the initial motivations for adapting and studying Alexander-type invariants in the context of plane curve complements.

Date: February 21, 2019.

2010 Mathematics Subject Classification. 32S25, 32S55, 32S05, 32S20, 57M27.

Key words and phrases. plane curve complement, essential line arrangement, singularities, higher order degrees, Alexander polynomial.

While in theory higher order degrees of a plane curve complement can be computed by Fox free calculus from a presentation of $G = \pi_1(\mathbb{C}^2 \setminus C)$, such calculations are in general tedious, see [12] for some examples. Furthermore, as these integers can also be interpreted as Betti-type invariants associated to the tower of coverings of $\mathbb{C}^2 \setminus C$ corresponding to the subgroups $G_r^{(i)}$ (the first of which is the universal abelian cover), a priori there is no reason to expect that such invariants have any good vanishing or finiteness properties. The main result of [11] proved that for curves in general position at infinity (i.e., whose projective completion is transversal to the line at infinity in $\mathbb{C}P^2$) these higher order degrees are in fact *finite*, and a uniform upper bound was given only in terms of the degree m of the curve by comparing with the corresponding invariants of the (m -Hopf) link at infinity. More precisely, one has the following result:

Theorem 1. [11, Corollary 4.8] *If $C \subset \mathbb{C}^2$ is a reduced plane curve of degree m , in general position at infinity, then:*

$$\delta_n(C) \leq m(m-2), \text{ for all } n.$$

One of the goals of this paper is to provide generalizations of Theorem 1 to various contexts in which the assumption of good behavior at infinity is relaxed. As the sequence of higher order degrees of a plane curve is an invariant of the fundamental group of the complement, a better understanding of its properties (such as finiteness) will impose new obstructions on the class of groups that can be realized as fundamental groups of affine plane curve complements.

Our first result generalizes Theorem 1 to the context of essential complex line arrangements. (Note that such line arrangements are not necessarily in general position at infinity.) We prove the following (see Theorem 7):

Theorem 2. *Assume that the complex affine plane curve C defines an essential line arrangement $\mathcal{A} = \{L_1, \dots, L_m\} \subset \mathbb{C}^2$ (that is, not all lines in \mathcal{A} are parallel). Then,*

$$\delta_n(C) \leq m(m-2), \text{ for all } n.$$

Moreover, the equality holds only in the case where \mathcal{A} consists of m lines going through one point, and in that case the equality holds for all $n \geq 0$.

In the special case when an arrangement contains one line which meets all other lines transversally, we show in Theorem 8 the following result (which was asserted without a proof in [12]):

Theorem 3. *Assume that the affine plane curve C defines a line arrangement $\mathcal{A} = \{L_1, \dots, L_m\} \subset \mathbb{C}^2$, which is obtained from an essential line arrangement $\mathcal{A}' = \{L_1, \dots, L_{m-1}\}$ by adjoining a line L_m that is transversal to every line in \mathcal{A}' (that is, the singularities of the curve C along the irreducible component L_m consist of $m-1$ nodes). Then*

$$\delta_n(C) = 0, \text{ for all } n \geq 0.$$

At the opposite spectrum, i.e., if the plane curve C defines a line arrangement $\mathcal{A} = \{L_1, \dots, L_m\} \subset \mathbb{C}^2$ consisting of m distinct parallel lines, then an easy calculation shows that (see Proposition 1):

$$\delta_n(C) = \begin{cases} \infty, & m > 1, \\ 0, & m = 1, \end{cases}$$

for all $n \geq 0$.

We also prove the following generalization of Theorem 1 in the context when the plane curve C is allowed to have mild singularities at infinity. More precisely, we show the following (see Theorem 9):

Theorem 4. *Let $C \subset \mathbb{C}^2$ be a reduced plane curve of degree m , let \overline{C} be its closure in $\mathbb{C}P^2$ and let L_∞ be the line at infinity. Suppose that the intersections of L_∞ and \overline{C} are either transversal or L_∞ is the tangent line to \overline{C} at a smooth point and it is a simple tangent there (i.e., it has multiplicity 2). If any of the following two conditions hold*

- (a) $m = 2$;
- (b) *at least one of the intersections of \overline{C} and L_∞ is transversal,*

then

$$\delta_n(C) \leq m(m - 2),$$

for all $n \geq 0$.

Moreover, we generalize Theorem 3 to the context of plane curves as follows (see Theorem 10):

Theorem 5. *Let $n \geq 0$. Assume that the affine plane curve C is of the form $C = L \cup C'$, where C' is a curve of degree $m - 1$ in \mathbb{C}^2 such that $\delta_n(C')$ is finite, and L is a line transversal to C' such that $L \cap C'$ consists of $m - 1$ distinct points. Then,*

$$\delta_n(C) = 0.$$

A natural question in the context of non-commutative Alexander-type invariants is to relate the higher order degrees of a plane curve complement to the previously studied Alexander-type invariants, such as the Alexander polynomials. Preliminary steps in this direction have already been made in [11], where the authors showed that if the (one-variable) Alexander polynomial of an irreducible plane curve is trivial then *all* higher-order degrees $\{\delta_n\}_n$ vanish. (If the curve is irreducible, then $\delta_0(C)$ is the degree of the Alexander polynomial of C .) In relation with the universal abelian invariants of a curve, it was also noted in [11] that if the codimension (in the character torus) of the first characteristic variety of the plane curve complement is > 1 then $\delta_0(C) = 0$ (this fact was first pointed out by A. Libgober in an informal conversation with the second author, see also Corollary 1). However, curves (e.g., in general position at infinity) may have supports of codimension one in the character variety (cf. [10]), and for this boundary case we show here that $\delta_0(C)$ is the degree of the multivariable Alexander polynomial Δ_C of the plane curve C (see Theorem 12).

The paper is structured as follows. In Section 2, we recall the definition of higher order degrees of an affine plane curve complement. Section 3 supplies proofs of Theorems 2 and 3. Theorems 4 and 5 are proved in Section 4. Finally, in Section 5, we indicate the relation between $\delta_0(C)$ and the degree of the multivariable Alexander polynomial of the plane curve complement in the case when C is not irreducible.

Acknowledgements. L. Maxim was partially supported by the Simons Foundation Collaboration Grant #567077 and by CNCS-UEFISCDI grant PN-III-P4-ID-PCE-2016-0030.

2. HIGHER-ORDER INVARIANTS OF A PLANE CURVE COMPLEMENT

Though most of the background material presented in this section applies to any finitely presented group, we focus mainly on fundamental groups of complex affine plane curve complements.

Let $C = \{f(x, y) = 0\}$ be a reduced curve in \mathbb{C}^2 of degree m , with complement

$$U := \mathbb{C}^2 \setminus C,$$

and denote by $G := \pi_1(U)$ the fundamental group of its complement. If C has s irreducible components, then

$$(1) \quad H_1(G; \mathbb{Z}) = H_1(U; \mathbb{Z}) = G/G' = \mathbb{Z}^s,$$

generated by meridian loops about the smooth parts of the irreducible components of C .

In this section we recall the definition of the higher-order Alexander-type invariants of the group G . These were originally used in the study of knots and, respectively, 3-manifolds, see e.g., [2, 6], and they were ported to the study of plane curve complements in [11], e.g., to show that certain groups cannot be realized as fundamental groups of such complements.

Definition 1. The *rational derived series* of the group G is defined as follows: $G_r^{(0)} = G$, and for $n \geq 1$,

$$G_r^{(n)} = \{g \in G_r^{(n-1)} \mid g^k \in [G_r^{(n-1)}, G_r^{(n-1)}], \text{ for some } k \in \mathbb{Z} \setminus \{0\}\}.$$

It is easy to see that $G_r^{(i)} \triangleleft G_r^{(j)} \triangleleft G$, if $i \geq j \geq 0$. The successive quotients of the rational derived series are torsion-free abelian groups. In fact (cf. [6, Lemma 3.5]),

$$G_r^{(n)} / G_r^{(n+1)} \cong \left(G_r^{(n)} / [G_r^{(n)}, G_r^{(n)}] \right) / \{\mathbb{Z} - \text{torsion}\}.$$

Therefore, for $G = \pi_1(\mathbb{C}^2 \setminus C)$ we get from (1) that $G' = G_r^{(1)}$.

The use of the rational derived series as opposed to the usual derived series is needed in order to avoid zero-divisors in the group ring $\mathbb{Z}\Gamma_n$, where

$$\Gamma_n := G / G_r^{(n+1)}.$$

By construction, Γ_n is a poly-torsion-free-abelian group, in short a PTFA ([6, Corollary 3.6]), i.e., it admits a normal series of subgroups such that each of the successive quotients of the series is torsion-free abelian. Then $\mathbb{Z}\Gamma_n$ is a right and left Ore domain, so it embeds in its classical right ring of quotients \mathcal{K}_n , a skew-field. Every module over \mathcal{K}_n is a free module, and such modules have a well-defined rank $\text{rk}_{\mathcal{K}_n}$ which is additive on short exact sequences. (These statements also apply to the right ring of quotients \mathcal{K} of the group ring $\mathbb{Z}\Gamma$ of any PTFA group Γ , e.g., see [11, Remark 2.4] and the references therein.)

Definition 2. The *n-th order Alexander module* of (the complement of) the plane curve C is defined as

$$\mathcal{A}_n^{\mathbb{Z}}(C) = H_1(U; \mathbb{Z}\Gamma_n) = H_1(U_{\Gamma_n}; \mathbb{Z}),$$

where U_{Γ_n} is the covering of U corresponding to the subgroup $G_r^{(n+1)}$. That is,

$$\mathcal{A}_n^{\mathbb{Z}}(C) = G_r^{(n+1)} / [G_r^{(n+1)}, G_r^{(n+1)}],$$

viewed as a right $\mathbb{Z}\Gamma_n$ -module.

The *n-th order rank* of (the complement of) C is:

$$r_n(C) = \text{rk}_{\mathcal{K}_n} H_1(U; \mathcal{K}_n).$$

Remark 1. Note that $\mathcal{A}_0^{\mathbb{Z}}(C) = G_r^{(1)} / [G_r^{(1)}, G_r^{(1)}] = G' / G''$, which is usually referred to as the Alexander invariant of the complement (see also Definition 7).

Example 1. If the curve C is in general position at infinity (i.e., the line at infinity in $\mathbb{C}P^2$ is transversal to the projective completion of C), and it is nonsingular or has only nodal singular points (i.e., locally defined by $x^2 - y^2 = 0$), then $G = \pi_1(\mathbb{C}^2 \setminus C)$ is abelian, and therefore $\mathcal{A}_n^{\mathbb{Z}}(C) = 0$ for all n (e.g., see [11, Remark 3.4]).

In [11], one associates to any plane curve C (or, equivalently, to the fundamental group G of its complement) a sequence of non-negative integers $\delta_n(C)$ as follows (it is more convenient to work over a principal ideal domain, or a PID for short, so we look for a “convenient” one): Let $\psi \in H^1(G; \mathbb{Z})$ be the primitive class representing the linking number homomorphism

$$G \xrightarrow{\psi} \mathbb{Z}, \alpha \mapsto \text{lk}(\alpha, C).$$

Since G' is in the kernel of ψ , we have a well-defined induced epimorphism $\bar{\psi} : \Gamma_n \rightarrow \mathbb{Z}$. Let $\bar{\Gamma}_n = \ker \bar{\psi}$. Then $\bar{\Gamma}_n$ is a PTFA group, so $\mathbb{Z}\bar{\Gamma}_n$ has a right ring of quotients

$$\mathbb{K}_n = (\mathbb{Z}\bar{\Gamma}_n)S_n^{-1},$$

where $S_n = \mathbb{Z}\bar{\Gamma}_n \setminus \{0\}$. Set

$$R_n := (\mathbb{Z}\Gamma_n)S_n^{-1}.$$

Then R_n is a flat left $\mathbb{Z}\Gamma_n$ -module.

A very important role in what follows is played by the fact that R_n is a PID; in fact, R_n is isomorphic to the ring of skew-Laurent polynomials $\mathbb{K}_n[t^{\pm 1}]$. This can be seen as follows: by choosing a $t \in \Gamma_n$ such that $\bar{\psi}(t) = 1$, we get a splitting ϕ of $\bar{\psi}$, and the embedding $\mathbb{Z}\bar{\Gamma}_n \subset \mathbb{K}_n$ extends to an isomorphism $R_n \cong \mathbb{K}_n[t^{\pm 1}]$. However this isomorphism depends in general on the choice of splitting of $\bar{\psi}$.

Definition 3. (1) The n -th order localized Alexander module of the plane curve C is defined to be

$$\mathcal{A}_n(C) = H_1(U; R_n),$$

viewed as a right R_n -module. If we choose a splitting ϕ to identify R_n with $\mathbb{K}_n[t^{\pm 1}]$, we define $\mathcal{A}_n^\phi(C) = H_1(U; \mathbb{K}_n[t^{\pm 1}])$.

(2) The n -th order degree of C is defined to be:

$$\delta_n(C) = \text{rk}_{\mathbb{K}_n} \mathcal{A}_n(C) = \text{rk}_{\mathbb{K}_n} \mathcal{A}_n^\phi(C).$$

Remark 2. Note that $\delta_n(C) < \infty$ if and only if $\text{rk}_{\mathcal{K}_n} H_1(U; \mathcal{K}_n) = 0$, i.e. $\mathcal{A}_n(C)$ is a torsion R_n -module.

Remark 3. If the plane curve C is irreducible, then $\delta_0(C)$ is the degree of the Alexander polynomial of C ; see [11, Remark 3.9].

The higher order degrees $\delta_n(C)$ are integer invariants of the fundamental group G of the complement (endowed with the linking number homomorphism). Indeed, by [6], one has:

$$(2) \quad \delta_n(C) = \text{rk}_{\mathbb{K}_n} \left(G_r^{(n+1)} / [G_r^{(n+1)}, G_r^{(n+1)}] \otimes_{\mathbb{Z}\bar{\Gamma}_n} \mathbb{K}_n \right).$$

In fact, the use of the linking number homomorphism makes these invariants depend on the pair (\mathbb{C}^2, C) , rather than on the curve complement U . Note that since the isomorphism between R_n and $\mathbb{K}_n[t^{\pm 1}]$ depends on the choice of splitting, one cannot define in a canonical way a higher-order version of the Alexander polynomial. However, for any choice of splitting, the degree of the associated higher-order Alexander polynomial is the same, hence this yields a well-defined invariant of the group G endowed with the linking number homomorphism, which is exactly the higher-order degree δ_n defined above.

The higher-order degrees of C may be computed by means of Fox free calculus from a presentation of $G = \pi_1(\mathbb{C}^2 \setminus C)$, see [6, Section 6] for details. Such computational techniques will be used freely in this paper.

It was shown in [11] that if C is an irreducible plane curve, or a curve in general position at infinity (i.e., for which the line at infinity in $\mathbb{C}P^2$ is transversal to the projective completion of C), then the higher-order degrees $\delta_n(C)$ are finite. More precisely, one has the following:

Theorem 6. *If $C \subset \mathbb{C}^2$ is a reduced plane curve of degree m , in general position at infinity, then:*

$$\delta_n(C) \leq m(m - 2), \text{ for all } n.$$

In particular, the n -th order Alexander module $\mathcal{A}_n^{\mathbb{Z}}(C)$ is a torsion $\mathbb{Z}\Gamma_n$ -module, for all n .

One of the goals of this paper is to provide generalizations of Theorem 6 to various contexts in which the assumption of good behavior at infinity is relaxed.

3. COMPLEX LINE ARRANGEMENTS

Our first result, Theorem 7 below, generalizes Theorem 6 to the context of essential complex line arrangements. In Theorem 8 we study the special class of arrangements containing a line with only nodal singularities.

Assume that all irreducible components of the reduced plane curve C are complex lines, i.e., the defining polynomial $f = \prod_{i=1}^m \ell_i$ factorizes into a product of affine forms $\ell_i : \mathbb{C}^2 \rightarrow \mathbb{C}$, $i = 1, \dots, m$. Let

$$L_i := \ker(\ell_i),$$

and let

$$\mathcal{A} := \{L_1, \dots, L_m\} \subset \mathbb{C}^2$$

be the corresponding complex line arrangement, with complement U . As before, we will use the notation $\delta_n(C)$ for the higher-order degrees of the complement $U := \mathbb{C}^2 \setminus C = \mathbb{C}^2 \setminus \mathcal{A}$.

3.1. Upper bounds on higher-order degrees. In this section, we prove the following generalization of Theorem 6 to the context of essential complex line arrangements.

Theorem 7. *Assume that the complex affine plane curve C defines an essential line arrangement $\mathcal{A} = \{L_1, \dots, L_m\} \subset \mathbb{C}^2$ (that is, not all lines in \mathcal{A} are parallel). Then,*

$$\delta_n(C) \leq m(m-2), \text{ for all } n.$$

Moreover, the equality holds only in the case where \mathcal{A} consists of m lines going through one point, and in that case the equality holds for all $n \geq 0$.

At the opposite spectrum (i.e., if the essentiality assumption is dropped), we have the following:

Proposition 1. *If the plane curve C defines a line arrangement $\mathcal{A} = \{L_1, \dots, L_m\} \subset \mathbb{C}^2$ consisting of m distinct parallel lines, then:*

$$\delta_n(C) = \begin{cases} \infty, & m > 1, \\ 0, & m = 1, \end{cases}$$

for all $n \geq 0$.

Proof. In this case, $\mathbb{C}^2 \setminus C$ is homotopy equivalent to a wedge sum of m circles. If $m = 1$, we have that $\pi_1(\mathbb{C}^2 \setminus C) \cong \mathbb{Z}$, so it is abelian. It then follows from (2) that

$$\delta_n(C) = 0, \text{ for all } n \geq 0.$$

Suppose now that $m > 1$. The chain complex computing $H_*(\mathbb{C}^2 \setminus C; R_n)$ looks like

$$\cdots \rightarrow 0 \rightarrow (R_n)^m \rightarrow R_n \rightarrow 0$$

Hence, $H_1(\mathbb{C}^2 \setminus C; R_n)$ is a non-zero free right R_n -module, so

$$\delta_n(C) = \infty, \text{ for all } n \geq 0.$$

□

Theorem 7 is a consequence of the following two preparatory lemmas (Lemma 1 and Lemma 2). In Lemma 1 we consider the case when there is a line in the arrangement which has no singularities at infinity, whereas in Lemma 2 every line is assumed to have singularities at infinity.

Lemma 1. *In the notations of Theorem 7, assume that there exists a line in \mathcal{A} such that no other line in \mathcal{A} is parallel to it. Then,*

$$\delta_n(C) \leq m(m-2), \text{ for all } n.$$

Moreover, the equality is achieved only in the case when C consists of m lines going through one point, and in that case the equality holds for all $n \geq 0$.

Proof. Reordering, we can assume that L_1 is not parallel to any other line in \mathcal{A} . Let P_1, \dots, P_r be the singular points of C in L_1 . Let

$$F = L_1 \setminus \bigsqcup_{i=1}^r (L_1 \cap \mathbb{B}_i^4)$$

be the (real) surface obtained by removing small balls $\mathbb{B}_i^4 \subset \mathbb{C}^2$ around the singular points P_i . Hence F is obtained from L_1 by removing a 2-dimensional open disk D_i around every singular point P_i .

Let

$$N = F \times S^1.$$

Here N should be thought of as the boundary of a tubular neighborhood around the non-singular part of L_1 . We have that $\partial N = \partial F \times S^1$, and since ∂F is a union of disjoint S^1 's (one from every disk D_i removed), then ∂N is a union of disjoint tori $\bigsqcup_{i=1}^r T_i$ (again, one from every disk D_i removed). Let us fix a point Q_i in the circle S^1 corresponding to the boundary of the disk D_i for every such disk removed.

Let d_i be the number of lines in \mathcal{A} going through the singular point P_i , $i = 1, \dots, r$. Let K_i be the link of the singularity at the point P_i (hence K_i is a Hopf link with d_i components), and let S_i^3 be the boundary of \mathbb{B}_i^4 . We consider the space

$$X = N \cup \left(\bigsqcup_{i=1}^r T_i \right) \left(\bigsqcup_{i=1}^r S_i^3 \setminus K_i \right) \subset \mathbb{C}^2 \setminus C,$$

with the neighborhood N assumed small enough, where the gluing is done as follows: a meridian around the component of K_i corresponding to the line L_1 is glued to $\{Q_i\} \times S^1 \subset N$, and a longitude of the component of K_i corresponding to L_1 is glued to the S^1 corresponding to the boundary of D_i .

The homology of the space X with R_n -coefficients can be computed from the corresponding Mayer-Vietoris sequence:

$$\begin{aligned} & \cdots \rightarrow H_2(N; R_n) \oplus \left(\bigoplus_{i=1}^r H_2(S_i^3 \setminus K_i; R_n) \right) \rightarrow H_2(X; R_n) \xrightarrow{\alpha} \\ (3) \quad & \xrightarrow{\alpha} \bigoplus_{i=1}^r H_1(T_i; R_n) \rightarrow H_1(N; R_n) \oplus \left(\bigoplus_{i=1}^r H_1(S_i^3 \setminus K_i; R_n) \right) \rightarrow H_1(X; R_n) \rightarrow \\ & \rightarrow \bigoplus_{i=1}^r H_0(T_i; R_n) \rightarrow H_0(N; R_n) \oplus \left(\bigoplus_{i=1}^r H_0(S_i^3 \setminus K_i; R_n) \right) \rightarrow H_0(X; R_n) \rightarrow 0 \end{aligned}$$

Hence, using the additivity of the rank of \mathbb{K}_n -modules (recall that \mathbb{K}_n is the right ring of quotients of the Ore domain $\mathbb{Z}\bar{\Gamma}_n$), we have that:

$$\begin{aligned} (4) \quad \text{rk}_{\mathbb{K}_n} H_1(X; R_n) &= \text{rk}_{\mathbb{K}_n} H_1(N; R_n) + \sum_{i=1}^r \text{rk}_{\mathbb{K}_n} H_1(S_i^3 \setminus K_i; R_n) - \sum_{i=1}^r \text{rk}_{\mathbb{K}_n} H_1(T_i; R_n) + \\ &+ \text{rk}_{\mathbb{K}_n} \text{Im}(\alpha) + \sum_{i=1}^r \text{rk}_{\mathbb{K}_n} H_0(T_i; R_n) - \text{rk}_{\mathbb{K}_n} H_0(N; R_n) - \\ &- \sum_{i=1}^r \text{rk}_{\mathbb{K}_n} H_0(S_i^3 \setminus K_i; R_n) + \text{rk}_{\mathbb{K}_n} H_0(X; R_n). \end{aligned}$$

Abusing notation, for any $i = 1, \dots, r$ we denote by $\psi : \pi_1(S_i^3 \setminus K_i) \rightarrow \mathbb{Z}$ the (local) linking number homomorphism induced by $\psi : \pi_1(\mathbb{C}^2 \setminus C) \rightarrow \mathbb{Z}$. Then the infinite cyclic cover of $S_i^3 \setminus K_i$ induced by the homomorphism ψ is homeomorphic to $F_i \times \mathbb{R}$, where F_i is the Milnor fiber corresponding to the singular point P_i . The Γ_n -cover of $S_i^3 \setminus K_i$ factors through this infinite cyclic cover, so we have the following isomorphism of \mathbb{K}_n -modules (e.g., see [8, Section 2.1])

$$(5) \quad H_j(S_i^3 \setminus K_i; R_n) \cong H_j(F_i; \mathbb{K}_n), \quad \text{for all } j \geq 0.$$

The Milnor fiber F_i has the homotopy type of a wedge sum of μ_i circles, where μ_i is the Milnor number associated to the singular point P_i . Together with (5), this yields that $H_2(S_i^3 \setminus K_i; R_n) = 0$. Moreover, since the singularity P_i consists of the intersection of d_i lines, one has

$$\mu_i = (d_i - 1)^2$$

and hence, since the Euler characteristic with coefficients on a 1-dimensional local system over a skew field does not depend on the local system, we have that

$$(6) \quad \begin{aligned} \text{rk}_{\mathbb{K}_n} H_2(S_i^3 \setminus K_i; R_n) &= 0 \\ \text{rk}_{\mathbb{K}_n} H_0(S_i^3 \setminus K_i; R_n) - \text{rk}_{\mathbb{K}_n} H_1(S_i^3 \setminus K_i; R_n) &= \chi(F_i) = 1 - (d_i - 1)^2, \end{aligned}$$

for all $i = 1, \dots, r$.

Similarly, abusing notation again, we denote by $\psi : \pi_1(N) \rightarrow \mathbb{Z}$ the homomorphism induced by the linking number homomorphism $\psi : \pi_1(\mathbb{C}^2 \setminus C) \rightarrow \mathbb{Z}$. Recall that $N = F \times S^1$, and F is homotopy equivalent to a wedge sum of r circles. From this, we see that the infinite cyclic cover of N associated to ψ is homeomorphic to $F \times \mathbb{R}$, so it is homotopy equivalent to F . Since the Γ_n -cover of N factors through this infinite cyclic cover, we get as before that

$$H_j(N; R_n) \cong H_j(F; \mathbb{K}_n), \quad \text{for all } j \geq 0,$$

and hence, we have that

$$(7) \quad \begin{aligned} \text{rk}_{\mathbb{K}_n} H_2(N; R_n) &= 0, \\ \text{rk}_{\mathbb{K}_n} H_0(N; R_n) - \text{rk}_{\mathbb{K}_n} H_1(N; R_n) &= \chi(F) = 1 - r. \end{aligned}$$

Similarly, the Γ_n -cover of the torus T_i factors through the infinite cyclic cover of T_i corresponding to the homomorphism induced by the linking number homomorphism ψ , and this infinite cyclic cover is homeomorphic to $S^1 \times \mathbb{R}$, hence homotopy equivalent to S^1 . Consequently, we have that

$$(8) \quad H_j(T_i; R_n) \cong H_j(S^1; \mathbb{K}_n), \quad \text{for all } j \geq 0, i = 1, \dots, r,$$

and hence

$$(9) \quad \text{rk}_{\mathbb{K}_n} H_0(T_i; R_n) - \text{rk}_{\mathbb{K}_n} H_1(T_i; R_n) = \chi(S^1) = 0$$

for all $i = 1, \dots, r$.

Note that the above calculation (more precisely, the vanishing of the second homology of N and $S_i^3 \setminus K_i$) also implies that the map α in (3) is injective. Thus,

$$\text{rk}_{\mathbb{K}_n} \text{Im}(\alpha) = \text{rk}_{\mathbb{K}_n} H_2(X; R_n).$$

Since X has the homotopy type of a 2-dimensional CW-complex (this can be seen from the way X is constructed), we have that $H_2(X; R_n)$ is a free (right) R_n -module. Thus, $\text{rk}_{\mathbb{K}_n} H_2(X; R_n)$ is either 0 or infinite. But

$$\text{rk}_{\mathbb{K}_n} \text{Im}(\alpha) \leq \sum_{i=1}^r \text{rk}_{\mathbb{K}_n} H_1(T_i; R_n),$$

and the right hand side of this inequality is a finite number (by (8)). Thus,

$$(10) \quad \text{rk}_{\mathbb{K}_n} \text{Im}(\alpha) = 0.$$

Finally, we show that

$$(11) \quad \text{rk}_{\mathbb{K}_n} H_0(X; R_n) = 0$$

by using Fox Calculus (e.g., see [6, Section 6]). Since \mathcal{A} is an essential line arrangement, we have that $m \geq 2$. Let $\gamma_1, \dots, \gamma_m \in \pi_1(X)$ be positively oriented meridians around L_1, \dots, L_m , respectively. We fix a presentation of $\pi_1(X)$ with $\{\gamma_1, \dots, \gamma_m\}$ as the first m generators. The complex of right R_n -modules that computes $H_1(X; R_n)$ using this fixed presentation is

$$\dots \xrightarrow{\partial_2} (R_n)^l \xrightarrow{\partial_1} R_n \xrightarrow{\partial_0} 0,$$

where $l \geq m$, and ∂_1 is given by the row matrix \overline{A} , with

$$A = \begin{pmatrix} \gamma_1 - 1 & \gamma_2 - 1 & \cdots & \gamma_m - 1 & \cdots \end{pmatrix}.$$

Here \overline{A} denotes the matrix obtained from A by taking the involution $\overline{}$ of all of its entries, and the involution in $\mathbb{Z}[\Gamma_n]$ is given by

$$\overline{\sum_{\lambda} n_{\lambda} g_{\lambda}} = \sum_{\lambda} n_{\lambda} g_{\lambda}^{-1}.$$

(The involution is needed here since we are dealing with a complex of right R_n -modules, as opposed to the usual formulation of Fox Calculus, where one works with left modules.) Hence,

$$\overline{A} = \begin{pmatrix} \gamma_1^{-1} - 1 & \gamma_2^{-1} - 1 & \cdots & \gamma_m^{-1} - 1 & \cdots \end{pmatrix}.$$

Let e_1, \dots, e_l be the canonical basis in $(R_n)^l$. We have that

$$\partial_1((e_1 - e_2)\gamma_1) = 1 - \gamma_2^{-1}\gamma_1,$$

which is a unit in R_n for all n , since $\gamma_2^{-1}\gamma_1 \in \overline{\Gamma}_n$ corresponds to a non-zero element in $\Gamma_0 = H_1(\mathbb{C}^2 \setminus C; \mathbb{Z})$. Hence ∂_1 is surjective, so

$$\text{rk}_{\mathbb{K}_n} H_0(X; R_n) = 0,$$

as desired.

Substituting (6), (7), (9), (10) and (11) in equation (4), we get that

$$(12) \quad \text{rk}_{\mathbb{K}_n} H_1(X; R_n) = \sum_{i=1}^r ((d_i - 1)^2 - 1) + r - 1 = \sum_{i=1}^r (d_i - 1)^2 - 1.$$

The next step in our proof is to relate $\text{rk}_{\mathbb{K}_n} H_1(X; R_n)$ to $\delta_n(C)$. Since L_1 is not parallel to any other line in the arrangement \mathcal{A} , the inclusion map $X \hookrightarrow \mathbb{C}^2 \setminus C$ induces an epimorphism

$$\pi_1(X) \longrightarrow \pi_1(\mathbb{C}^2 \setminus C).$$

This can be seen as follows. Let T be a tubular neighborhood of L_1 in \mathbb{C}^2 such that X is a deformation retract of $T \setminus (T \cap C)$. Since L_1 is not parallel to any other line in the arrangement, there exists a generic line L (a line transversal to every other line in the arrangement) such that all of the intersections with lines in the arrangement happen in the interior of T . Thus, the map

$$\pi_1((\mathbb{C}^2 \setminus C) \cap L \cap T) \longrightarrow \pi_1((\mathbb{C}^2 \setminus C) \cap L)$$

induced by inclusion is an epimorphism, as one can see a set of generators of $\pi_1((\mathbb{C}^2 \setminus C) \cap L)$ inside of $(\mathbb{C}^2 \setminus C) \cap L \cap T$. By a Zariski theorem of Lefschetz type ([4, Theorem 6.5, Chapter 1]), we have that the map induced by inclusion

$$\pi_1((\mathbb{C}^2 \setminus C) \cap L) \longrightarrow \pi_1(\mathbb{C}^2 \setminus C)$$

is an epimorphism. Then the following commutative diagram yields that $\pi_1(X) \longrightarrow \pi_1(\mathbb{C}^2 \setminus C)$ is an epimorphism, where all the arrows in the diagram are induced by inclusion maps.

$$\begin{array}{ccc}
\pi_1(X) & & \\
\downarrow & \searrow & \\
\pi_1(T \setminus (T \cap C)) & \longrightarrow & \pi_1(\mathbb{C}^2 \setminus C) \\
\uparrow & & \uparrow \\
\pi_1((\mathbb{C}^2 \setminus C) \cap L \cap T) & \twoheadrightarrow & \pi_1((\mathbb{C}^2 \setminus C) \cap L)
\end{array}$$

We have thus shown that $\pi_1(X) \rightarrow \pi_1(\mathbb{C}^2 \setminus C)$ is an epimorphism. This implies (as in the proof of [11, Theorem 4.1]) that there is an R_n -module epimorphism

$$H_1(X; R_n) \twoheadrightarrow H_1(\mathbb{C}^2 \setminus C; R_n),$$

and hence

$$(13) \quad \delta_n(C) \leq \text{rk}_{\mathbb{K}_n} H_1(X; R_n) = \sum_{i=1}^r (d_i - 1)^2 - 1.$$

Since L_1 is not parallel to any other line in the arrangement, we have that

$$\sum_{i=1}^r (d_i - 1) = m - 1.$$

Furthermore,

$$d_i - 1 \leq m - 1 \quad \text{for all } i = 1, \dots, r,$$

where the equality is only satisfied in the case where \mathcal{A} consists of m lines going through a single point. Altogether,

$$\begin{aligned}
\delta_n(C) &\leq \sum_{i=1}^r (d_i - 1)^2 - 1 \leq (m - 1) \cdot \left(\sum_{i=1}^r (d_i - 1) \right) - 1 \\
&= (m - 1)^2 - 1 = m(m - 2),
\end{aligned}$$

where the second inequality can only be an equality in the case where \mathcal{A} consists of m lines going through a single point. In fact, if \mathcal{A} consists of m lines going through a single point, then X is a deformation retract of $\mathbb{C}^2 \setminus C$, so in that case the first inequality is also an equality (since (13) becomes an equality) and $\delta_n(C) = m(m - 2)$ for all n . (An alternative proof of the fact that $\delta_n(C) = m(m - 2)$ in the case when the arrangement consists of m lines passing through a point was given in [16] by using Fox Calculus.) \square

Remark 4. In concrete examples, one can use (13) to get a better (combinatorial) upper bound for $\delta_n(C)$. Moreover, if there are several lines in \mathcal{A} such that no other line in \mathcal{A} is parallel to them, we can take the tubes around each of those lines to get different bounds for $\delta_n(C)$ similar to (13), and then take the minimum of all of these bounds.

Example 2. Consider the line arrangement of m lines given by $m - 1$ parallel lines L_2, \dots, L_m and a line L_1 transversal to all of them. In this case, the tube X around L_1 is homotopy equivalent to the arrangement complement, so by (12) we have that

$$\delta_n(C) = \text{rk}_{\mathbb{K}_n} H_1(X; R_n) = m - 2$$

for all n .

In view of Lemma 1, the following result completes the proof of Theorem 7.

Lemma 2. *In the notations of Theorem 7, assume that for every line in \mathcal{A} there exists a different line in \mathcal{A} that is parallel to it. Then,*

$$\delta_n(C) \leq (m-2)(m-1) - 1, \text{ for all } n.$$

In particular,

$$\delta_n(C) \leq m(m-2), \text{ for all } n.$$

Proof. Reordering, we can assume that the lines L_1, \dots, L_k are all parallel, with L_j not parallel to L_1 for all $k+1 \leq j \leq m$, and $k \geq 2$.

Let $\overline{L_1}$ be the closure of L_1 in $\mathbb{C}P^2$, and let T be a tubular neighborhood of $\overline{L_1}$ in $\mathbb{C}P^2$ with boundary ∂T , constructed so that $\partial T \setminus (C \cup L_\infty)$ is the space

$$X_\infty = N \cup \left(\left(\bigsqcup_{i=1}^r T_i \right) \sqcup_{T_\infty} \left(\left(\bigsqcup_{i=1}^r S_i^3 \setminus K_i \right) \sqcup (S_\infty^3 \setminus K_\infty) \right) \right) \subset \mathbb{C}^2 \setminus C$$

defined similarly as the space X from the proof of Lemma 1. Here, $L_\infty \subset \mathbb{C}P^2$ is the line at infinity, S_∞^3 is a 3-sphere centered at the intersection point P_∞ of $\overline{L_1}$ with the line at infinity, K_∞ is the link of P_∞ , and T_∞ is the torus along which we glue N to $S_\infty^3 \setminus K_\infty$.

By construction, X_∞ is a deformation retract of $T \setminus (C \cup L_\infty)$, and by a similar argument using a Zariski theorem of Lefschetz type (like in the proof of Lemma 1), we get that the inclusion map $X_\infty \hookrightarrow \mathbb{C}^2 \setminus C$ induces an epimorphism

$$\pi_1(X_\infty) \twoheadrightarrow \pi_1(\mathbb{C}^2 \setminus C),$$

which in turn implies that

$$(14) \quad \delta_n(C) \leq \text{rk}_{\mathbb{K}_n} H_1(X_\infty; R_n).$$

To compute $\text{rk}_{\mathbb{K}_n} H_1(X_\infty; R_n)$, we follow the same steps as in the proof of Lemma 1, based on a Mayer-Vietoris argument. The only difference will appear when computing $H_j(S_\infty^3 \setminus K_\infty; R_n)$ for $j = 0, 1, 2$, since the linking number homomorphism ψ satisfies that $\psi(\gamma_\infty) = -m$, where γ_∞ is a positively oriented meridian around the line at infinity.

Following the same computation as in the proof of (11), we get that

$$(15) \quad H_0(S_\infty^3 \setminus K_\infty; R_n) = 0$$

To compute $\text{rk}_{\mathbb{K}_n} H_2(S_\infty^3 \setminus K_\infty; R_n)$ and $\text{rk}_{\mathbb{K}_n} H_1(S_\infty^3 \setminus K_\infty; R_n)$, we will use Fox Calculus, since we cannot relate these groups to the homology of a Milnor fiber. For this, we first need to find a nice presentation of $\pi_1(S_\infty^3 \setminus K_\infty)$.

By the choices made in the first paragraph of our proof, K_∞ is the Hopf link on $k+1$ components, with $k \geq 2$. A presentation of $\pi_1(S_\infty^3 \setminus K_\infty)$ is given by (e.g., see [13, Lemma 2.7])

$$\pi_1(S_\infty^3 \setminus K_\infty) = \langle \gamma_1, \gamma_2, \dots, \gamma_k, y \mid \gamma_i y \gamma_i^{-1} y^{-1} \text{ for all } i = 1, \dots, k \rangle,$$

where $\gamma_1, \gamma_2, \dots, \gamma_k$ are positively oriented meridians around L_1, \dots, L_k respectively. An equivalent presentation of $\pi_1(S_\infty^3 \setminus K_\infty)$ can be given so that y is the product of meridian loops $\gamma_1, \gamma_2, \dots, \gamma_k, \gamma_\infty$ (in a certain order that is not important here), e.g., see [13, Remark 2.8]. In particular, from this second presentation we get that

$$\psi(y) = k - m.$$

For simplicity, let $a_1 = \gamma_1$, and $a_j = \gamma_j \gamma_1^{-1}$ for all $j = 2, \dots, k$. Then, we get

$$(16) \quad \pi_1(S_\infty^3 \setminus K_\infty) = \langle a_1, a_2, \dots, a_k, y \mid a_i y a_i^{-1} y^{-1} \text{ for all } i = 1, \dots, k \rangle,$$

with

$$\begin{aligned}\psi(a_1) &= 1, \\ \psi(a_j) &= 0, \text{ for all } j = 2, \dots, k, \\ \psi(y) &= k - m.\end{aligned}$$

We compute $H_1(S_\infty^3 \setminus K_\infty; R_n)$ and $H_2(S_\infty^3 \setminus K_\infty; R_n)$ as right R_n -modules, by using the presentation of $\pi_1(S_\infty^3 \setminus K_\infty)$ given in (16). The chain complex computing these groups looks like

$$\dots \rightarrow (R_n)^k \xrightarrow{\partial_2} (R_n)^{k+1} \xrightarrow{\partial_1} R_n \rightarrow 0,$$

where ∂_2 is given by the matrix

$$\begin{aligned}& \begin{pmatrix} 1-y & 0 & \dots & 0 \\ 0 & 1-y & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1-y \\ a_1-1 & a_2-1 & \dots & a_k-1 \end{pmatrix} = \\ & = \begin{pmatrix} 1-y^{-1} & 0 & \dots & 0 \\ 0 & 1-y^{-1} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1-y^{-1} \\ a_1^{-1}-1 & a_2^{-1}-1 & \dots & a_k^{-1}-1 \end{pmatrix}\end{aligned}$$

Note that $1 - a_j^{-1}$ is not zero in $\mathbb{Z}\bar{\Gamma}_n$, since a_j^{-1} is not the identity in Γ_0 , for $j = 2, \dots, k$. Note also that y commutes with a_1, \dots, a_k in $\pi_1(S_\infty^3 \setminus K_\infty)$. Multiply the k -th row by $1 - a_k^{-1}$ on the left (this is a unit in R_n). Add the first row times $1 - a_1^{-1}$, the second row times $1 - a_2^{-1}$, \dots , the $(k-1)$ -st row times $1 - a_{k-1}^{-1}$, and the last row times $1 - y^{-1}$ to the k -th row (all the multiplications are on the left) to get

$$\begin{pmatrix} 1-y^{-1} & 0 & \dots & 0 & 0 \\ 0 & 1-y^{-1} & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 1-y^{-1} & 0 \\ 0 & 0 & \dots & 0 & 0 \\ a_1^{-1}-1 & a_2^{-1}-1 & \dots & a_{k-1}^{-1}-1 & a_k^{-1}-1 \end{pmatrix}$$

Note that $a_k^{-1} - 1$ is a unit in R_n , and we multiply the last column by $(a_k^{-1} - 1)^{-1}$ on the right. Add the last column times $1 - a_j^{-1}$ to the j -th column for all $j = 1, \dots, k-1$. We get

$$\begin{pmatrix} 1-y^{-1} & 0 & \dots & 0 & 0 \\ 0 & 1-y^{-1} & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 1-y^{-1} & 0 \\ 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & \dots & 0 & 1 \end{pmatrix}$$

This matrix corresponds to ∂_2 after a change of basis in both $(R_n)^k$ and $(R_n)^{k+1}$. Therefore, we get $H_2(S_\infty^3 \setminus K_\infty; R_n) = 0$ and

$$H_1(S_\infty^3 \setminus K_\infty, x_0; R_n) = R_n \oplus (R_n / (1 - y^{-1}))^{\oplus(k-1)},$$

where x_0 is a point in $S_\infty^3 \setminus K_\infty$, and there are $k - 1$ direct summands of the form $R_n/(1 - y^{-1})$. By [6, Proposition 5.6], we get that

$$H_1(S_\infty^3 \setminus K_\infty; R_n) = \bigoplus_{k-1 \text{ copies}} R_n/(1 - y^{-1}),$$

so

$$\text{rk}_{\mathbb{K}_n} H_1(S_\infty^3 \setminus K_\infty; R_n) = (k - 1)(m - k).$$

Now, as in the proof of Lemma 1, we get that

$$\begin{aligned} \text{rk}_{\mathbb{K}_n} H_1(X_\infty; R_n) &= \sum_{i=1}^r ((d_i - 1)^2 - 1) + (k - 1)(m - k) + r - 1 \\ (17) \qquad \qquad \qquad &= \sum_{i=1}^r (d_i - 1)^2 + (k - 1)(m - k) - 1. \end{aligned}$$

Since $k \geq 2$, we have that $m - k \leq m - 2$. Also, $d_i \leq m - k + 1$ for all $i = 1, \dots, r$, and $\sum_{i=1}^r (d_i - 1) = m - k$. Thus,

$$\begin{aligned} \sum_{i=1}^r (d_i - 1)^2 + (k - 1)(m - k) - 1 &\leq (m - k) \cdot \left(\sum_{i=1}^r (d_i - 1) + k - 1 \right) - 1 \\ &= (m - k)(m - 1) - 1 \\ &\leq (m - 2)(m - 1) - 1, \end{aligned}$$

which completes the proof. \square

Remark 5. The higher order degrees $\delta_n(C)$ are *not* homeomorphism invariants of $\mathbb{C}^2 \setminus C$. For example, if $m \geq 3$, the case discussed in Example 2 and that of m lines going through a point have homeomorphic complements. However, as we have discussed, their higher order degrees $\delta_n(C)$ are $m - 2$ and $m(m - 2)$, respectively, for all $n \geq 0$. This is due to the dependence of higher order degrees on the local system given by the linking number homomorphism.

3.2. Vanishing of higher-order degrees. In the case when an arrangement contains one line which meets all other lines transversally, higher order degrees are particularly simple. In this section, we prove the following result, which was asserted (without proof) in [12, Section 3.1].

Theorem 8. *Assume that the affine plane curve C defines a line arrangement $\mathcal{A} = \{L_1, \dots, L_m\} \subset \mathbb{C}^2$, which is obtained from an essential line arrangement $\mathcal{A}' = \{L_1, \dots, L_{m-1}\}$ by adjoining a line L_m that is transversal to every line in \mathcal{A}' (that is, the singularities of the curve C along the irreducible component L_m consist of $m - 1$ nodes). Then*

$$\delta_n(C) = 0, \text{ for all } n \geq 0.$$

Before proving the theorem, we recall some notation. Let C' be the curve defined by \mathcal{A}' , let $U = \mathbb{C}^2 \setminus C$ denote as before the complement of \mathcal{A} , and let $U' = \mathbb{C}^2 \setminus C'$ be the complement of \mathcal{A}' . Let $u_0 \in U$, which we will take as the base point for the fundamental groups of both $\pi_1(U)$ and $\pi_1(U')$.

By [15, Lemma 2], we have that

$$(18) \qquad 1 \longrightarrow \mathbb{Z} \xrightarrow{g_1} \pi_1(U) \xrightarrow{g_2} \pi_1(U') \longrightarrow 1$$

is a central extension, where $g_1(1)$ is a positively oriented meridian around L_m , and the map g_2 is induced by inclusion. By the Zariski-Van Kampen theorem (see, e.g., [4, Chapter 4, Section 3]), we find a

presentation of $\pi_1(U)$ of the form

$$(19) \quad \pi_1(U) = \langle y_1, \dots, y_m \mid s_j(y_1, \dots, y_m) \rangle$$

where y_1, \dots, y_{m-1} are certain positively oriented meridians about irreducible components of C' and y_m is a positively oriented meridian about L_m . All of the y_i 's are contained in a generic line section of $\mathbb{C}^2 \setminus C$, and $s_j(y_1, \dots, y_m)$ are certain words on the generators given by braid monodromy. Note that since y_m , and $g_1(1)$ are both positively oriented meridians about L_m , they must be conjugate, and since $g_1(1)$ is in the center of $\pi_1(U)$, then $y_m = g_1(1)$ in $\pi_1(U)$.

Consider the following splitting of g_1

$$\begin{aligned} h : \pi_1(U) &\longrightarrow \mathbb{Z} \\ y_i &\mapsto 0 \quad \text{for } i = 1, \dots, m-1 \\ y_m &\mapsto 1 \end{aligned}$$

which is well defined because it factors through the abelianization of $\pi_1(U)$. Hence, $h(s_j(y_1, \dots, y_m)) = 0$ for all j , which, along with the fact that y_m is in the center of $\pi_1(U)$, allows us to find an equivalent presentation

$$(20) \quad \pi_1(U) = \langle y_1, \dots, y_m \mid [y_i, y_m] \text{ for all } i = 1, \dots, m-1; r_j(y_1, \dots, y_{m-1}) \text{ for } j = 1, \dots, l \rangle$$

where the r_j 's are words in the letters y_1, \dots, y_{m-1} . By setting $x_m = y_m$ and $x_i = y_i y_m^{-1}$ for $i = 1, \dots, m-1$, and taking into account that $\psi(r_j(y_1, \dots, y_{m-1})) = 0$ for all $j = 1, \dots, l$, we obtain the following equivalent presentation for $\pi_1(U)$

$$(21) \quad \pi_1(U) = \langle x_1, \dots, x_m \mid [x_i, x_m] \text{ for all } i = 1, \dots, m-1; r_j(x_1, \dots, x_{m-1}) \text{ for } j = 1, \dots, l \rangle.$$

Using (18) and (20), we get the following presentation for $\pi_1(U')$

$$(22) \quad \pi_1(U') = \langle y_1, \dots, y_{m-1} \mid r_j(y_1, \dots, y_{m-1}) \text{ for } j = 1, \dots, l \rangle.$$

Note that the following map is an isomorphism

$$\begin{aligned} f : \pi_1(U') \times \mathbb{Z} &\longrightarrow \pi_1(U) \\ (y_i, t) &\longmapsto x_i \cdot x_m^t. \end{aligned}$$

For any $n \geq 0$, we denote by $\Gamma_n(U)$ (resp., $\Gamma_n(U')$) the PTFA group corresponding to $\pi_1(U)$ (resp., $\pi_1(U')$), as in Section 2. We then have that f induces an isomorphism

$$f_n : \Gamma_n(U') \times \mathbb{Z} \longrightarrow \Gamma_n(U).$$

Moreover, if $\psi : \pi_1(U) \longrightarrow \mathbb{Z}$ is the linking number homomorphism, then $\bar{\Gamma}_n(U)$ is identified with $\Gamma_n(U')$ via f_n , where

$$\bar{\Gamma}_n(U) = \ker(\bar{\psi} : \Gamma_n(U) \longrightarrow \mathbb{Z}),$$

with $\bar{\psi}$ induced from ψ . As in Section 2, we let $S_n = \mathbb{Z}[\bar{\Gamma}_n(U)] \setminus \{0\}$, $\mathbb{K}_n = \mathbb{Z}[\bar{\Gamma}_n(U)] S_n^{-1}$, and $R_n = \mathbb{Z}[\Gamma_n(U)] S_n^{-1}$. Let $\mathcal{K}_n(U')$ denote the (skew) field of quotients of the Ore domain $\mathbb{Z}[\Gamma_n(U')]$.

Remark 6. Notice that f_n identifies \mathbb{K}_n with $\mathcal{K}_n(U')$.

Consider the matrix of Fox derivatives for $\pi_1(U')$, that is,

$$\left(\frac{\partial r_j(y_1, \dots, y_{m-1})}{\partial y_i} \right)_{i,j}, \quad 1 \leq i \leq m-1, 1 \leq j \leq l,$$

which has entries in $\mathbb{Z}[\pi_1(U')]$, and we take its involution

$$A = \overline{\left(\frac{\partial r_j(y_1, \dots, y_{m-1})}{\partial y_i} \right)_{i,j}}.$$

Let $q'_n : \pi_1(U') \rightarrow \Gamma_n(U')$ be the projection, and let

$$B(n) = A^{q'_n},$$

that is, the matrix formed by the images of the entries of A by q'_n . Since $\mathcal{K}_n(U')$ is flat over $\mathbb{Z}[\Gamma_n(U')]$, we have that $B(n)$ is a presentation matrix for the right $\mathcal{K}_n(U')$ -module $H_1(U', u_0; \mathcal{K}_n(U'))$; again, we refer to [6, Section 6] for more details about Fox calculus.

Lemma 3. *The rank of the left $\mathcal{K}_n(U')$ -module generated by the rows of $B(n)$ is $m - 2$.*

Proof. Since U' is the complement of an essential line arrangement, we get by Theorem 7 that $\delta_n(C')$ is finite. By [11, Remark 3.8], this means that

$$\mathrm{rk}_{\mathcal{K}_n(U')} H_1(U'; \mathcal{K}_n(U')) = 0,$$

and by [6, Proposition 5.6], we get that

$$\mathrm{rk}_{\mathcal{K}_n(U')} H_1(U', u_0; \mathcal{K}_n(U')) = \mathrm{rk}_{\mathcal{K}_n(U')} H_1(U'; \mathcal{K}_n(U')) + 1 = 1.$$

Since $B(n)$ is an $(m - 1) \times l$ matrix, the rank of the left $\mathcal{K}_n(U')$ -module generated by the rows of $B(n)$ (which is the same as the rank of the right $\mathcal{K}_n(U')$ -module generated by the columns of $B(n)$) must be $m - 2$. \square

We are now ready to prove Theorem 8.

Proof of Theorem 8. Let $n \geq 0$. We start by considering the presentation matrix for $H_1(U, u_0; R_n)$ as a right R_n -module given by the involution of the matrix of Fox derivatives corresponding to the presentation of $\pi_1(U)$ from (21), which is

$$(23) \quad \left(\begin{array}{cccc|c} 1 - x_m^{-1} & 0 & \cdots & 0 & B(n) \\ 0 & 1 - x_m^{-1} & \cdots & 0 & \\ \vdots & \vdots & \ddots & \vdots & \\ 0 & 0 & \cdots & 1 - x_m^{-1} & \\ \hline x_1^{-1} - 1 & x_2^{-1} - 1 & \cdots & x_{m-1}^{-1} - 1 & 0 \quad \cdots \quad 0 \end{array} \right)$$

where $B(n)$ is seen as a matrix in $\mathbb{K}_n \subset R_n$ by the identification of \mathbb{K}_n and $\mathcal{K}_n(U')$ given by f_n , and the rest of the entries are seen in R_n . By Lemma 3, the rank of the left \mathbb{K}_n -module spanned by the rows of $B(n)$ is $m - 2$. We denote this by $\mathrm{rk}_{\mathbb{K}_n} B(n) = m - 2$.

Note that $(1 - x_j^{-1})$ are non-zero elements of S_n for all $j = 1, \dots, m - 1$. We multiply the first row (on the left) by $(1 - x_1^{-1})$, and then add the j -th row times $(1 - x_j^{-1})$ to the first row for all $j = 2, \dots, m$. Taking into account that x_m commutes with everything else, we get

$$\left(\begin{array}{cccc|c} 0 & 0 & \cdots & 0 & B_1(n) \\ 0 & 1 - x_m^{-1} & \cdots & 0 & \\ \vdots & \vdots & \ddots & \vdots & \\ 0 & 0 & \cdots & 1 - x_m^{-1} & \\ \hline x_1^{-1} - 1 & x_2^{-1} - 1 & \cdots & x_{m-1}^{-1} - 1 & 0 \quad \cdots \quad 0 \end{array} \right)$$

where $\text{rk}_{\mathbb{K}_n} B_1(n) = m - 2$, since we just did row operations in \mathbb{K}_n to get from $B(n)$ to $B_1(n)$. Multiplying the first column (on the right) by $(x_1^{-1} - 1)^{-1}$ and then doing column operations, we get

$$(24) \quad \left(\begin{array}{cccc|cccc} 0 & 0 & \cdots & 0 & & & & \\ 0 & 1 - x_m^{-1} & \cdots & 0 & & & & \\ \vdots & \vdots & \ddots & \vdots & & & & \\ 0 & 0 & \cdots & 1 - x_m^{-1} & & & & \\ \hline 1 & 0 & \cdots & 0 & 0 & \cdots & 0 & \end{array} \right) B_1(n)$$

Performing row and column operations in \mathbb{K}_n , and using that $\text{rk}_{\mathbb{K}_n} B_1(n) = m - 2$, we get a matrix of the form

$$\left(\begin{array}{cccc|cccc} 0 & * & \cdots & * & 0 & \cdots & 0 & 0 & \cdots & 0 \\ 0 & * & \cdots & * & & & & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & & & & \vdots & \ddots & \vdots \\ 0 & * & \cdots & * & & & & 0 & \cdots & 0 \\ \hline 1 & 0 & \cdots & 0 & 0 & \cdots & 0 & 0 & \cdots & 0 \end{array} \right) I_{m-2}$$

where I_{m-2} is the identity matrix of dimension $m - 2$. Performing column operations we can get this matrix to look like

$$\left(\begin{array}{cccc|cccc} 0 & * & \cdots & * & 0 & \cdots & 0 & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 & & & & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & & & & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 & & & & 0 & \cdots & 0 \\ \hline 1 & 0 & \cdots & 0 & 0 & \cdots & 0 & 0 & \cdots & 0 \end{array} \right) I_{m-2}$$

Permuting the first and last rows, and putting the columns corresponding to I_{m-2} as columns 2, 3, \dots , $m - 1$, we get

$$\left(\begin{array}{cccc|cccc} & & & & 0 & \cdots & 0 & 0 & \cdots & 0 \\ & & & & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ & & & & 0 & \cdots & 0 & 0 & \cdots & 0 \\ & & & & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ & & & & 0 & \cdots & 0 & 0 & \cdots & 0 \\ \hline & & & & 0 & \cdots & 0 & * & \cdots & * \end{array} \right) I_{m-1}$$

Let \mathcal{K}_n be the skew field of quotients of $\mathbb{Z}[\Gamma_n(U)]$. By a similar argument as in the proof of Lemma 3, the rank of the left \mathcal{K}_n -module spanned by the rows of this matrix must be $m - 1$, so the last row is actually identically 0. Hence, $\delta_n(C) = 0$. \square

4. PLANE CURVES

In this section, we adapt some of the results of Section 3 to the context of affine plane curve complements.

4.1. Upper bounds on higher-order degrees. The goal of this section is to prove the following generalization of Theorem 6, in which the plane curve C is allowed to have *mild* singularities at infinity.

Theorem 9. *Let $C \subset \mathbb{C}^2$ be a reduced plane curve of degree m , let \overline{C} be its closure in $\mathbb{C}P^2$ and let L_∞ be the line at infinity. Suppose that the intersections of L_∞ and \overline{C} are either transversal or L_∞ is the tangent line to \overline{C} at a smooth point and it is a simple tangent there (i.e., it has multiplicity 2). If any of the following two conditions hold*

- (a) $m = 2$;
- (b) at least one of the intersections of \overline{C} and L_∞ is transversal,

then

$$\delta_n(C) \leq m(m - 2),$$

for all $n \geq 0$.

Proof. Assume that the intersection $\overline{C} \cap L_\infty$ consists of r distinct points. The case $r = m$ corresponds to the curve C being in general position at infinity, which was already considered in Theorem 6. So, without any loss of generality, we may assume that $r \leq m - 1$. The proof of the theorem in this case relies on a Mayer-Vietoris argument similar to the one we used in the proofs of Lemma 1 and Lemma 2.

Let T be a tube in $\mathbb{C}P^2$ around L_∞ , and let $\{P_1, \dots, P_r\} = \overline{C} \cap L_\infty$. If $m > 2$, we can assume (after reordering) that the intersection of \overline{C} and L_∞ is transversal at P_{r-1} . Then $T \setminus (\overline{C} \cup L_\infty)$ deformation retracts to the space

$$X = \partial T \setminus C = N \cup \left(\bigsqcup_{i=1}^r T_i \right) \left(\bigsqcup_{i=1}^r S_i^3 \setminus K_i \right) \subset \mathbb{C}^2 \setminus C$$

where, as in the proof of Lemma 1, N is the boundary of a tube around the non-singular part of L_∞ , that is, a tube around L_∞ minus a disk around every point P_i , $S_i^3 \setminus K_i$ is the link complement of the singularity of $\overline{C} \cup L_\infty$ at P_i , and T_i is a 2-torus described as in the proof of Lemma 1. Note that X is the link (complement) at infinity, which is the space used in [11] for proving Theorem 6, with the difference that if C is in general position at infinity, X is just the complement of the Hopf link on m components.

By a Zariski-Lefschetz type theorem again, the inclusion $X \hookrightarrow \mathbb{C}^2 \setminus C$ induces an epimorphism

$$\pi_1(X) \twoheadrightarrow \pi_1(\mathbb{C}^2 \setminus C),$$

which in turn implies that

$$\delta_n(C) \leq \text{rk}_{\mathbb{K}_n} H_1(X; R_n).$$

It thus suffices to show that $\text{rk}_{\mathbb{K}_n} H_1(X; R_n) \leq m(m - 2)$.

The Mayer-Vietoris sequence for the homology of X with R_n -coefficients yields the same equality as in formula (4) of Lemma 1, so it remains to compute (or bound) all of the terms on the right-hand side of (4).

We begin by noticing that N is homotopy equivalent to the cartesian product of a wedge sum of $r - 1$ circles and S^1 if $r > 1$, and to S^1 if $r = 1$ (and $m = 2$). If $r = 1$, a direct Fox Calculus computation yields that

$$(25) \quad \begin{aligned} H_2(N; R_n) &= 0, \\ H_1(N; R_n) &= 0, \\ \text{rk}_{\mathbb{K}_n} H_0(N; R_n) &= m, \end{aligned}$$

where one only uses the fact that a positively oriented meridian γ_∞ around L_∞ generates $\pi_1(N) \cong \mathbb{Z}$ and $\psi(\gamma_\infty) = -m$.

If $r > 1$, a presentation for the fundamental group of $N \simeq \left(\bigvee_{r-1} S^1 \right) \times S^1$ is given as:

$$\pi_1(N) = \langle a_1, \dots, a_{r-1}, b \mid [a_i, b] \text{ for } i = 1, \dots, r - 1 \rangle,$$

where each a_i corresponds to a circle in the wedge sum, which in turn corresponds to the boundary of a disk centered at P_i , while b is a positively oriented meridian about L_∞ . In particular, since the intersection of \overline{C} and L_∞ is transversal at P_{r-1} , the loop a_{r-1} can be chosen to be an oriented meridian about the irreducible component of \overline{C} going through P_{r-1} . Hence, $\psi(a_{r-1}) = 1$, where ψ denotes as before the

linking number homomorphism. Setting $x_{r-1} = a_{r-1}$ and $x_j = a_j a_{r-1}^{-\psi(a_j)}$ for all $j = 1, \dots, r-2$, we get the equivalent presentation

$$\pi_1(N) = \langle x_1, \dots, x_{r-1}, b \mid [x_i, b] \text{ for } i = 1, \dots, r-1 \rangle.$$

The involution of the matrix of Fox derivatives looks like the left-hand side of the matrix in equation (23) of Section 3.2, after changing m for r and x_m for b .

If $r > 2$, we have two possible cases: either there exists $j \in \{1, \dots, r-2\}$ such that $x_j \neq 0$ in Γ_n (in which case, by reordering, we can assume that $j = 1$), or $x_j = 0$ in Γ_n for all $j = 1, \dots, r-2$.

In the first case, the same computations as in Section 3.2 yield the left-hand side of the matrix in equation (24) of Section 3.2. Using that $\psi(b) = -m$, we get that

$$\begin{aligned} H_2(N; R_n) &= 0, \\ \text{rk}_{\mathbb{K}_n} H_1(N; R_n) &= m(r-2). \end{aligned}$$

We can also see by using Fox Calculus and the fact that $x_1^{-1} - 1$ is a unit in \mathbb{K}_n that $\text{rk}_{\mathbb{K}_n} H_0(N; R_n) = 0$, so we get that

$$(26) \quad \begin{aligned} H_2(N; R_n) &= 0, \\ \text{rk}_{\mathbb{K}_n} H_1(N; R_n) - \text{rk}_{\mathbb{K}_n} H_0(N; R_n) &= m(r-2). \end{aligned}$$

In fact, these equalities also hold for the case $r = 1$ considered in (25).

If $r > 2$ and $x_j = 0$ in Γ_n for all $j = 1, \dots, r-2$, the complex that computes $H_*(N; R_n)$ by Fox Calculus looks like

$$\longrightarrow (R_n)^{r-1} \xrightarrow{\partial_2} (R_n)^r \xrightarrow{\partial_1} R_n \longrightarrow 0$$

where ∂_2 is given by the matrix

$$\begin{pmatrix} 1 - b^{-1} & 0 & 0 & \cdots & 0 \\ 0 & 1 - b^{-1} & 0 & \cdots & 0 \\ 0 & 0 & 1 - b^{-1} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 - b^{-1} \\ 0 & 0 & \cdots & 0 & x_{r-1}^{-1} - 1 \end{pmatrix}$$

and ∂_1 by

$$\begin{pmatrix} 0 & 0 & \cdots & 0 & x_{r-1}^{-1} - 1 & b^{-1} - 1 \end{pmatrix},$$

and we can see directly that (26) also holds in this case.

Finally, let us analyze the case $r = 2$. In this case, N is homotopy equivalent to a torus, and the Γ_n -cover of N factors through the infinite cyclic cover of N corresponding to ψ , which is homotopy equivalent to S^1 (a similar argument was used in the proof of Lemma 1). Thus, in this case,

$$\begin{aligned} \text{rk}_{\mathbb{K}_n} H_1(N; R_n) - \text{rk}_{\mathbb{K}_n} H_0(N; R_n) &= \text{rk}_{\mathbb{K}_n} H_1(S^1; \mathbb{K}_n) - \text{rk}_{\mathbb{K}_n} H_0(S^1; \mathbb{K}_n) \\ &= \chi(S^1) \\ &= 0, \end{aligned}$$

so the equalities in (26) also hold.

Let us next compute the local contributions in (4), i.e., corresponding to the link complements of the P_i 's. Suppose that the intersection of \overline{C} and L_∞ is transversal at P_i . Then the link K_i of P_i is the Hopf link on 2 components, and we can pick meridians a, b around the components of the link to generate

$\pi_1(S_i^3 \setminus K_i) \cong \mathbb{Z}^2$, and which satisfy $\psi(a) = 1$ and $\psi(b) = -m$. In this case, $S_i^3 \setminus K_i$ deformation retracts onto a torus, so using again an argument involving the infinite cyclic cover, we get that

$$(27) \quad \text{rk}_{\mathbb{K}_n} H_1(S_i^3 \setminus K_i; R_n) = \text{rk}_{\mathbb{K}_n} H_0(S_i^3 \setminus K_i; R_n),$$

where both ranks are finite, and $H_2(S_i^3 \setminus K_i; R_n) = 0$.

On the other hand, if L_∞ is the tangent line to \overline{C} at P_i , with multiplicity 2, then K_i is a type (2, 4) torus link, which corresponds to the following braid



Again, we can find a presentation for $\pi_1(S_i^3 \setminus K_i)$ with two generators and one relation, namely

$$\pi_1(S_i^3 \setminus K_i) = \langle a, b \mid ababa^{-1}b^{-1}a^{-1}b^{-1} \rangle,$$

where a corresponds to a positively oriented meridian around the irreducible component of \overline{C} going through P_i and b corresponds to a positively oriented meridian around L_∞ . We then have that $\psi(a) = 1$ and $\psi(b) = -m$. By setting $x = a$ and $y = ba^m$, we get the equivalent presentation

$$\pi_1(S_i^3 \setminus K_i) = \langle x, y \mid xyx^{-m+1}yx^{-1}y^{-1}x^{m-1}y^{-1} \rangle.$$

Let $s = xyx^{-m+1}yx^{-1}y^{-1}x^{m-1}y^{-1}$. When regarded in $\mathbb{K}_n[t^{\pm 1}]$, $\frac{\partial s}{\partial x}$ is a polynomial of degree at most $m - 1$, and $\frac{\partial s}{\partial x} = 0$ if and only if $y = 1$ in $\mathbb{K}_n[t^{\pm 1}]$. Suppose that $y \neq 1$ in $\mathbb{K}_n[t^{\pm 1}]$. Since there is only one relation, a Fox Calculus computation yields that

$$\text{rk}_{\mathbb{K}_n} H_1(S_i^3 \setminus K_i; R_n) \leq m - 1,$$

and

$$\text{rk}_{\mathbb{K}_n} H_0(S_i^3 \setminus K_i; R_n) = 0.$$

Hence, if $y \neq 1$ in $\mathbb{K}_n[t^{\pm 1}]$, then

$$(28) \quad \text{rk}_{\mathbb{K}_n} H_1(S_i^3 \setminus K_i; R_n) - \text{rk}_{\mathbb{K}_n} H_0(S_i^3 \setminus K_i; R_n) \leq m - 1.$$

If $y = 1$ in $\mathbb{K}_n[t^{\pm 1}]$, we note that $\frac{\partial s}{\partial y}$ is a polynomial of degree m , so $\text{rk}_{\mathbb{K}_n} H_1(S_i^3 \setminus K_i; R_n) = m$. Moreover, if $\frac{\partial s}{\partial x} = 0$, then a Fox Calculus computation yields that $\text{rk}_{\mathbb{K}_n} H_0(S_i^3 \setminus K_i; R_n) = 1$. Hence, if $y = 1$ in $\mathbb{K}_n[t^{\pm 1}]$, the same inequality as in (28) holds. In both cases ($y = 1$ and $y \neq 1$ in $\mathbb{K}_n[t^{\pm 1}]$), we see that $H_2(S_i^3 \setminus K_i; R_n) = 0$.

Using a presentation of $\pi_1(X)$ in which there is a generator a such that $\psi(a) = 1$ (for example, taking a to be a positively oriented meridian around the irreducible component of \overline{C} going through P_i), we also get that

$$(29) \quad \text{rk}_{\mathbb{K}_n} H_0(X; R_n) \leq 1$$

via a Fox Calculus computation.

Next, we deal with the contributions to formula (4) of the tori T_i , for any $i = 1, \dots, r$. We have the following presentation of the fundamental group

$$\pi_1(T_i) = \langle c, b \mid [c, b] \rangle,$$

where b is a positively oriented meridian about L_∞ , $\psi(c) = 1$ if the intersection of \overline{C} with L_∞ is transversal at P_i , and $\psi(c) = 2$ otherwise.

If the intersection of \overline{C} with L_∞ is transversal at P_i , then the Γ_n -cover of T_i factors through the infinite cyclic cover corresponding to ψ . Moreover, since $\psi(c) = 1$, this infinite cyclic cover is homeomorphic to $S^1 \times \mathbb{R}$, thus homotopy equivalent to S^1 . Therefore,

$$(30) \quad \begin{aligned} \operatorname{rk}_{\mathbb{K}_n} H_0(T_i; R_n) - \operatorname{rk}_{\mathbb{K}_n} H_1(T_i; R_n) &= \operatorname{rk}_{\mathbb{K}_n} H_0(S^1; \mathbb{K}_n) - \operatorname{rk}_{\mathbb{K}_n} H_1(S^1; \mathbb{K}_n) = \chi(S^1) = 0, \\ \operatorname{rk}_{\mathbb{K}_n} H_2(T_i; R_n) &= \operatorname{rk}_{\mathbb{K}_n} H_2(S^1; \mathbb{K}_n) = 0. \end{aligned}$$

If L_∞ is tangent to \overline{C} at P_i (note that there are $m - r$ such P_i 's), then we have as before that $\psi(c) = 2$. There is a \mathbb{K}_n -module isomorphism (e.g., see [8, Section 2.1])

$$H_*(T_i; R_n) \cong H_*(\tilde{T}_i; \mathbb{K}_n),$$

where \tilde{T}_i is the (possibly disconnected) infinite cyclic cover of T_i corresponding to ψ . Note that \tilde{T}_i is either homeomorphic to $S^1 \times \mathbb{R}$, or to the disjoint union $(S^1 \times \mathbb{R}) \sqcup (S^1 \times \mathbb{R})$, depending on whether m is odd or even, respectively. In both cases, \tilde{T}_i is homotopy equivalent to a one-dimensional finite CW-complex with vanishing Euler characteristic. Hence

$$(31) \quad \begin{aligned} \operatorname{rk}_{\mathbb{K}_n} H_0(T_i; R_n) - \operatorname{rk}_{\mathbb{K}_n} H_1(T_i; R_n) &= \operatorname{rk}_{\mathbb{K}_n} H_0(\tilde{T}_i; \mathbb{K}_n) - \operatorname{rk}_{\mathbb{K}_n} H_1(\tilde{T}_i; \mathbb{K}_n) = \chi(\tilde{T}_i) = 0, \\ \operatorname{rk}_{\mathbb{K}_n} H_2(T_i; R_n) &= \operatorname{rk}_{\mathbb{K}_n} H_2(\tilde{T}_i; \mathbb{K}_n) = 0. \end{aligned}$$

Arguing as in the proof of Lemma 1, we also get that

$$(32) \quad \operatorname{rk}_{\mathbb{K}_n} \operatorname{Im}(\alpha) = 0.$$

Altogether, substituting (26), (27), (28), (29), (30), (31) and (32) into (4), we get that

$$\operatorname{rk}_{\mathbb{K}_n} H_1(X; R_n) \leq m(r - 2) + (m - r)(m - 1) + 1 = m^2 - 3m + r + 1.$$

Moreover, since we assumed that $r \leq m - 1$, we get that

$$\operatorname{rk}_{\mathbb{K}_n} H_1(X; R_n) \leq m^2 - 2m = m(m - 2),$$

thus concluding the proof. \square

4.2. Vanishing of higher-order degrees. In the case when an irreducible component of a plane curve C is a line L which meets all other components of $C \cup L_\infty$ transversally (with L_∞ denoting the line at infinity in $\mathbb{C}P^2$), higher order degrees are particularly simple. This is exemplified in the following generalization of Theorem 8.

Theorem 10. *Let $n \geq 0$. Assume that the affine plane curve C is of the form $C = L \cup C'$, where C' is a curve of degree $m - 1$ in \mathbb{C}^2 such that $\delta_n(C')$ is finite, and L is a line transversal to C' , such that $L \cap C'$ consists of $m - 1$ distinct points. Then,*

$$\delta_n(C) = 0.$$

Proof. The proof follows the same steps as in the proof of Theorem 8. Let $U = \mathbb{C}^2 \setminus C$ and $U' = \mathbb{C}^2 \setminus C'$. The Zariski-Van Kampen theorem (see, e.g., [4, Chapter 4, Section 3]) can be used to find a presentation of $\pi_1(U)$ such as the one described in (19). Using [15, Lemma 2] and the same arguments as in the proof of Theorem 8, we find presentations of $\pi_1(U)$ and $\pi_1(U')$ such as the ones described in (20) and (22), respectively. Thus, Remark 6 still holds in this setting. The rest of the proof follows the same Fox Calculus computation as in the proof of Theorem 8, except for the proof of Lemma 3, which in this case follows from the hypothesis that $\delta_n(C')$ is finite. \square

Remark 7. The fact, used in the proof of Theorem 10, that $\pi_1(\mathbb{C}^2 \setminus C) \cong \pi_1(\mathbb{C}^2 \setminus C') \times \mathbb{Z}$, can also be deduced from the Oka-Sakamoto theorem [14]. Indeed, under our assumptions, we have by [14] that

$$\pi_1(\mathbb{C}^2 \setminus C) \cong \pi_1(\mathbb{C}^2 \setminus C') \times \pi_1(\mathbb{C}^2 \setminus L) \cong \pi_1(\mathbb{C}^2 \setminus C') \times \mathbb{Z}.$$

In particular, this also shows that Theorem 10 can be generalized as follows:

Theorem 11. *Let $n \geq 0$. Assume that the affine plane curve C is of the form $C = C' \cup C''$, where C' is a curve of degree m' in \mathbb{C}^2 such that $\delta_n(C')$ is finite, and C'' is a smooth curve of degree m'' , such that $C' \cap C''$ consists of $m'm''$ distinct points in \mathbb{C}^2 . Then,*

$$\delta_n(C) = 0.$$

5. RELATIONSHIP WITH THE FIRST CHARACTERISTIC VARIETY

In this section, we relate the higher-order degrees to more classical Alexander-type invariants. We begin by recalling the following result:

Proposition 2. ([11, Proposition 5.1]) *If C is an irreducible affine plane curve, then*

$$\delta_0(C) = \deg \Delta_C(t),$$

where $\Delta_C(t)$ denotes the Alexander polynomial of the curve complement. If, moreover, the Alexander polynomial is trivial, then all higher-order degrees vanish.

In what follows, we generalize the above result to non-irreducible affine plane curves. Let us first introduce some notation, following the conventions from [17].

Let R be a Noetherian commutative ring with unit. Assume also that R is a unique factorization domain. Let M be a finitely generated R -module. Then M admits a finite presentation of the form

$$R^q \xrightarrow{\Phi} R^m \longrightarrow M.$$

Definition 4. The i -th elementary ideal of M , denoted $E_i(M)$, is the ideal of R generated by the minors of size $m - i$ of the $m \times q$ matrix Φ , with the convention that $E_i(M) = R$ if $i \geq m$ and $E_i(M) = 0$ if $m - i > q$.

Remark 8. The i -th elementary ideal (also referred to as the $(i + 1)$ -st Fitting ideal) does not depend on the choice of representation of M as an R -module.

Remark 9. It follows immediately from Definition 4 that

$$E_i(M) \subset E_{i+1}(M)$$

for all $i \geq 0$.

Definition 5. Let $i \geq 0$. We define $\Delta_i(M) \in R$ to be the generator of the smallest principal ideal in R containing $E_i(M)$, that is, the greatest common divisor of all elements of $E_i(M)$.

Remark 10. $\Delta_i(M)$ is well-defined up to multiplication by a unit of R .

Let $G = \langle x_1, \dots, x_m \mid r_1, \dots, r_q \rangle$ be a finitely presented group, let H be its maximal, torsion free abelian quotient, and let $\pi : G \longrightarrow H$ be the quotient map. The group ring $\mathbb{Z}H$ is a commutative Noetherian ring with unit, which is also a unique factorization domain.

Let F_m be the free group with generators x_1, \dots, x_m . For each $1 \leq j \leq m$, there is a linear operator

$$\frac{\partial}{\partial x_j} : \mathbb{Z}F_m \longrightarrow \mathbb{Z}F_m$$

(called the j -th Fox derivative) uniquely determined by the following properties:

- (a) $\frac{\partial 1}{\partial x_j} = 0$,
- (b) $\frac{\partial x_i}{\partial x_j} = \delta_{ij}$,
- (c) $\frac{\partial uv}{\partial x_j} = \frac{\partial u}{\partial x_j} + u \frac{\partial v}{\partial x_j}$, for any $u, v \in F_m$.

Let $\phi : F_m \rightarrow G$ be the presenting homomorphism.

Definition 6. The *Alexander matrix* of the given presentation of G is

$$\Phi_G = \left(\frac{\partial r_i}{\partial x_j} \right)^{\pi \circ \phi} : (\mathbb{Z}H)^q \rightarrow (\mathbb{Z}H)^m.$$

Now, let X be a connected CW-complex with a unique 0-cell x_0 , and finitely many 1-cells. Let $G = \pi_1(X, x_0)$ be the fundamental group, and let H be its maximal torsion-free abelian quotient, that is, $H \cong \mathbb{Z}^{b_1(G)}$. The canonical projection $\pi : G \rightarrow H$ defines a local system of $\mathbb{Z}H$ -modules. The long exact sequence for the homology of the pair (X, x_0) with coefficients in $\mathbb{Z}H$ given by this local system is

$$\dots \rightarrow 0 \rightarrow H_1(X; \mathbb{Z}H) \rightarrow H_1(X, x_0; \mathbb{Z}H) \rightarrow H_0(x_0; \mathbb{Z}H) \rightarrow H_0(X; \mathbb{Z}H) \rightarrow 0,$$

where

$$\ker(H_0(x_0; \mathbb{Z}H) = \mathbb{Z}H \rightarrow H_0(X; \mathbb{Z}H))$$

can be identified with the augmentation ideal $I_H = \ker(\epsilon : \mathbb{Z}H \rightarrow \mathbb{Z})$, and ϵ is defined as

$$\begin{aligned} \epsilon : \mathbb{Z}H &\rightarrow \mathbb{Z} \\ \sum n_i h_i &\mapsto \sum n_i \end{aligned}$$

for $n_i \in \mathbb{Z}$, $h_i \in H$.

The $\mathbb{Z}H$ -modules $H_1(X; \mathbb{Z}H)$ and $H_1(X, x_0; \mathbb{Z}H)$ depend only on the fundamental group G , so we denote them by B_G and A_G , respectively. From the above discussion, these modules fit into the following short exact sequence of $\mathbb{Z}H$ -modules

$$0 \rightarrow B_G \rightarrow A_G \rightarrow I_H \rightarrow 0.$$

Definition 7. The $\mathbb{Z}H$ -module B_G is called the *Alexander invariant* of X , and A_G is called the *Alexander module* of X .

Definition 8. The *Alexander polynomial* of the group G , denoted by Δ_G , is defined by

$$\Delta_G = \Delta_1(A_G) = \gcd(E_1(A_G)) \in \mathbb{Z}H.$$

Remark 11. Sometimes, the Alexander polynomial of the fundamental group of a CW-complex X is defined as $\Delta_0(B_G) = \gcd(E_0(B_G))$, but the definition using the Alexander module instead of the Alexander invariant is more suitable for our purposes. In the case when $b_1(X) = 1$ (e.g., $X = \mathbb{C}^2 \setminus C$, where C is an irreducible affine plane curve), the two definitions coincide. Besides, the Alexander matrix Φ_G provides a presentation for A_G , so we can compute Δ_G from it.

Definition 9. Let X be a connected CW-complex with finite k -skeleton, and let $G = \pi_1(X)$. The *homology jump loci* of X (over \mathbb{C}) are the Zariski closed sets

$$\mathcal{V}_d^i(X) = \{\rho \in \text{Hom}(G, \mathbb{C}^*) \mid \dim_{\mathbb{C}} H_i(X; \mathbb{C}_\rho) \geq d\},$$

where \mathbb{C}_ρ is the rank-one \mathbb{C} -local system on X induced by ρ , $0 \leq i \leq k$, and $d > 0$. When $i = 1$, we use the simplified notation $\mathcal{V}_d(X)$ for $\mathcal{V}_d^1(X)$.

Let X be a connected CW complex with a unique 0-cell and finitely many 1-cells. Let $\text{Hom}(G, \mathbb{C}^*)^0$ be the identity component of the algebraic group $\text{Hom}(G, \mathbb{C}^*)$. The projection map $\pi : G \rightarrow H$ induces an isomorphism $\pi^* : \text{Hom}(H, \mathbb{C}^*) \rightarrow \text{Hom}(G, \mathbb{C}^*)^0$.

Definition 10. The *characteristic varieties* of X (over \mathbb{C}), denoted by $V_d(X)$, are the subvarieties of $(\mathbb{C}^*)^{b_1(X)} \cong \text{Spec } \mathbb{C}H$ given by

$$V_d(X) = V(E_{d-1}(B_G \otimes \mathbb{C})).$$

We recall the following result from [7], see also [17, Proposition 4.7].

Proposition 3. Let $\rho : H \rightarrow \mathbb{C}^*$ be a non-trivial character. Then, for all $d \geq 1$,

$$\pi^*(\rho) \in \mathcal{V}_d(X) \iff \rho \in V(E_d(A_G \otimes \mathbb{C})) \iff \rho \in V_d(X).$$

Remark 12. If $X = \mathbb{C}^2 \setminus C$ is a plane curve complement, then π^* is the identity, and $\text{Hom}(G, \mathbb{C}^*) \cong \text{Hom}(H, \mathbb{C}^*)$ can be identified with $(\mathbb{C}^*)^m$, where m is the number of irreducible components of the plane curve. In this case, Proposition 3 asserts that, away from $(1, \dots, 1)$, \mathcal{V}_d and V_d coincide. Moreover, by Remark 11, away from $(1, \dots, 1)$, we can compute V_d from the dimension $m - d$ minors of the Alexander matrix $\bar{\Phi}_G$.

From now on, let C be a plane curve in \mathbb{C}^2 with m irreducible components, with $m \geq 2$, and let $G = \pi_1(U, u_0)$, where $U = \mathbb{C}^2 \setminus C$. We denote by $\Delta_C \in \mathbb{Z}[t_1^{\pm 1}, \dots, t_m^{\pm 1}]$ the Alexander polynomial Δ_G of Definition 8.

Definition 11. Let q be a Laurent polynomial in $\mathbb{C}[t_1^{\pm 1}, \dots, t_m^{\pm 1}]$. We can write q as

$$q = \sum_k a_k t_1^{i_1^k} \cdot \dots \cdot t_m^{i_m^k},$$

for some $a_k \in \mathbb{C}^*$, and $(i_1^k, \dots, i_m^k) \neq (i_1^j, \dots, i_m^j)$ for all $k \neq j$. We define the *degree* of q as

$$\deg(q) = \max_k \left(\sum_{l=1}^m i_l^k \right) - \min_j \left(\sum_{l=1}^m i_l^j \right)$$

Remark 13. The degree of a Laurent polynomial q is 0 if and only if q is a homogeneous polynomial up to multiplication by a unit of $\mathbb{C}[t_1^{\pm 1}, \dots, t_m^{\pm 1}]$. Multiplying by a unit of $\mathbb{C}[t_1^{\pm 1}, \dots, t_m^{\pm 1}]$ does not change the degree, so $\deg(\Delta_C)$ is well-defined.

Lemma 4.

$$E_0(H_1(U; R_0)) = E_1(H_1(U, u_0; R_0)).$$

Proof. Note that R_0 is a commutative ring. From the long exact sequence of a pair with coefficients in R_0 , we get

$$0 \rightarrow H_1(U; R_0) \rightarrow H_1(U, u_0; R_0) \rightarrow H_0(u_0; R_0) \rightarrow H_0(U; R_0)$$

If $m \geq 2$, following the same Fox Calculus computation as in the proof of Lemma 1 for $H_0(X; R_0)$, we get that $H_0(U; R_0) = 0$, so we have the short exact sequence of R_0 -modules

$$0 \rightarrow H_1(U; R_0) \rightarrow H_1(U, u_0; R_0) \rightarrow R_0 \rightarrow 0$$

which splits, so the result follows. \square

In the above notation, we have the following result relating the multivariate Alexander polynomial and the zero-th higher order degree, thus extending Proposition 2 to the non-irreducible case.

Theorem 12. Suppose $m \geq 2$, and assume that $\delta_0(C)$ is finite. Then,

$$\delta_0(C) = \deg(\Delta_C).$$

Proof. Note that all the rings that we will deal with in this proof are commutative, so we will not distinguish between left and right modules. Let $\gamma_1, \dots, \gamma_m$ be positively oriented meridians around the different components of C , let $\psi : \pi_1(U) \rightarrow \mathbb{Z}$ be the linking number homomorphism. Consider the splitting of ψ given by

$$\begin{aligned} \phi : \mathbb{Z} &\longrightarrow \pi_1(U) \\ 1 &\longmapsto \gamma_1 \end{aligned}$$

which we use to identify R_0 with $\mathbb{K}_0[t^{\pm 1}]$. With this identification, we can think of R_0 as

$$Q \left(\mathbb{Z} \left[\left(\frac{t_2}{t_1} \right)^{\pm 1}, \dots, \left(\frac{t_m}{t_1} \right)^{\pm 1} \right] \right) [t_1^{\pm 1}]$$

where $\mathbb{K}_0 = Q \left(\mathbb{Z} \left[\left(\frac{t_2}{t_1} \right)^{\pm 1}, \dots, \left(\frac{t_m}{t_1} \right)^{\pm 1} \right] \right)$ is the field of quotients of $\mathbb{Z} \left[\left(\frac{t_2}{t_1} \right)^{\pm 1}, \dots, \left(\frac{t_m}{t_1} \right)^{\pm 1} \right]$.

We can think of $\mathbb{Z}[\Gamma_0]$ as $\mathbb{Z}[t_1^{\pm 1}, \dots, t_m^{\pm 1}]$ seen inside of R_0 this way. Note that the degree of any $q \in \mathbb{Z}[t_1^{\pm 1}, \dots, t_m^{\pm 1}]$ is the same as the degree of q in t seen as an element of $\mathbb{K}_0[t^{\pm 1}]$.

Let f_1, \dots, f_r be a set of generators of $E_1(A_G)$. We have that

$$\Delta_C = \gcd_{\mathbb{Z}[t_1^{\pm 1}, \dots, t_m^{\pm 1}]}(f_1, \dots, f_r).$$

Since R_0 is a flat $\mathbb{Z}[\Gamma_0]$ -module, f_1, \dots, f_r are also a set of generators of $E_1(H_1(U, u_0; R_0))$. By assumption, $\delta_0(C)$ is finite, which means that $E_1(H_1(U, u_0; R_0))$ is not the zero ideal. Moreover, $R_0 \cong \mathbb{K}_0[t^{\pm 1}]$ is a PID, so there exists $p \in \mathbb{K}_0[t^{\pm 1}]$ such that

$$E_1(H_1(U, u_0; R_0)) = (p)$$

and, by definition, $\delta_0(C) = \deg_{\mathbb{K}_0[t^{\pm 1}]}(p)$. Moreover, by considering the prime decomposition of f_1, \dots, f_r in $\mathbb{Z}[t_1^{\pm 1}, \dots, t_m^{\pm 1}]$ and applying Lemma 5 (below), we can choose such $p \in \mathbb{Z}[t_1^{\pm 1}, \dots, t_m^{\pm 1}]$ such that there exists $a \in \mathbb{Z}[t_1^{\pm 1}, \dots, t_m^{\pm 1}]$, where a is a unit in R_0 , with

$$p \cdot a = \Delta_C$$

But units in R_0 which are contained in $\mathbb{Z}[t_1^{\pm 1}, \dots, t_m^{\pm 1}]$ all have degree 0 as elements of $\mathbb{Z}[t_1^{\pm 1}, \dots, t_m^{\pm 1}]$, therefore

$$\deg_{\mathbb{Z}[t_1^{\pm 1}, \dots, t_m^{\pm 1}]}(\Delta_C) = \deg_{\mathbb{Z}[t_1^{\pm 1}, \dots, t_m^{\pm 1}]}(p) = \delta_0(C). \quad \square$$

Lemma 5. *Let $R = Q \left(\mathbb{Z} \left[\left(\frac{t_2}{t_1} \right)^{\pm 1}, \dots, \left(\frac{t_m}{t_1} \right)^{\pm 1} \right] \right) [t_1^{\pm 1}]$, and $S = \mathbb{Z}[t_1^{\pm 1}, \dots, t_m^{\pm 1}]$. Let q be a prime element in S . Then, q is either prime or a unit in R .*

Proof. This is an exercise in commutative algebra, which is a direct consequence of [1, Proposition 3.11, iv)], for example. \square

As a consequence of Proposition 3, Proposition 2, Theorem 12 and [5, Corollary 3.2], we get the following:

Corollary 1.

$$\text{codim } V_1(U) > 1 \iff \Delta_C \in \mathbb{Z} \setminus 0 \implies \delta_0(C) = 0.$$

Moreover, Proposition 2 and Theorem 12 imply the following:

Corollary 2. $\delta_0(C) = 0$ if and only if Δ_C has a representative in $\mathbb{Z}[t_1^{\pm 1}, \dots, t_m^{\pm 1}]$ that is a non-zero homogeneous polynomial.

Remark 14. Note that since \mathcal{K}_0 is flat over $\mathbb{Z}[\Gamma_0]$, by [6, Proposition 5.6] we have that

$$\text{codim } V_1(U) = 0 \iff E_1(A_G) = (0) \iff \text{rk}_{\mathcal{K}_0} H_1(U, u_0; \mathcal{K}_0) > 1 \iff \delta_0 \text{ is not finite.}$$

Hence, if $\delta_0(C)$ is finite (as is the case for all curves in general position at infinity ([11]), the ones described in Theorem 9, or all line arrangements except the one consisting of m parallel lines, for $m \geq 2$), then $\text{codim } V_1(U) \geq 1$.

Example 3. Let C be the line arrangement described in Example 2, and let t_i the variable corresponding to the component L_i , for $i = 1, \dots, m$. Then

$$\Delta_C = (t_1 - 1)^{m-2}.$$

Let us now specialize our results to the case when C is an essential line arrangement. We recall the following result from [17].

Proposition 4. ([17, Theorem 9.15]) *Let C be an essential line arrangement. Then:*

(1) *If C consists of m lines going through a point (a pencil of lines), with $m \geq 3$, then*

$$\Delta_C = (t_1 t_2 \dots t_m - 1)^{m-2}.$$

(2) *If C is as in Example 2, then $\Delta_C = (t_1 - 1)^{m-2}$.*

(3) *For all other essential arrangements, $\Delta_C \in \mathbb{Z} \setminus \{0\}$.*

Then Theorem 12 and Proposition 4 have the following consequence:

Corollary 3. *Let C be an essential line arrangement. Then:*

(1) *$\text{codim } V_1(C) > 1 \iff \delta_0(C) = 0$. That is, the result from Corollary 1 is an “if and only if” for line arrangements.*

(2) *If C is a pencil of lines, then $\delta_0(C) = m(m - 2)$.*

(3) *If C is as in Example 2, then $\delta_0(C) = (m - 2)$.*

(4) *If C is not as in items (2) or (3), then $\delta_0(C) = 0$.*

Remark 15. By Lemma 1 and Example 3, $\delta_n(C) = \delta_0(C)$ for all n for the cases described in items (2) and (3) of the above Corollary.

REFERENCES

- [1] M. F. Atiyah and I. G. Macdonald, *Introduction to commutative algebra*, Addison-Wesley Series in Mathematics, Westview Press, Boulder, CO, 2016.
- [2] T. D. Cochran, *Noncommutative knot theory*, *Algebr. Geom. Topol.* **4** (2004), 347–398.
- [3] J. I. Cogolludo and V. Florens, *Twisted Alexander polynomials of plane algebraic curves*, *J. Lond. Math. Soc.* **76** (2007), no. 1, 105–121.
- [4] A. Dimca, *Singularities and topology of hypersurfaces*, Universitext, Springer-Verlag, New York, 1992.
- [5] A. Dimca, S. Papadima, and A. Suciu, *Alexander polynomials: essential variables and multiplicities*, *Int. Math. Res. Not. IMRN* **3** (2008), Art. ID rmm119, 36.
- [6] S. L. Harvey, *Higher-order polynomial invariants of 3-manifolds giving lower bounds for the Thurston norm*, *Topology* **44** (2005), no. 5, 895–945.
- [7] E. Hironaka, *Alexander stratifications of character varieties*, *Ann. Inst. Fourier (Grenoble)* **47** (1997), no. 2, 555–583.
- [8] P. Kirk and C. Livingston, *Twisted Alexander invariants, Reidemeister torsion, and Casson-Gordon invariants*, *Topology* **38** (1999), no. 3, 635–661.
- [9] A. Libgober, *Alexander polynomials of plane algebraic curves and cyclic multiple planes*, *Duke Math. J.* **49** (1982), 833–851.
- [10] ———, *On the homology of finite abelian coverings*, *Topology Appl.* **43** (1992), no. 2, 157–166.
- [11] C. Leidy and L. Maxim, *Higher-order Alexander invariants of plane algebraic curves*, *Int. Math. Res. Not.* (2006), Art. ID 12976, 23.

- [12] ———, *Obstructions on fundamental groups of plane curve complements*, Real and complex singularities, Contemp. Math., vol. 459, Amer. Math. Soc., Providence, RI, 2008, pp. 117–130.
- [13] L. Maxim and K. Wong, *Twisted Alexander invariants of complex hypersurface complements*, Proc. Roy. Soc. Edinburgh Sect. A (2018), 1–25.
- [14] M. Oka and K. Sakamoto, *Product theorem of the fundamental group of a reducible curve*, J. Math. Soc. Japan **30** (1978), no. 5, 599–602.
- [15] M. Oka, *A survey on Alexander polynomials of plane curves*, Singularités Franco-Japonaises, Sémin. Congr., vol. 10, Soc. Math. France, Paris, 2005, pp. 209–232.
- [16] Y. Su, *Higher-order Alexander invariants of hypersurface complements*, arXiv:1510.03467.
- [17] A. I. Suci, *Fundamental groups, Alexander invariants, and cohomology jumping loci*, Topology of algebraic varieties and singularities, Contemp. Math., vol. 538, Amer. Math. Soc., Providence, RI, 2011, pp. 179–223.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN-MADISON, 480 LINCOLN DRIVE, MADISON WI 53706-1388, USA

E-mail address: evaelduque@math.wisc.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN-MADISON, 480 LINCOLN DRIVE, MADISON WI 53706-1388, USA

E-mail address: maxim@math.wisc.edu