Department of Mathematics, University of Wisconsin-Madison

Math 431 — Final Exam — Fall 2024

NAME : (as it appears on Canvas)

EMAIL: @wisc.edu

PROFESSOR: David Clancy or Mikhail Ivanov

Time: **120 minutes**

- This exam contains 8 questions some with multiple parts, 14 pages (including the cover) for the total of 109 points. Read the problems carefully and budget your time wisely.
- You are allowed a single sheet of hand-written notes.
- **NO CALCULATORS** or other electronic devices are to be used. Turn off your phone so as to not disturb others.
- You **do not** need to simplify binomial coefficients or factorials.
- Please present your solutions in a clear manner. Cross out any writing that you do not wish to be graded.
- Justify your steps.
- For binomial random variable approximations, you will NOT get full credit for your answer if you do not include some explanation of why that approximation is valid.
- You can use the attached table of the values of the $\Phi(x)$ function.
- If you use an additional page for a particular problem, be sure to **CLEARLY** indicate this on the problem's page so I know to look further.

Question:	1	2	3	4	5	6	7	8	Total
Points:	12	15	8	20	11	20	15	8	109
Score:									

1. (12 points) For this problem, you need to provide **answers only**.

   (a) Let X be the random variable with moment generating function

   $$M_X(t) = \frac{1}{4} + \frac{1}{8}e^{2t} + \frac{1}{8}e^{3t} + \frac{1}{2}e^{5t}.$$

   What is the p.m.f. of X?

   (b) A full house in poker consists of 5 cards where three are the same value and the remaining two are the same value (e.g. three Q's and two 4's). If 5 cards are dealt from a shuffled deck without replacement, what is the probability of drawing a full house?

   (c) Let X and Y be i.i.d. Unif$[0, 1]$. Compute the covariance of $X + Y$ and $X - Y$.

   (d) We flip a coin over and over again. Let N_n be the number of flips required to see the nth Heads. What is $\lim_{n\to\infty} P(N_n > 3n)$?

2. (15 points) **Please provide explanation for all parts.**

Suppose that X and Y are random variables with

$$E[X] = 2, \qquad E[X^2] = 5, \qquad E[Y] = 4, \qquad \mathrm{Cov}(X, Y) = -4.$$

(a) What is $\mathrm{Var}(X)$.

(b) What is $E[XY]$?

(c) What is $\mathrm{Cov}(2X + Y, X)$?

3. (8 points) Let X and Y be two independent random variables where $X \sim \text{Exp}(1)$ and $Y \sim \text{Unif}(-1, 1)$. What is the probability density function of $Z = X + Y$?

4. Let X and Y be jointly continuous random variables with joint density function

$$f_{X,Y}(x,y) = \begin{cases} \frac{2}{y}e^{-2y} & : 0 < x < y \\ 0 & : \text{else} \end{cases}$$

(a) (5 points) Find the marginal density of Y, i.e. $f_Y(y)$. Specify the name of the distribution and its parameters.

(b) (5 points) Determine the conditional distribution $f_{X|Y}(x|y)$ for each $y > 0$?

Problem 4, cont.

(c) (5 points) What is $E[X|Y = y]$ for each $y > 0$?

(d) (5 points) What is $E[X]$?

5. There is an urn with 30 red balls, 20 green balls, and 20 yellow balls. You sample 40 balls from the urn without replacement, one after the other.

   (a) (5 points) Given that the last ball drawn is red, what is the probability the first ball drawn is yellow?

   (b) (6 points) Let X be the number of times that you draw three consecutive balls of the same color. What is $E[X]$?

6. Let X be a non-negative random variable. Each part requires an explanation.

(a) (5 points) We know that $E[X] = 200$, what can we say about $P(X > 300)$?

(b) (5 points) Suppose we also know that $\text{Var}(X) = 3600$, what can we say about $P(X > 300)$?

(c) (5 points) Suppose we also know that X is the sum of 50 i.i.d. random variables. What is the expectation and variance of the summand? That is if $X = \sum_{j=1}^{50} Y_j$, where Y_j are i.i.d., then what is $E[Y_j]$ and $\text{Var}(Y_j)$?

(d) (5 points) Given the information in (c) provide an estimate for the precise value of $P(X > 300)$, assuming it is reasonable.

7. A fair 6-sided die is rolled, and then a fair coin is flipped as many times as the die roll says, e.g., if the result of the die roll is a 3, then the coin is flipped 3 times. Let X be the result of the die roll and Y be the number of times the coin lands Heads.

   (a) (5 points) Find the joint pmf of X and Y.

   (b) (5 points) Find $P(Y = 5)$.

   (c) (5 points) What is $E[X|Y = 5]$?

8. We have two bins. The first bin has 6 blue marbles and 4 yellow marbles. The second bin has 3 blue marbles and 4 yellow marbles. We flip a coin. If it is heads, we sample from the first bin, otherwise we sample from the second.

   (a) (4 points) If the marble we select is yellow, what is the probability that we chose the first bin?

   (b) (4 points) Now suppose we put the yellow marble from (a) back in the bin it was drawn from and then draw a marble from the same bin. This marble is also yellow. What is the probability we chose the first bin?

SCRATCH PAPER - DO NOT REMOVE FROM YOUR EXAM.
SCRATCH WORK WILL NOT BE GRADED

SCRATCH PAPER - DO NOT REMOVE FROM YOUR EXAM
SCRATCH WORK WILL NOT BE GRADED

Table of Named Distributions

Name	p.m.f. or p.d.f.	$E[X]$	$Var(X)$
$\text{Ber}(p)$	$p_X(0) = 1-p,\ p_X(1) = p$	p	$p(1-p)$
$\text{Bin}(n,p)$	$p_X(k) = \binom{n}{k}p^k(1-p)^{n-k},\ 0 \le k \le n$	np	$np(1-p)$
$\text{Geom}(p)$	$p_X(k) = p(1-p)^{k-1},\ k \ge 1$	$\frac{1}{p}$	$\frac{1-p}{p^2}$
$\text{NegBin}(k,p)$	$p_X(n) = \binom{n-1}{k-1}p^k(1-p)^{n-k},\ n \ge k$	$\frac{k}{p}$	$\frac{k(1-p)}{p^2}$
$\text{HyperGeom}(N,N_A,n)$	$p_X(k) = \frac{\binom{N_A}{k}\binom{N-N_A}{n-k}}{\binom{N}{n}}$	$\frac{nN_A}{N}$	$\frac{nN_A(N-N_A)(N-n)}{N^2(N-1)}$
$\text{Poisson}(\lambda)$	$p_X(k) = \frac{\lambda^k}{k!}e^{-\lambda},\ k \ge 0$	λ	λ
$\text{Unif}[a,b]$	$f_X(x) = \frac{1}{b-a},\ x \in [a,b]$	$\frac{a+b}{2}$	$\frac{(b-a)^2}{12}$
$\text{N}(\mu,\sigma^2)$	$f_X(x) = \frac{1}{\sqrt{2\pi}\sigma}2e^{-\frac{(x-\mu)^2}{2\sigma^2}}$	μ	σ^2
$\text{Exp}(\lambda)$	$f_X(x) = \lambda e^{-\lambda x},\ x \ge 0$	$\frac{1}{\lambda}$	$\frac{1}{\lambda^2}$
$\text{Gamma}(n,\lambda)$	$f_X(x) = \frac{\lambda^n x^{n-1}e^{-\lambda}}{(n-1)!},\ x \ge 0$	$\frac{n}{\lambda}$	$\frac{n}{\lambda^2}$

Table of values for $\Phi(x)$, the CDF of a standard normal random variable

	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0.0	0.5000	0.5040	0.5080	0.5120	0.5160	0.5199	0.5239	0.5279	0.5319	0.5359
0.1	0.5398	0.5438	0.5478	0.5517	0.5557	0.5596	0.5636	0.5675	0.5714	0.5753
0.2	0.5793	0.5832	0.5871	0.5910	0.5948	0.5987	0.6026	0.6064	0.6103	0.6141
0.3	0.6179	0.6217	0.6255	0.6293	0.6331	0.6368	0.6406	0.6443	0.6480	0.6517
0.4	0.6554	0.6591	0.6628	0.6664	0.6700	0.6736	0.6772	0.6808	0.6844	0.6879
0.5	0.6915	0.6950	0.6985	0.7019	0.7054	0.7088	0.7123	0.7157	0.7190	0.7224
0.6	0.7257	0.7291	0.7324	0.7357	0.7389	0.7422	0.7454	0.7486	0.7517	0.7549
0.7	0.7580	0.7611	0.7642	0.7673	0.7704	0.7734	0.7764	0.7794	0.7823	0.7852
0.8	0.7881	0.7910	0.7939	0.7967	0.7995	0.8023	0.8051	0.8078	0.8106	0.8133
0.9	0.8159	0.8186	0.8212	0.8238	0.8264	0.8289	0.8315	0.8340	0.8365	0.8389
1.0	0.8413	0.8438	0.8461	0.8485	0.8508	0.8531	0.8554	0.8577	0.8599	0.8621
1.1	0.8643	0.8665	0.8686	0.8708	0.8729	0.8749	0.8770	0.8790	0.8810	0.8830
1.2	0.8849	0.8869	0.8888	0.8907	0.8925	0.8944	0.8962	0.8980	0.8997	0.9015
1.3	0.9032	0.9049	0.9066	0.9082	0.9099	0.9115	0.9131	0.9147	0.9162	0.9177
1.4	0.9192	0.9207	0.9222	0.9236	0.9251	0.9265	0.9279	0.9292	0.9306	0.9319
1.5	0.9332	0.9345	0.9357	0.9370	0.9382	0.9394	0.9406	0.9418	0.9429	0.9441
1.6	0.9452	0.9463	0.9474	0.9484	0.9495	0.9505	0.9515	0.9525	0.9535	0.9545
1.7	0.9554	0.9564	0.9573	0.9582	0.9591	0.9599	0.9608	0.9616	0.9625	0.9633
1.8	0.9641	0.9649	0.9656	0.9664	0.9671	0.9678	0.9686	0.9693	0.9699	0.9706
1.9	0.9713	0.9719	0.9726	0.9732	0.9738	0.9744	0.9750	0.9756	0.9761	0.9767
2.0	0.9772	0.9778	0.9783	0.9788	0.9793	0.9798	0.9803	0.9808	0.9812	0.9817
2.1	0.9821	0.9826	0.9830	0.9834	0.9838	0.9842	0.9846	0.9850	0.9854	0.9857
2.2	0.9861	0.9864	0.9868	0.9871	0.9875	0.9878	0.9881	0.9884	0.9887	0.9890
2.3	0.9893	0.9896	0.9898	0.9901	0.9904	0.9906	0.9909	0.9911	0.9913	0.9916
2.4	0.9918	0.9920	0.9922	0.9925	0.9927	0.9929	0.9931	0.9932	0.9934	0.9936
2.5	0.9938	0.9940	0.9941	0.9943	0.9945	0.9946	0.9948	0.9949	0.9951	0.9952
2.6	0.9953	0.9955	0.9956	0.9957	0.9959	0.9960	0.9961	0.9962	0.9963	0.9964
2.7	0.9965	0.9966	0.9967	0.9968	0.9969	0.9970	0.9971	0.9972	0.9973	0.9974
2.8	0.9974	0.9975	0.9976	0.9977	0.9977	0.9978	0.9979	0.9979	0.9980	0.9981
2.9	0.9981	0.9982	0.9982	0.9983	0.9984	0.9984	0.9985	0.9985	0.9986	0.9986
3.0	0.9987	0.9987	0.9987	0.9988	0.9988	0.9989	0.9989	0.9989	0.9990	0.9990
3.1	0.9990	0.9991	0.9991	0.9991	0.9992	0.9992	0.9992	0.9992	0.9993	0.9993
3.2	0.9993	0.9993	0.9994	0.9994	0.9994	0.9994	0.9994	0.9995	0.9995	0.9995
3.3	0.9995	0.9995	0.9995	0.9996	0.9996	0.9996	0.9996	0.9996	0.9996	0.9997
3.4	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9998