

# Billiard arrays and finite-dimensional irreducible $U_q(\mathfrak{sl}_2)$ -modules

Paul Terwilliger

University of Wisconsin-Madison

In this talk we will describe the notion of a **Billiard Array**.

This is a triangular array of one-dimensional subspaces of a finite-dimensional vector space, subject to several conditions that specify which sums are direct.

We will use Billiard Arrays to describe the finite-dimensional irreducible modules for the quantum algebra  $U_q(\mathfrak{sl}_2)$  and the Lie algebra  $\mathfrak{sl}_2$ .

# Billiard Arrays


Our topic is informally described as follows.

As in the game of Billiards, we start with an array of billiard balls arranged to form an equilateral triangle.

We assume that there are  $N + 1$  balls along each boundary, with  $N \geq 0$ .

## Billiard Arrays, Cont.

For  $N = 3$  the balls are centered at the following locations:



For us, each ball in the array represents a one-dimensional subspace of an  $(N + 1)$ -dimensional vector space  $V$  over a field  $\mathbb{F}$ .

## Billiard Arrays, Cont.

We impose two conditions on the array, that specify which sums are direct.

The first condition is that, for each set of balls on a line parallel to a boundary, their sum is direct.

The second condition is described as follows.

Three mutually adjacent balls in the array are said to form a **3-clique**. There are two kinds of 3-cliques:  $\Delta$  (**black**) and  $\nabla$  (**white**).

The second condition is that, for any three balls in the array that form a black 3-clique, their sum is not direct.

## Billiard Arrays, Cont.

Whenever the above two conditions are met, our array is called a **Billiard Array** on  $V$ .

We say that the Billiard Array is **over**  $\mathbb{F}$ , and call  $N$  the **diameter**.

Notation: For  $1 \leq i \leq N + 1$  let  $\mathcal{P}_i(V)$  denote the set of subspaces of  $V$  that have dimension  $i$ .

# The set $\Delta_N$

To describe our Billiard Array on  $V$ , we use the following construction.

Recall the natural numbers  $\mathbb{N} = \{0, 1, 2, \dots\}$ .

## Definition

Let  $\Delta_N$  denote the set consisting of the 3-tuples of natural numbers whose sum is  $N$ .

Thus

$$\Delta_N = \{(r, s, t) \mid r, s, t \in \mathbb{N}, r + s + t = N\}.$$

## The set $\Delta_N$ , cont.

We arrange the elements of  $\Delta_N$  in a triangular array.

For  $N = 3$ , the array looks as follows after deleting all punctuation:

```
      030
     120 021
    210 111 012
   300 201 102 003
```

An element in  $\Delta_N$  is called a **location**.



## The set $\Delta_N$ , cont.

We view our Billiard Array on  $V$  as a function  
 $B : \Delta_N \rightarrow \mathcal{P}_1(V), \lambda \mapsto B_\lambda.$

For  $\lambda \in \Delta_N$ ,  $B_\lambda$  is the billiard ball/subspace at location  $\lambda$ .

# Summary of Results

In this talk we obtain three main results, which are summarized as follows:

- (i) We show that the Billiard Arrays on  $V$  are in bijection with the 3-tuples of totally opposite flags on  $V$ ;
- (ii) We classify the Billiard Arrays up to isomorphism;
- (iii) We use Billiard Arrays to describe the finite-dimensional irreducible modules for the quantum algebra  $U_q(\mathfrak{sl}_2)$  and the Lie algebra  $\mathfrak{sl}_2$ .

We now describe our results in more detail.

Until further notice,  $V$  denotes a vector space over  $\mathbb{F}$  with dimension  $N + 1$ .

## Definition

By a **flag on  $V$**  we mean a sequence  $\{U_i\}_{i=0}^N$  such that  $U_i \in \mathcal{P}_{i+1}(V)$  for  $0 \leq i \leq N$  and  $U_{i-1} \subseteq U_i$  for  $1 \leq i \leq N$ .

# Constructing Flags

We now show how to construct a flag on  $V$ .

## Definition

By a **decomposition** of  $V$  we mean a sequence  $\{V_i\}_{i=0}^N$  of one-dimensional subspaces of  $V$  whose direct sum is  $V$ .

Given a decomposition  $\{V_i\}_{i=0}^N$  of  $V$ .

Define  $U_i = V_0 + \cdots + V_i$  for  $0 \leq i \leq N$ . Then the sequence  $\{U_i\}_{i=0}^N$  is a flag on  $V$ .

This flag is said to be **induced** by the decomposition  $\{V_i\}_{i=0}^N$ .

# Two opposite flags on $V$

Let  $\{U_i\}_{i=0}^N$  and  $\{U'_i\}_{i=0}^N$  denote flags on  $V$ .

## Definition

The above flags are called **opposite** whenever  $U_i \cap U'_j = 0$  if  $i + j < N$  ( $0 \leq i, j \leq N$ ).

## Lemma

*The flags  $\{U_i\}_{i=0}^N$  and  $\{U'_i\}_{i=0}^N$  are opposite if and only if there exists a decomposition  $\{V_i\}_{i=0}^N$  of  $V$  that induces  $\{U_i\}_{i=0}^N$  and whose inversion  $\{V_{N-i}\}_{i=0}^N$  induces  $\{U'_i\}_{i=0}^N$ .*

*In this case  $V_i = U_i \cap U'_{N-i}$  for  $0 \leq i \leq N$ .*

# Three totally opposite flags on $V$

## Definition

Suppose we are given three flags on  $V$ , denoted

$$\{U_i\}_{i=0}^N, \quad \{U'_i\}_{i=0}^N, \quad \{U''_i\}_{i=0}^N.$$

These flags are said to be **totally opposite** whenever

$$U_{N-r} \cap U'_{N-s} \cap U''_{N-t} = 0$$

for all integers  $r, s, t$  ( $0 \leq r, s, t \leq N$ ) such that  $r + s + t > N$ .

# Totally opposite vs mutually opposite

If three flags on  $V$  are totally opposite, then they are mutually opposite.

However, the converse is not true in general.

# From three totally opposite flags to a Billiard Array

Suppose we are given three totally opposite flags on  $V$ , denoted

$$\{U_i\}_{i=0}^N, \quad \{U'_i\}_{i=0}^N, \quad \{U''_i\}_{i=0}^N.$$

Using these flags we now construct a Billiard Array on  $V$ .

## Lemma

For each location  $\lambda = (r, s, t)$  in  $\Delta_N$  define

$$B_\lambda = U_{N-r} \cap U'_{N-s} \cap U''_{N-t}.$$

Then  $B_\lambda$  has dimension one. Moreover the map  $B : \Delta_N \rightarrow \mathcal{P}_1(V)$ ,  $\lambda \mapsto B_\lambda$  is a Billiard Array on  $V$ .



# From Billiard a Array to three totally opposite flags

We just went from flags to Billiard Arrays; we now reverse the direction.

Let  $B$  denote a Billiard Array on  $V$ .

By the **1-corner** of  $\Delta_N$  we mean the location  $(N, 0, 0)$ .

The **2-corner** and **3-corner** of  $\Delta_N$  are similarly defined.

# From Billiard Arrays to three totally opposite flags

## Lemma

For our Billiard Array  $B$  on  $V$ , for  $0 \leq i \leq N$  let  $U_i$  (resp.  $U'_i$ ) (resp.  $U''_i$ ) denote the sum of the balls in  $B$  that are at most  $i$  balls over from the 1-corner (resp. 2-corner) (resp. 3-corner). Then

$$\{U_i\}_{i=0}^N, \quad \{U'_i\}_{i=0}^N, \quad \{U''_i\}_{i=0}^N$$

are totally opposite flags on  $V$ .

Consider the following two sets:

- (i) the Billiard Arrays on  $V$ ;
- (ii) the 3-tuples of totally opposite flags on  $V$ .

In the previous two lemmas we described a function from (i) to (ii) and a function from (ii) to (i).

## Theorem

*The above functions are inverses, and hence bijections.*

# The classification of Billiard Arrays

Our next goal is to classify the Billiard Arrays up to isomorphism.

Let  $B$  denote a Billiard Array on the above vector space  $V$ .

Let  $V'$  denote a vector space over  $\mathbb{F}$  with dimension  $N + 1$ , and let  $B'$  denote a Billiard Array on  $V'$ .

The Billiard Arrays  $B, B'$  are called **isomorphic** whenever there exists an  $\mathbb{F}$ -linear bijection  $V \rightarrow V'$  that sends  $B_\lambda \mapsto B'_\lambda$  for all  $\lambda \in \Delta_N$ .

# Value functions

By a **value function on**  $\Delta_N$  we mean a function  $\Delta_N \rightarrow \mathbb{F} \setminus \{0\}$ .

For  $N \leq 1$ , up to isomorphism there exists a unique Billiard Array over  $\mathbb{F}$  that has diameter  $N$ .

For  $N \geq 2$  we will obtain a bijection between the following two sets:

- (i) the isomorphism classes of Billiard Arrays over  $\mathbb{F}$  that have diameter  $N$ ;
- (ii) the value functions on  $\Delta_{N-2}$ .

# The affine braces for a Billiard Array

We now describe the bijection from the previous slide.

Let  $B$  denote a Billiard Array on  $V$ .

## Definition

Let  $\lambda, \mu, \nu$  denote locations in  $\Delta_N$  that form a black 3-clique. By an **affine brace** (or **abrace**) for this clique, we mean a set of vectors

$$u \in B_\lambda, \quad v \in B_\mu, \quad w \in B_\nu$$

that are not all zero, and  $u + v + w = 0$ . (In fact each of  $u, v, w$  is nonzero).

# The affine braces for a Billiard Array, cont.

Here is an example of an abrace.

## Example

Let  $\lambda, \mu, \nu$  denote locations in  $\Delta_N$  that form a black 3-clique. Pick any nonzero vectors

$$u \in B_\lambda, \quad v \in B_\mu, \quad w \in B_\nu.$$

The vectors  $u, v, w$  are linearly dependent. So there exist scalars  $a, b, c$  in  $\mathbb{F}$ , not all zero, such that  $au + bv + cw = 0$ . The vectors  $au, bv, cw$  form an abrace for the clique.

# The affine braces for a Billiard Array, cont.

Affine braces have the following property.

## Lemma

*Let  $\lambda, \mu, \nu$  denote locations in  $\Delta_N$  that form a black 3-clique. Then each nonzero vector in  $B_\lambda$  is contained in a unique abrace for this clique.*



# The braces for a Billiard Array

We have been discussing affine braces.

We now consider a variation on this concept, called a brace.

## Definition

Let  $\lambda, \mu$  denote adjacent locations in  $\Delta_N$ . Note that there exists a unique location  $\nu \in \Delta_N$  such that  $\lambda, \mu, \nu$  form a black 3-clique. We call  $\nu$  the **completion** of the pair  $\lambda, \mu$ .

# The braces for a Billiard Array, cont.

## Definition

Let  $\lambda, \mu$  denote adjacent locations in  $\Delta_N$ . By a **brace** for  $\lambda, \mu$  we mean a set of nonzero vectors

$$u \in B_\lambda, \quad v \in B_\mu$$

such that  $u + v \in B_\nu$ . Here  $\nu$  denotes the completion of  $\lambda, \mu$ .

# The braces for a Billiard Array, cont.

Braces have the following property.

## Lemma

*Let  $\lambda, \mu$  denote adjacent locations in  $\Delta_N$ . Each nonzero vector in  $B_\lambda$  is contained in a unique brace for  $\lambda, \mu$ .*

We now define some maps  $\tilde{B}_{\lambda,\mu}$ .

## Definition

Let  $\lambda, \mu$  denote adjacent locations in  $\Delta_N$ . We define an  $\mathbb{F}$ -linear map  $\tilde{B}_{\lambda,\mu} : B_\lambda \rightarrow B_\mu$  as follows. This map sends each nonzero  $u \in B_\lambda$  to the unique  $v \in B_\mu$  such that  $u, v$  is a brace for  $\lambda, \mu$ .

# The maps $\tilde{B}_{\lambda,\mu}$ , cont.

Let  $\lambda, \mu$  denote adjacent locations in  $\Delta_N$ .

We just defined an  $\mathbb{F}$ -linear map  $\tilde{B}_{\lambda,\mu} : B_\lambda \rightarrow B_\mu$ .

We now consider what happens when we compose the maps of this kind.

# The maps $\tilde{B}_{\lambda,\mu}$ , cont.

## Lemma

Let  $\lambda, \mu$  denote adjacent locations in  $\Delta_N$ . Then the maps  $\tilde{B}_{\lambda,\mu} : B_\lambda \rightarrow B_\mu$  and  $\tilde{B}_{\mu,\lambda} : B_\mu \rightarrow B_\lambda$  are inverses.

# The maps $\tilde{B}_{\lambda,\mu}$ , cont.

## Lemma

Let  $\lambda, \mu, \nu$  denote locations in  $\Delta_N$  that form a black 3-clique.  
Then the composition around the clique:

$$B_\lambda \xrightarrow{\tilde{B}_{\lambda,\mu}} B_\mu \xrightarrow{\tilde{B}_{\mu,\nu}} B_\nu \xrightarrow{\tilde{B}_{\nu,\lambda}} B_\lambda$$

is equal to the identity map on  $B_\lambda$ .

# The maps $\tilde{B}_{\lambda,\mu}$ , cont.

## Definition

Let  $\lambda, \mu, \nu$  denote locations in  $\Delta_N$  that form a white 3-clique. Then the composition around the clique:

$$B_\lambda \xrightarrow{\tilde{B}_{\lambda,\mu}} B_\mu \xrightarrow{\tilde{B}_{\mu,\nu}} B_\nu \xrightarrow{\tilde{B}_{\nu,\lambda}} B_\lambda$$

is a nonzero scalar multiple of the identity map on  $B_\lambda$ . The scalar is called the **clockwise  $B$ -value** (resp. **c.clockwise  $B$ -value**) of the clique whenever the sequence  $\lambda, \mu, \nu$  runs clockwise (resp. c.clockwise) around the clique.



# Clockwise and c.clockwise $B$ -values

## Lemma

*For each white 3-clique in  $\Delta_N$ , its clockwise  $B$ -value and c.clockwise  $B$ -value are reciprocal.*

## Definition

For each white 3-clique in  $\Delta_N$ , by its  **$B$ -value** we mean the clockwise  $B$ -value.

We have now assigned a nonzero scalar value to each white 3-clique in  $\Delta_N$ .

# The value function $\hat{B}$

For  $N \leq 1$  the set  $\Delta_N$  has no white 3-clique.

For  $N \geq 2$  we now give a bijection from  $\Delta_{N-2}$  to the set of white 3-cliques in  $\Delta_N$ .

The bijection sends each element  $(r, s, t)$  in  $\Delta_{N-2}$  to the white 3-clique in  $\Delta_N$  consisting of the locations

$$(r, s + 1, t + 1), \quad (r + 1, s, t + 1), \quad (r + 1, s + 1, t).$$

# The value function $\hat{B}$

Recall our Billiard Array  $B$  on  $V$ .

## Definition

Using  $B$  we define a function  $\hat{B} : \Delta_{N-2} \rightarrow \mathbb{F}$  as follows:  $\hat{B}$  sends each element  $(r, s, t)$  in  $\Delta_{N-2}$  to the  $B$ -value of the corresponding white 3-clique in  $\Delta_N$ .

By construction  $\hat{B}$  is a value function on  $\Delta_{N-2}$ .

We call  $\hat{B}$  the **value function for  $B$** .

# The classification of Billiard Arrays

We now classify the Billiard Arrays up to isomorphism.

Recall the Billiard Array  $B$  and its value function  $\hat{B}$ .

## Theorem

*The map  $B \mapsto \hat{B}$  induces a bijection between the following two sets:*

- (i) the isomorphism classes of Billiard Arrays over  $\mathbb{F}$  that have diameter  $N$ ;*
- (ii) the value functions on  $\Delta_{N-2}$ .*

# Billiard Arrays and $U_q(\mathfrak{sl}_2)$

We now use Billiard Arrays to describe the finite-dimensional irreducible modules for  $U_q(\mathfrak{sl}_2)$  and  $\mathfrak{sl}_2$ .

# $U_q(\mathfrak{sl}_2)$ and its modules

We recall the quantum algebra  $U_q(\mathfrak{sl}_2)$ .

We will use the **equitable presentation**.

Fix a nonzero  $q \in \mathbb{F}$  such that  $q^2 \neq 1$ .

# The definition of $U_q(\mathfrak{sl}_2)$

## Definition

Let  $U_q(\mathfrak{sl}_2)$  denote the associative  $\mathbb{F}$ -algebra with generators  $x, y^{\pm 1}, z$  and relations  $yy^{-1} = y^{-1}y = 1$ ,

$$\frac{qxy - q^{-1}yx}{q - q^{-1}} = 1,$$

$$\frac{qyz - q^{-1}zy}{q - q^{-1}} = 1,$$

$$\frac{qzx - q^{-1}xz}{q - q^{-1}} = 1.$$

The  $x, y^{\pm 1}, z$  are called the **equitable generators** for  $U_q(\mathfrak{sl}_2)$ .

# The elements $\nu_x, \nu_y, \nu_z$

The defining relations for  $U_q(\mathfrak{sl}_2)$  can be reformulated as follows:

$$q(1 - yz) = q^{-1}(1 - zy),$$

$$q(1 - zx) = q^{-1}(1 - xz),$$

$$q(1 - xy) = q^{-1}(1 - yx).$$

Denote these common values by  $\nu_x, \nu_y, \nu_z$  respectively.



# How $x, y, z$ are related to $\nu_x, \nu_y, \nu_z$

The  $x, y, z$  are related to  $\nu_x, \nu_y, \nu_z$  as follows:

$$\begin{aligned}x\nu_y &= q^2\nu_yx, & x\nu_z &= q^{-2}\nu_zx, \\y\nu_z &= q^2\nu_zy, & y\nu_x &= q^{-2}\nu_xy, \\z\nu_x &= q^2\nu_xz, & z\nu_y &= q^{-2}\nu_yz.\end{aligned}$$

Assume that  $q$  is not a root of unity.

Pick  $N \in \mathbb{N}$ , and let  $V$  denote an irreducible  $U_q(\mathfrak{sl}_2)$ -module with dimension  $N + 1$ .

Then each of  $\nu_x, \nu_y, \nu_z$  is **nilpotent** on  $V$ .

Moreover, each of the following is a flag on  $V$ :

$$\{\nu_x^{N-i} V\}_{i=0}^N, \quad \{\nu_y^{N-i} V\}_{i=0}^N, \quad \{\nu_z^{N-i} V\}_{i=0}^N$$

## Theorem

Assume that  $q$  is not a root of unity. Let  $V$  denote an irreducible  $U_q(\mathfrak{sl}_2)$ -module of dimension  $N + 1$ . Then:

(i) the flags

$$\{\nu_x^{N-i} V\}_{i=0}^N, \quad \{\nu_y^{N-i} V\}_{i=0}^N, \quad \{\nu_z^{N-i} V\}_{i=0}^N$$

are totally opposite;

(ii) for the corresponding Billiard Array on  $V$ , the value of each white 3-clique is a constant  $q^{-2}$ .

# Billiard Arrays and $\mathfrak{sl}_2$

We just obtained a Billiard Array from each finite-dimensional irreducible  $U_q(\mathfrak{sl}_2)$ -module, under the assumption that  $q$  is not a root of unity.

For the Lie algebra  $\mathfrak{sl}_2$  over  $\mathbb{F}$ , one similarly obtains a Billiard Array from each finite-dimensional irreducible  $\mathfrak{sl}_2$ -module, under the assumption that  $\mathbb{F}$  has characteristic zero.

For these Billiard Arrays the value of each white 3-clique is 1.

# Summary

In this talk, we introduced the notion of a Billiard Array.

We showed that the Billiard Arrays on a vector space  $V$  are in bijection with the 3-tuples of totally opposite flags on  $V$ .

We classified the Billiard Arrays up to isomorphism.

We used Billiard Arrays to describe the finite-dimensional irreducible modules for the quantum algebra  $U_q(\mathfrak{sl}_2)$  and the Lie algebra  $\mathfrak{sl}_2$ .

Thank you for your attention!

## THE END

T. Ito, P. Terwilliger, C. Weng. The quantum algebra  $U_q(\mathfrak{sl}_2)$  and its equitable presentation. *J. Algebra* **298** (2006) 284–301.  
arXiv:math/0507477.

P. Terwilliger. The universal Askey-Wilson algebra and the equitable presentation of  $U_q(\mathfrak{sl}_2)$ . *SIGMA* **7** (2011) 099, 26 pages,  
arXiv:1107.3544.

P. Terwilliger. Finite-dimensional irreducible  $U_q(\mathfrak{sl}_2)$ -modules from the equitable point of view. arXiv:1303.6134.

P. Terwilliger. Billiard Arrays and finite-dimensional irreducible  $U_q(\mathfrak{sl}_2)$ -modules. arXiv:1408.0143.