

MATH 240; EXAM # 2, 100 points, November 8, 2002 (R.A.Brualdi)

TOTAL SCORE (10 problems):

Name: Solutions

TA (circle): Matthew Petro Dilip Raghavan

Disc. (circle) TUES. THURS. TIME:

1. [15 points] Calculate explicitly the following:

(a)  $P(7, 4) = 7!/3! = 840$

(b)  $C(8, 5) = \binom{8}{5} = 8!/5!3! = 56$

(c) The **coefficient** of  $x^5y^3$  in the expansion of  $(2x - y)^8$ :

$$\binom{8}{3}2^5(-1)^3 = -1792$$

(d) The **Boolean matrix product**

$$\begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} \odot \begin{bmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}.$$

(e) Let  $S$  be a program segment, and let  $p$  and  $q$  be propositions. **Explain the meaning** of the symbol  $p\{S\}q$ .

If proposition  $p$  holds and the program segment  $S$  terminates, Then proposition  $q$  holds.

2. [12 points] Use **mathematical induction** to prove that

$$3^{2n} \equiv 1 \pmod{8} \text{ for all } n \geq 0.$$

**Be sure to specify the two main steps in carrying out the induction.**

(1) basis step ( $n = 0$  in this case): One easily checks that  $3^{2 \cdot 0} \equiv 1 \pmod{8}$ .

(2) inductive step: one needs to show that  $3^{2n} \equiv 1 \pmod{8}$  implies  $3^{2(n+1)} \equiv 1 \pmod{8}$ .

So suppose that  $3^{2n} \equiv 1 \pmod{8}$ . Then  $3^{2(n+1)} = 3^{2n} \cdot 3^2 = 3^{2n} \cdot 9 \equiv 1 \cdot 1 \equiv 1 \pmod{8}$ , since  $9 \equiv 1 \pmod{8}$ .

Thus the statement follows by induction.

3. [5 points]

Consider a theorem of the form:

**Theorem: If  $p$ , then  $q$ .**

Explain the difference between a **direct proof** and an **indirect proof** of the Theorem.

In a direct proof, one assumes that  $p$  is true and shows that  $q$  is true; i.e. the line  $p$  true and  $q$  false cannot occur in the truth table of the implication.

In an indirect proof, one assumes that  $q$  is false, and shows that  $p$  is false, leading to the same conclusion.

4. [12 points] Let  $s_n = 1^5 + 2^5 + \dots + n^5$ . Give recursive and iterative algorithms for computing  $s_n$ .

(i) **Recursive Algorithm:**

If  $n = 1$ , then  $s_1 := 1$ .

Else  $s_n := s_{n-1} + n^5$ .

(ii) **Iterative Algorithm:**

$s_1 := 1$

For  $i = 2, 3, \dots, n$ ,  $s_i := s_{i-1} + i^5$ .

5. [12 points] A password is to be 7 characters long with each character equal to one of the 10 digits  $0, 1, \dots, 9$  or one of the 26 lowercase letters  $a, b, c, \dots, x, y, z$ . (Repeated characters are allowed.)

(i) **How many** passwords are there such that the **characters are distinct and one of the characters equals 0**?

$$7 \cdot 35 \cdot 34 \cdot \dots \cdot 30$$

(ii) If a password is chosen at random from all possible passwords, what is the **probability** its characters are distinct and one of the characters equals 0?

$$\text{above answer}/36^7$$

6. [12 points] A bakery has 6 different kinds of bagels and has on hand a large number of each kind. You want to buy a bag of 2 dozen bagels containing at least one bagel of each type. **How many different bags could you buy?**

Number of solutions in nonnegative integers of  $x_1 + x_2 + \dots + x_6 = 18$  and so  $\binom{18+6-1}{5} = \binom{23}{5}$

7. [10 points] Messages are to be transmitted over a communications channel using three different signals: one requires 1 nanosecond, another requires 2 nanoseconds, and a third requires 4 nanoseconds. A message consists of sequences of these three signals where each signal is immediately followed by the next signal.

- (i) Give a **recurrence relation** for the number  $f_n$  of different messages that can be sent in exactly  $n$  nanoseconds,  $n \geq 1$ .

$$f_n = f_{n-1} + f_{n-2} + f_{n-4}, m \geq 5.$$

- (ii) What are the **initial conditions** which together with the recurrence relation in (i) determine  $f_n$  for all  $n$ . **Do NOT solve the recurrence relation.**

$$f_1 = 1, f_2 = 2, f_3 = 3, f_4 = 6.$$

8. [12 points] Find an **explicit formula** for the general term  $a_n$  of the sequence  $a_1, a_2, \dots, a_n, \dots$  that satisfies:

$$a_n = a_{n-1} + 6a_{n-2}, n \geq 2 \text{ with } a_0 = 3, a_1 = 1.$$

Characteristic equation is  $x^2 - x - 6 = 0$  which has roots 3 and  $-2$ , giving the general solution  $a_n = c3^n + d(-2)^n$ . Using the initial conditions we get  $c = 7/5$  and  $d = 8/5$ .

9. [10 points] **How many permutations** are there of the 9 letters Y,O,U,A,R,E,H,I,T that contain at least one of the words YOU, ARE, HIT?

Let  $A$  be those permutations that contain YOU, let  $B$  be those containing ARE, and let  $C$  be those containing HIT. We want  $A \cup B \cup C$ . Using the Inclusion-Exclusion Principle we get

$$3 \cdot 7! - 3 \cdot 5! + 3!.$$