## Math 104: Homework 11 (due April 29)

- 1. Ross exercise 34.3
- 2. Ross exercise 34.12
- 3. Ross exercise 36.6
- 4. Find a sequence of integrable functions  $(f_n)$  on  $\mathbb{R}$  where  $\int_{-\infty}^{\infty} f_n = 1$  for all n, but  $f_n \to 0$  uniformly on  $\mathbb{R}$ .
- 5. (a) By using simple properties of sin *x* and cos *x*, show how to define the function tan :  $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right) \rightarrow \mathbb{R}$ . Prove that it is differentiable, strictly increasing, and neither bounded above nor below.
  - (b) By using inverse function theorems, define  $\tan^{-1} : \mathbb{R} \to (-\frac{\pi}{2}, \frac{\pi}{2})$  and show that

$$(\tan^{-1})'(x) = \frac{1}{1+x^2}.$$

(c) Prove that for |x| < 1,

$$\tan^{-1} x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{2n+1}.$$

(d) By making use of Abel's theorem, or otherwise, show that

$$\frac{\pi}{4} = \sum_{n=0}^{\infty} \frac{(-1)^n}{2n+1}.$$

- 6. Let  $I_n = \int_0^{\pi/2} \sin^n x \, dx$ .
  - (a) Prove that  $I_0 = \frac{\pi}{2}$  and  $I_1 = 1$ .
  - (b) Use integration by parts to prove that  $(n + 1)I_n = (n + 2)I_{n+2}$  for all  $n \ge 0$ .
  - (c) Prove that  $I_{2m+1} \leq I_{2m} \leq (1 + \frac{1}{2m})I_{2m+1}$  for all  $m \in \mathbb{N}$ , and hence that  $I_{2m}/I_{2m+1} \rightarrow 1$  as  $m \rightarrow \infty$ .
  - (d) Prove that for  $m \in \mathbb{N}$ ,

$$\frac{\pi}{2} = \frac{2}{1} \frac{4}{3} \frac{6}{5} \dots \frac{2m}{2m-1} I_{2m}, \qquad 1 = \frac{3}{2} \frac{5}{4} \frac{7}{6} \dots \frac{2m+1}{2m} I_{2m+1},$$

and hence that

$$\frac{\pi}{2} = \lim_{m \to \infty} \frac{2}{1} \frac{2}{3} \frac{4}{3} \frac{4}{5} \frac{6}{5} \frac{6}{7} \dots \frac{2m}{2m-1} \frac{2m}{2m+1}$$