Math 104: Homework 9 (due April 15)

- 1. Ross exercise 28.4
- 2. Suppose that f_n is a sequence of real-valued functions defined on an interval [a, b] that converges uniformly to a function f. Let $x_0 \in [a, b]$, and suppose that

$$\lim_{x\to x_0}f_n(x)=l_n$$

for all $n \in \mathbb{N}$.

- (a) Prove that l_n is a Cauchy sequence, and hence that it converges to a limit l.
- (b) Prove that $\lim_{x\to x_0} f(x) = l$.
- 3. (a) Let $f : \mathbb{R} \to \mathbb{R}$ be a twice differentiable function, where f(0) = 0 and $f''(x) \ge 0$ for all x > 0. Prove that f(x)/x is increasing for x > 0.
 - (b) If $f : \mathbb{R} \to \mathbb{R}$ is twice differentiable, f(0) = 0 and if f(x)/x is increasing for all x > 0, show that $f''(x) \ge 0$ for some x > 0, but not necessarily for all x > 0. [*Hint: consider* $f(x) = x(1 e^{-x})$.]
- 4. Ross exercise 29.17
- 5. Let $f(x) = |x|^3$. Compute f'(x), f''(x), and show that $f^{(3)}(0)$ does not exist.
- 6. Ross exercise 30.1
- 7. Suppose that *f* is differentiable at a point *a*. Define

$$L_{1}(a,h) = \frac{f(a+h) - f(a-h)}{2h},$$

$$L_{2}(a,h) = \frac{-f(a+2h) + 8f(a+h) - 8f(a-h) + f(a-2h)}{12h}$$

- (a) Prove that $\lim_{h\to 0} L_i(a,h) = f'(a)$ for i = 1, 2.
- (b) Consider the case when $f(x) = x^5$. How does $|L_i(a, h) f'(a)|$ behave as $h \to 0$ for i = 1, 2? Is there a difference in the rate of convergence?