

# COMPUTABILITY-THEORETIC AND PROOF-THEORETIC ASPECTS OF PARTIAL AND LINEAR ORDERINGS

RODNEY G. DOWNEY  
DENIS R. HIRSCHFELDT  
STEFFEN LEMPP  
REED SOLOMON

ABSTRACT. Szpilrajn's Theorem states that any partial order  $\mathcal{P} = \langle S, <_{\mathcal{P}} \rangle$  has a linear extension  $\mathcal{L} = \langle S, <_{\mathcal{L}} \rangle$ . This is a central result in the theory of partial orderings, allowing one to define, for instance, the dimension of a partial ordering. It is now natural to ask questions like "Does a well-partial ordering always have a well-ordered linear extension?" Variations of Szpilrajn's Theorem state, for various (but not for all) linear order types  $\tau$ , that if  $\mathcal{P}$  does not contain a subchain of order type  $\tau$ , then we can choose  $\mathcal{L}$  so that  $\mathcal{L}$  also does not contain a subchain of order type  $\tau$ . In particular, a well-partial ordering always has a well-ordered extension.

We show that several effective versions of variations of Szpilrajn's Theorem fail, and use this to narrow down their proof-theoretic strength in the spirit of reverse mathematics.

## 1. INTRODUCTION

The results of the present paper come from a fruitful interaction of combinatorics and logic. The context of the investigations is the attempt to understand the *effective* content and *proof-theoretical* content of classical mathematics. Before we discuss our results and their ramifications in detail, we give a brief outline of these two programs.

The study of the effective content of mathematics is that part of the work of mathematical logic that seeks to understand and classify the underlying algorithmics inherent in mathematics. Up until the beginning of the 20th century, virtually all mathematics was algorithmic, in the sense that if one claimed that a certain object existed, one gave a computable procedure to generate the object. It was Hilbert in his famous twenty-three problems who asked, essentially, if one

---

1991 *Mathematics Subject Classification.* 03B30, 03C57, 03D45, 06A06.

*Key words and phrases.* partial and linear orderings, well-founded, linear extension, reverse mathematics.

This research was carried out while the first and second authors were visiting the third and fourth authors at the University of Wisconsin. The first and second authors' research was partially supported by the Marsden Fund of New Zealand. The third author's research is partially supported by NSF grant DMS-9732526. The fourth author is supported by an NSF postdoctoral fellowship.

could build a machine to generate all the theorems of, for example, Peano arithmetic. This consideration gave rise to Gödel's powerful incompleteness theorems and, indirectly, to computer science through the work of Turing, von Neumann, and others. Another classical example of such questions was Dehn's [De12] word, conjugacy, and isomorphism problems in finitely presented group theory, which led to the formation of the subject of combinatorial group theory.

In such studies one asks questions like "If one is (computably) *given* a structure  $\mathcal{A}$ , does this guarantee that one can effectively generate a related structure  $\mathcal{B}$  of a particular kind?" A pretty example can be obtained from the work of Rabin, Fröhlich and Shepherdson, and Metakides and Nerode. Rabin [Ra60] demonstrated that if one is computably given a field  $\langle F, \times, +, ^{-1}, 0, 1 \rangle$  (so that  $F$  is a computable set coded by the natural numbers, upon which the normal field operations are computable) then one can effectively find a computable algebraic closure. Fröhlich and Shepherdson [FS56] showed that one can be given two computable algebraic closures of the same computable field which are not computably the same. This is interesting because the *usual* method of generating algebraic closures is to adjoin roots, and this necessarily specifies a unique computable closure. So, in particular, *Rabin's theorem must use a different method of constructing algebraic closures.* Metakides and Nerode [MN79] explained the phenomenon by proving that a computable field has a computably unique computable algebraic closure iff it has a (*separable*) *splitting algorithm*, which means, roughly speaking, that a computable field has a computably unique computable algebraic closure iff one can decide if a given polynomial over the field is irreducible, and hence perform the usual root adjoining process computably.

What does all this tell us? Firstly, we see that there is a demonstrably different way of constructing algebraic closures. This is typical: Clarifying levels of effectiveness involves far greater algebraic or analytic understanding of the structures under consideration. Secondly, we obtain a precise algorithmic equivalence between two processes: adjoining roots and constructing isomorphisms. So, aside from the intrinsic logical interest, we obtain significant insight into the classical algebra. We refer the reader to the *Handbook of Recursive Mathematics* [EGNR98] for more details.

Hand in hand with the above line of research is the attempt to understand the proof-theoretical strength of theorems of classical mathematics. One program here is the "reverse mathematics" of Friedman and Simpson. The idea is to ask whether, given a theorem, one can prove its equivalence to some axiomatic system, with the aim of determining what proof-theoretical resources are necessary for the theorems of mathematics. (A very old example of this line of investigation is Euclid's question of the necessity of the parallel axiom.)

One modern incarnation of this type of analysis comes from the fragment of mathematics living in second-order arithmetic. Second-order arithmetic is a system strong enough to encompass most of classical mathematics. Its underlying language is a two-sorted one with variables for numbers  $(x, y, z, \dots)$  and for sets of numbers  $(X, Y, \dots)$  with the usual logical connectives and quantifiers, together with the normal Peano axioms for number variables (e.g.  $n + 1 = m + 1 \rightarrow m = n$ ), the

induction scheme

$$(0 \in X \wedge \forall x(x \in X \rightarrow x + 1 \in X)) \rightarrow \forall n(n \in X),$$

plus what are called *comprehension schemes*, which assert, roughly speaking, that if we specify an object  $X$  by a formula  $\varphi$  of a particular type, with  $X$  not occurring freely in  $\varphi$ , then the object exists. More formally, a comprehension scheme for a class of formulas is the collection of axioms stating that

$$\exists X \forall n(n \in X \leftrightarrow \varphi(n))$$

for each formula  $\varphi$  in the given class such that  $X$  does not occur freely in  $\varphi$ .

The fundamental idea of reverse mathematics is to calibrate the proof-theoretical strength of a classical theorem by classifying how much comprehension is needed to establish the existence of the structures needed to prove the theorem. That is, we “reverse” the theorem to derive some sort of comprehension scheme. The calibrating measure is that of the allowable “logical complexity” of the  $\varphi$ ’s. Typically, this complexity might be the allowable quantifier depth and type of definition of  $\varphi$  when defined over some quantifier-free formula, although other measures are used as we will see below.

More precisely, a formula  $\varphi$  is called  $\Sigma_0^0$ , or  $\Pi_0^0$ , if it has no unbounded quantifiers. For example, the formula asserting the fact that “ $x$  is prime”, i. e.,

$$\text{Prime}(x) \equiv \forall y \leq x \forall z \leq x (y \cdot z = x \rightarrow y = 1 \vee z = 1),$$

is an example of a  $\Sigma_0^0$  formula. We now adjoin unbounded quantifiers and measure the complexity according to the number of alternations of quantifiers: we say that a formula  $\varphi$  is  $\Sigma_{n+1}^0$  iff there is a  $\Pi_n^0$  formula  $\psi$  such that  $\varphi(x)$  holds iff  $\exists y \psi(x, y)$  and similarly we define  $\Pi_{n+1}^0$  with the roles of  $\Sigma$  and  $\Pi$  reversed. Finally, a formula that is both  $\Sigma_n^0$  and  $\Pi_n^0$  is called  $\Delta_n^0$ . The superscript “0” refers to the fact that there are no *set* quantifiers. Saying that a function is continuous is  $\Pi_3^0$  with the normal  $\epsilon$ - $\delta$  definition. If a formula is  $\Sigma_n^0$  or  $\Pi_n^0$  for some  $n$  we say that it is *arithmetical*. We obtain a similar hierarchy if we allow set quantification by putting a superscript “1” and measuring the number of alternations of set quantifiers over an arithmetical matrix. Thus, for instance, a formula  $\varphi$  of the form  $\exists X \forall Y \psi(X, Y, n)$  with  $\psi$  arithmetical is said to be  $\Sigma_2^1$ , since it begins existentially and has one alternation of set quantifiers.

After all these definitions we can formulate the reverse mathematics program initiated by H. Friedman and Simpson. The goal of this program is to find the minimal “set-theoretical” axioms needed to prove theorems in “ordinary mathematics” by finding “set-theoretical” axioms in second-order arithmetic which not only prove the theorem in “ordinary mathematics” but such that the theorem can also prove the axioms (over some weaker axiom system, typically the axiom system  $RCA_0$  of recursive (i. e.,  $\Delta_1^0$ -)comprehension, together with the basic axioms of a discretely ordered semiring, as above, and  $\Sigma_1^0$ -induction, as we define below.) In his address to the International Congress of Mathematicians, Friedman [Fr74] identified five

systems of second-order arithmetic specified by starting with the basic axioms of a discretely ordered semiring and the  $\Sigma_1^0$ -induction scheme

$$(\varphi(0) \wedge \forall x(\varphi(x) \rightarrow \varphi(x + 1))) \rightarrow \forall n(\varphi(n))$$

where  $\varphi$  is  $\Sigma_1^0$ , and then classifying the types of *allowable comprehension*. The base system is called  $RCA_0$  and allows only  $\Delta_1^0$  comprehension. The next system,  $WKL_0$ , includes the base system plus the comprehension scheme which says that every infinite binary tree has an infinite branch. The third system,  $ACA_0$ , allows for comprehension of sets described by arithmetic formulas. The fourth,  $ATR_0$ , is slightly technical to state, but is equivalent to the statement that any two countable well-orderings are comparable. And, finally, there is the system called  $\Pi_1^1\text{-}CA_0$  which allows comprehension over  $\Pi_1^1$  formulas. (Naturally, there are other possible systems such as  $\Pi_2^1\text{-}CA_0$ , which will, in fact, be relevant to our studies here.)

An important fact is that virtually all of classical “non-set-theoretical” mathematics can be carried out in  $\Pi_1^1\text{-}CA_0$ . It is a remarkable fact that almost all of the classical theorems of mathematics are equivalent to one of the five comprehension schemes above. Pursuing our field example, we note that Friedman, Simpson, and Smith [FSS83] re-interpreted and extended the computability results of Rabin-Fröhlich-Shepherdson-Metakides-Nerode mentioned above to show that the statement “Every countable field has an algebraic closure” is provable in  $RCA_0$ , whereas the uniqueness of the closure is equivalent to  $WKL_0$ .

The existence of a prime ideal in a countable commutative ring with 1 is equivalent to  $WKL_0$ , whereas the existence of a maximal ideal is equivalent to  $ACA_0$  (again meaning that *another* construction of the prime ideal needs to be used than the usual one, which first uses Zorn’s Lemma to construct a maximal ideal and then argues that it is prime). Finally, the existence of an Ulm resolution for a reduced abelian  $p$ -group is equivalent to  $ATR_0$ , and the existence of a decomposition of a countable abelian group into a maximal divisible subgroup and a reduced group is equivalent to  $\Pi_1^1\text{-}CA_0$ . We refer the reader to Simpson [Si99] for more details.

Again we can ask “What is the point of all this?” At one level, we can mention the greater insight one obtains from calibrating the precise resources needed to prove a theorem. We can, in some sense, quantify the intuition that some theorems are “harder” than others. A beautiful example of this is the work on “fast growing Ramsey functions”. One result in this area is the celebrated Paris-Harrington version of the finite Ramsey Theorem [PH77], which is not provable in Peano Arithmetic. Here one shows that the theorem is equivalent to  $ACA_0$ , and hence *although the theorem is concerned only with finite sets, any proof must nevertheless use infinite sets*. An even more striking example of this phenomenon is the work of Friedman, Robertson, and Seymour [FRS87], who proved that the Graph Minor Theorem (even for graphs of bounded tree-width) is *not* provable in  $\Pi_1^1\text{-}CA_0$  and hence the very complicated iterated minimal bad sequence arguments are, in some sense, necessary. (Actually, we remark that the original proof of the Graph Minor Theorem for graphs of bounded tree-width used Friedman’s version of Kruskal’s theorem with the “gap condition”, which was specifically designed to construct stronger incompleteness in Peano Arithmetic as part of the reverse

mathematical program, so that the metamathematical considerations had a huge classical spin-off!)

Another use of reverse mathematics for classical mathematics is in *showing that reasonable classifications are not possible, or at least determining the level that any classification must have*. To make this more precise, let us turn to the area of this paper, in which we will be analyzing extensions of partial orderings to linear ones. Already, we know that this area should be full of metamathematical complexities because of the work of Slaman and Woodin [SW98]. They answered a question of Loś, who had asked for *a classification of those partial orderings with a dense linear extension*. They showed that the collection is not Borel, that is, not  $\Pi_1^1$ , and hence admits *no reasonable classification*.

For the logician, we remark that from a model-theoretic point of view, partial and linear orderings are badly behaved: The existence of an infinite chain necessarily implies instability of the first-order theory. From a computability-theoretic point of view, however, partial and linear orderings are very interesting as they allow a wide variety of codings (see, e. g., Downey [Do98]).

In this paper, we prove some computability-theoretic results, as well as some corollaries for reverse mathematics, on partial and linear orderings.

The starting point of our investigations is

**Szpilrajn’s Theorem** (Szpilrajn [Sz30]). *Any partial order  $\mathcal{P} = \langle S, <_P \rangle$  has a linear extension  $\mathcal{L} = \langle S, <_L \rangle$ .*

We note that Szpilrajn’s Theorem is easily seen to be effective (see Downey [Do98, Observation 6.1]).

Given a property  $P$  of partial orderings, it is natural to ask whether  $\mathcal{P}$  satisfying property  $P$  implies that  $\mathcal{L}$  can be chosen to satisfy property  $P$  as well. Call a linear order type  $\tau$  *extendible*<sup>1</sup> if any partial order  $\mathcal{P} = \langle S, <_P \rangle$  which does not contain a subchain of order type  $\tau$  has a linear extension  $\mathcal{L} = \langle S, <_L \rangle$  which also does not contain a subchain of order type  $\tau$ .

The extendibility of various linear order types was studied extensively by Bonnet, Corominas, Fraïssé, Jullien, and Pouzet in France, as well as independently by Galvin, Kostinsky, and McKenzie in the United States. A complete characterization of the countable extendible linear order types was obtained by Bonnet [Bo69]. In his thesis [Ju69], Jullien obtained a characterization of all countable *weakly extendible* linear order types  $\tau$  (i. e., those  $\tau$  such that any *countable* partial ordering  $\mathcal{P} = \langle S, <_P \rangle$  not containing a chain of order type  $\tau$  can be extended to a linear ordering  $\mathcal{L} = \langle S, <_L \rangle$  not containing a chain of order type  $\tau$ ). (Interestingly enough, there are indeed countable linear order types which are weakly extendible but not extendible, e. g.,  $\omega + 1$ .)

The easiest example of an infinite extendible linear order type is  $\omega^*$  (i. e., the order type of  $\omega$  under the reverse ordering). This is simply to say that any well-founded partial ordering can be extended to a well-ordering. (Of course, by symmetry, this is equivalent to saying that  $\omega$  is extendible.)

---

<sup>1</sup>In the literature, this is sometimes referred to as “enforceable”. Perhaps a better name might be “omittable”.

By Bonnet [Bo69], and independently by Galvin and McKenzie (unpublished), the countable dense linear order type without endpoints,  $\eta$ , is also extendible: If we call a partial ordering without a densely ordered linear chain *scattered*, then “ $\eta$  is extendible” simply means that any scattered partial ordering can be extended to a scattered linear ordering. (We refer the reader to the survey papers by Bonnet and Pouzet [BP82] and by Downey [Do98] for more background on linear extensions of partial orderings, and on computability-theoretic aspects of linear orderings, respectively.)

In the present paper, we analyze the extendibility of the order types  $\omega^*$ ,  $\eta$ , and  $\zeta$  (the order type of the integers) in computability-theoretic and proof-theoretic terms. In particular, we study the extendibility of these three order types along the lines of the program of *reverse mathematics* as outlined above. The axiom systems we will use here are  $WKL_0$ ,  $ACA_0$ ,  $ATR_0$ , and  $\Pi_2^1\text{-}CA_0$ .

We note that there is a strong connection between results in reverse mathematics and effective versions of classical theorems. E. g., loosely speaking, a classical theorem is provable in  $RCA_0$  alone iff the classical theorem holds effectively. Since, in the below, we show that the extendibility of the three linear order types  $\omega^*$ ,  $\eta$ , and  $\zeta$  fails effectively in a very strong sense, we establish lower bounds for the proof-theoretic strength of their extendibility.

Consider first the extendibility of  $\omega^*$ , i. e., the fact that, classically, any well-founded partial ordering has a well-ordered linear extension. Surprisingly, this result holds effectively if the partial ordering is assumed to be *classically well-founded* (i. e., there is no infinite descending sequence) by Rosenstein and Kierstead, but *not* if the partial ordering is only assumed to be *computably well-founded* (i. e., there is no *computable* infinite descending sequence) by Rosenstein and Statman. On the other hand, by Rosenstein, any computable, computably well-founded partial ordering has a computably well-ordered linear extension which is computable in  $\mathbf{0}'$ , the Turing degree of the halting problem. (See Rosenstein [Ro84] for all these results, and Rosenstein [Ro82] for more background.)

We sharpen these results as follows:

**Theorem 1.** (1) “ $\omega^*$  is extendible” is provable in  $ACA_0$ .

(2) “ $\omega^*$  is extendible” proves  $WKL_0$  over  $RCA_0$ .

(3) “ $\omega^*$  is extendible” is not provable in  $WKL_0$ .

The exact proof-theoretic strength of the extendibility of  $\omega^*$  thus remains open. We remark that the *particular proof* of the extendibility of  $\omega^*$  from  $ACA_0$  which we present below reverses to  $ACA_0$ .

Consider next the extendibility of  $\eta$ , i. e., the fact that, classically, any scattered partial ordering has a scattered linear extension. By Rosenstein [Ro82, Ro84], any computable, computably scattered partial ordering has a computably scattered linear extension which is computable in  $\mathbf{0}'$ .

**Theorem 2.** (1) “ $\eta$  is extendible” is provable in  $\Pi_2^1\text{-}CA_0$ .<sup>2</sup>

---

<sup>2</sup>Upon hearing of our result, Howard Becker (personal communication) has found a proof of “ $\eta$  is extendible” from  $\Pi_1^1\text{-}CA_0$ . We show below how our proof can be modified to give Becker’s result.

(2) “ $\eta$  is extendible” is not provable in  $WKL_0$ .

In particular, our proof of Theorem 2 answers long-standing open questions from Rosenstein [Ro84] by the following

**Theorem 2A.** *There is a classically scattered, computable partial ordering such that every computable linear extension has a computable densely ordered subchain.*

However, the exact proof-theoretic strength of the extendibility of  $\eta$  remains open.

We finally classify precisely the proof-theoretic strength of the extendibility of  $\zeta$ , and add to the small collection of classical theorems equivalent to  $ATR_0$  over  $RCA_0$ :

**Theorem 3.** *“ $\zeta$  is extendible” is equivalent to  $ATR_0$  over  $RCA_0$ .*

The rest of the paper is devoted to the proofs of these theorems. Henceforth, we assume that the reader is familiar with the rudiments of reverse mathematics, referring to Simpson [Si99] where necessary, and assume that the reader is familiar with the rudiments of computability theory, as found in an initial segment of Soare [So86] or Rogers [Ro67].

## 2. THE PROOF OF THEOREM 1

To prove part (1) of Theorem 1, simply observe that the proof of Kierstead and Rosenstein [Ro84] (see also [Do98, p. 909]) can be used: Fix a partial ordering  $\mathcal{P} = \langle N, <_P \rangle$ . (Note here that we may assume without loss of generality that the universe of  $\mathcal{P}$  is  $N$ , the set of integers in the sense of the model  $\mathcal{N}$  of  $ACA_0$ .) We define a linear extension  $\mathcal{L} = \langle N, <_L \rangle$  by approximations  $\mathcal{L}_s = \mathcal{L} \upharpoonright [0, s]$  as follows: At stage 0, set  $L_0 = \{0\}$ . At stage  $s + 1$ , fix the  $<_L$ -least element  $a \in L_s$  such that  $s + 1 <_P a$  and let  $s + 1$  be the immediate  $\mathcal{L}_{s+1}$ -predecessor of  $a$ . (If  $a$  fails to exist, make  $s + 1$  the greatest element of  $\mathcal{L}_{s+1}$ . So the idea is to insert  $s + 1$  into  $\mathcal{L}_s$  at the rightmost place consistent with  $<_P$ .)

Now suppose  $\{a_s\}_{s \in N}$  is a  $<_L$ -descending sequence (coded in the model  $\mathcal{N}$ ). By Ramsey’s Theorem, we may assume without loss of generality that  $\{a_s \mid s \in \mathcal{N}\}$  is a  $<_P$ -antichain. (Note that by Jockusch [Jo72], Ramsey’s Theorem can be used inside the model  $\mathcal{N}$ .) We may also assume that  $\{a_s\}_{s \in N}$  is  $<$ -increasing (in the usual ordering of  $N$ ). We can now construct (inside the model  $\mathcal{N}$ ) a subsequence  $\{a_{j_t}\}_{t \in N}$  and a  $<_P$ -descending sequence  $\{b_t\}_{t \in N}$  with  $a_{j_{t+1}} <_P b_t < a_{j_t}$  for all  $t$  as follows: Set  $a_{j_0} = a_0$ . Given  $t$ , and since the  $a$ ’s form a  $<_P$ -antichain, for each  $s > j_t$  there is a ( $<$ -least) element  $d_s$  with  $a_s <_P d_s <_L a_{j_t}$ . By the construction (since we always place elements “rightmost” in  $\mathcal{L}$ ), we have  $d_s < a_{j_t}$ . Among these  $d_s$ , fix the  $d$   $<$ -least such that  $d = d_s$  for  $\mathcal{N}$ -infinitely many  $s$ , and thin out the sequence of  $a$ ’s to only contain  $a_s$  with  $d_s = d$  for the definition of  $b_{t'}$  ( $t' > t$ ). We then set  $b_t = d$  and  $a_{j_{t+1}} = \text{some } a_s <_P d$ . Using the minimality in the choice of  $d$ , we can then argue that  $\{b_t\}_{t \in N}$  is a  $<_P$ -descending sequence, contradicting the wellfoundedness of  $\mathcal{P}$ .

To prove part (2) of Theorem 1, note that the proof of Rosenstein and Statman [Ro84] (see also [Do98, p. 910]) can be adapted: Fix an infinite tree  $T \subseteq 2^{<\omega}$  (coded

in the model  $\mathcal{N}$  of  $RCA_0 + \text{“}\omega^* \text{ is extendible”}$ ), and view it as partially ordered by reverse inclusion (which we denote by  $<_P$ ). It now suffices to show that any linear extension  $<_L$  of  $<_P$  has an infinite  $<_L$ -descending chain, since then “ $\omega^*$  extendible” implies that  $T$  has an infinite  $<_P$ -descending chain, i. e., an infinite path.

So fix an arbitrary linear extension  $<_L$  of  $<_P$ . We define an infinite  $<_L$ -descending sequence  $\{c_s\}_{s \in \mathbb{N}}$  as follows: Let  $c_0$  be the root of  $T$ . Given  $s$ , let  $C_s = \{c_0, \dots, c_s\}$ , and let  $D_s$  be the set of immediate  $T$ -successors (i. e., immediate  $<_P$ -predecessors) of elements of  $C_s$ . Then we let  $c_{s+1}$  be the  $<_L$ -maximal element of  $D_s - C_s$ . It is now easy to check that the maximality condition in the choice of  $c_s$  ensures that they form a  $<_L$ -descending sequence as desired.

To prove part (3) of Theorem 1, we will show that, given a sequence  $X_0 \leq_T X_1 \leq_T \dots$  of uniformly low, uniformly  $\Delta_2^0$ -sets, there is a computable partial ordering  $\mathcal{P} = \langle S, <_P \rangle$  such that, for any infinite  $<_P$ -descending sequence  $\{c_n\}_{n \in \mathbb{N}}$ , there is some  $i$  such that  $X_i \oplus \{c_n\}_{n \in \mathbb{N}}$  can compute the halting set  $K$ ; and such that for any  $i$ , any  $X_i$ -computable linear extension  $<_L$  of  $<_P$  contains an infinite  $<_L$ -descending chain Turing computable in  $<_L$ . Since, by Jockusch and Soare [JS72] and Simpson [Si99, Theorem VIII.2], there is a model of  $WKL_0$  whose second-order part consists of all sets in the Turing ideal generated by a sequence  $X_0 \leq_T X_1 \leq_T \dots$  of uniformly low, uniformly  $\Delta_2^0$ -sets (which thus in particular does not contain the halting set  $K$ ), this implies that  $WKL_0$  does not imply the extendibility of  $\omega^*$ .

The construction of  $\mathcal{P}$  is a finite-injury priority argument. We construct  $\mathcal{P}$  as the disjoint union of sub-partial orderings  $\mathcal{P}_{i,e}$  (for  $e, i \in \omega$ ) such that each  $\mathcal{P}_{i,e}$  is a connected component of  $\mathcal{P}$  (when viewed as a directed graph). Each  $\mathcal{P}_{i,e}$  will be devoted to showing that if the  $e$ th binary  $X_i$ -computable relation  $\mathcal{L}_{i,e} = L_e^{X_i}$  is a linear extension of  $<_P$  then it has an infinite descending sequence inside  $P_{i,e}$  (the domain of  $\mathcal{P}_{i,e}$ ) which is computable in  $\mathcal{L}_{i,e}$ . At the same time, we have to show that any infinite  $<_P$ -descending sequence inside  $P_{i,e}$  can compute the halting set  $K$ . (Note here that any  $<_P$ -descending sequence in  $\mathcal{P}$  must be completely contained in a single  $P_{i,e}$ .)

We can thus fix indices  $i$  and  $e$  and concentrate on the construction of the subordering  $\mathcal{P}_{i,e}$ . (The constructions for the various  $i$  and  $e$  can be fit together using a computable partition of  $\omega$ . Since if  $\mathcal{L}_{i,e}$  is not a linear extension of  $<_P$  on  $P_{i,e}$ ,  $\mathcal{P}_{i,e}$  may turn out to be finite, this computable partition cannot be fixed beforehand but must be constructed simultaneously with the components  $\mathcal{P}_{i,e}$ .)

*Remark.* Since we are assuming that  $X_i$  is low (uniformly in  $i$ ), we may assume that  $L_e^{X_i}$  is either total (and thus can be approximated effectively à la Shoenfield’s Limit Lemma), or else  $L_e^{X_i}$  is finite. For simplicity, we will suppress the details of this approximation.

We will now construct a partial ordering  $\mathcal{P}$  such that

- (1) if  $\mathcal{L}_{i,e} \upharpoonright P_{i,e}$  is a linear extension  $<_L$  of  $<_P$  on  $P_{i,e}$ , then we have an infinite  $\mathcal{L}_{i,e}$ -computable  $<_L$ -descending sequence  $\{c_n\}_{n \in \mathbb{N}}$  of elements of  $P_{i,e}$ , and
- (2) from the join of  $X_i$  with any infinite set of elements of  $P_{i,e}$ , each of which has infinitely many elements  $<_P$ -below it, we can compute  $K$ .

We first illustrate our construction by showing how to code whether  $0 \in K$  into



any infinite  $\langle_P$ -descending chain in  $P_{i,e}$  while at the same time, computably in  $\langle_L$ , fixing an element  $c_0 \rangle_P$  infinitely many elements of  $P_{i,e}$ . We start with seven elements  $a, a_1, a_2, \dots, a_6$ , declaring

$$a_3 \langle_P a_2 \langle_P a_1 \langle_P a \text{ and } a_6 \langle_P a_5 \langle_P a_4 \langle_P a$$

with no other comparabilities. We call  $a_2$  and  $a_5$  the *0-critical elements* and wait for  $\mathcal{L}_{i,e}$  to decide whether or not  $a_2 \rangle_L a_5$ . If  $a_2 \rangle_L a_5$  then we let  $a_1$  be the first element  $c_0$  of our  $\langle_L$ -descending chain, and we build the rest of  $P_{i,e}$  in the  $\langle_P$ -interval  $(a_2, a_1)$  until 0 enters  $K$ ; when 0 enters  $K$  then we switch to building the rest of  $P_{i,e}$  in the  $\langle_P$ -interval  $(a_6, a_5)$ . Symmetrically, if  $a_5 \rangle_L a_2$  then we let  $a_4$  be the first element  $c_0$  of our  $\langle_L$ -descending chain, and we build the rest of  $P_{i,e}$  in the  $\langle_P$ -interval  $(a_5, a_4)$  until 0 enters  $K$ ; when 0 enters  $K$  then we switch to building the rest of  $P_{i,e}$  in the  $\langle_P$ -interval  $(a_3, a_2)$ . If  $\langle_L$  eventually decides whether or not  $a_2 \rangle_L a_5$ , then, since, if  $0 \in K$ , we will eventually see 0 enter  $K$ , only one of the four  $\langle_P$ -intervals  $[a_2, a_1)$ ,  $[a_3, a_2)$ ,  $[a_5, a_4)$ , and  $[a_6, a_5)$  will be infinite (we will call this  $\langle_P$ -interval the *0-active interval*), and we will have put only elements  $\langle_L$ -above this  $\langle_P$ -interval in our  $\langle_L$ -descending chain. (If  $\mathcal{L}_{i,e}$  does not converge on whether or not  $a_2 \rangle_L a_5$  then  $P_{i,e}$  will be finite.) Note that from  $\langle_L$  we can compute the first element  $c_0$  of our  $\langle_L$ -descending chain. And any infinite  $\langle_P$ -descending chain must contain either elements  $\leq_P a_1$  or elements  $\leq_P a_4$  (but not both), and so from this and the  $\langle_L$ -ordering of the 0-critical elements, we can compute whether  $0 \in K$ . (We note here that the definition of the special element  $c_0$  does not change when 0 enters  $K$ , but the definition of the 0-critical interval does. The same will be true in the full construction in the next paragraph, and there the definition of the  $n$ -critical elements will also change depending on what elements  $\langle n$  enter  $K$ .)

The full construction simply nests the above: The previous paragraph describes the definition of the 0-active interval. Given the  $n$ -active interval  $[b, c)$ , we create six new elements  $b_1, \dots, b_6$  in it, declaring

$$b_3 \langle_P b_2 \langle_P b_1 \langle_P b \text{ and } b_6 \langle_P b_5 \langle_P b_4 \langle_P b$$

with no other comparabilities in the  $\langle_P$ -interval  $[b, c)$ . We call  $b_2$  and  $b_5$  the  $(n+1)$ -critical elements and wait for  $\mathcal{L}_{e,i}$  to decide whether or not  $b_2 \rangle_L b_5$ . If  $b_2 \rangle_L b_5$  then we let  $b_1$  be the element  $c_{n+1}$  of our  $\langle_L$ -descending chain, and we build the rest of the  $\langle_P$ -interval  $[b, c)$  in the  $\langle_P$ -interval  $(b_2, b_1)$  until  $n+1$  enters  $K$ ; when  $n+1$  enters  $K$  then we switch to building the rest of the  $\langle_P$ -interval  $[b, c)$  in the  $\langle_P$ -interval  $(b_6, b_5)$ . Symmetrically, if  $b_5 \rangle_L b_2$  then we let  $b_4$  be the element  $c_{n+1}$  of our  $\langle_L$ -descending chain, and we build the rest of the  $\langle_P$ -interval  $[b, c)$  in the  $\langle_P$ -interval  $(b_5, b_4)$  until  $n+1$  enters  $K$ ; when  $n+1$  enters  $K$  then we switch to building the rest of the  $\langle_P$ -interval  $[b, c)$  in the  $\langle_P$ -interval  $(b_3, b_2)$ . If  $\langle_L$  eventually decides whether or not  $b_2 \rangle_L b_5$ , then, since, if  $n+1 \in K$ , we will eventually see  $n+1$  enter  $K$ , only one of the four  $\langle_P$ -intervals  $[b_2, b_1)$ ,  $[b_3, b_2)$ ,  $[b_5, b_4)$ , and  $[b_6, b_5)$  will be infinite (we will call this  $\langle_P$ -interval the  $(n+1)$ -active interval), and we will have put only elements  $\langle_L$ -above this  $\langle_P$ -interval in our  $\langle_L$ -descending chain. (If  $\mathcal{L}_{i,e}$  does not converge on whether  $b_2 \rangle_L b_5$  then  $P_{i,e}$  will be finite.) Note again

that from  $<_L$  we can compute the element  $c_{n+1}$  of our  $<_L$ -descending chain. And any infinite  $<_P$ -descending chain must contain either elements  $\leq_P b_1$  or elements  $\leq_P b_4$  (but not both), and so from this and the  $<_L$ -ordering of the  $(n+1)$ -critical elements, we can compute whether  $n+1 \in K$ .

As mentioned in the Remark above, the above nested construction for each  $n$  is controlled by our computable approximation of how  $<_L$  orders the  $n$ -critical elements. If  $<_L$  does not decide the ordering of the  $n$ -critical elements, then  $P_{i,e}$  will be finite; otherwise,  $P_{i,e}$  will be infinite, and for each  $n$ , we will eventually settle on the correct  $n$ -active interval (while making finitely many mistakes before then). In the latter case, it is not hard to check that we are building an infinite  $<_L$ -computable  $<_L$ -descending chain  $\{c_n\}_{n \in \mathbb{N}}$ ; and that any infinite subset of elements of  $P_{i,e}$ , each of which has infinitely many elements  $<_P$ -below it, can compute  $K$  with an additional oracle for  $<_L$ , and thus a fortiori with an additional oracle for  $X_i$ .

This concludes the proof of part (3) of Theorem 1.

### 3. THE PROOF OF THEOREMS 2 AND 2A

To prove part (1)<sup>3</sup> of Theorem 2, simply observe that the proof of Bonnet and Pouzet [BP69] (see also [BP82, p. 140]) can be adapted here: Given a partial ordering  $\mathcal{P} = \langle P, <_P \rangle$ , we say an element  $a$  of  $P$  is  $<_P$ -good if  $\langle P(\leq_P a), <_P \rangle$  has a scattered linear extension; and  $>_P$ -good if  $\langle P(\geq_P a), <_P \rangle$  has a scattered linear extension. Define the  $\eta$ -kernel  $K(\mathcal{P})$  of  $\mathcal{P}$  to be the set of all elements of  $\mathcal{P}$  which are neither  $<_P$ -good nor  $>_P$ -good. (Here  $P(\leq_P a)$  and  $P(\geq_P a)$  are the sets of elements of  $P$  below or above  $a$ , respectively.) It is now easy to check that

$$K(\mathcal{P}) = \{a \in P \mid$$

$$\forall \text{ linear extension } \prec \text{ of } \langle P(\leq a), <_P \rangle \exists \prec\text{-chain } S \text{ (} S \text{ is densely ordered)}$$

$$\text{and}$$

$$\forall \text{ linear extension } \prec \text{ of } \langle P(\geq a), <_P \rangle \exists \prec\text{-chain } S \text{ (} S \text{ is densely ordered)}\}.$$

So  $K(\mathcal{P})$  is  $\Pi_2^1$ -definable, and thus the  $\eta$ -kernel of any partial ordering  $\mathcal{P}$  in our model  $\mathcal{N}$  of  $\Pi_2^1\text{-CA}_0$  also exists in  $\mathcal{N}$ .

We now observe that the collection of  $<_P$ -good elements, and the collection of  $>_P$ -good elements, form an initial segment, or a final segment, of  $\mathcal{P}$ , respectively. We claim that  $\Pi_2^1\text{-CA}_0$  proves  $\mathcal{P}$  has a scattered linear extension iff  $K(\mathcal{P}) = \emptyset$ . First, if  $\mathcal{P}$  has a scattered linear extension, then by restriction, it is clear that every cone in  $\mathcal{P}$  has a scattered linear extension. Therefore,  $K(\mathcal{P}) = \emptyset$ . To establish the other direction, suppose that  $K(\mathcal{P}) = \emptyset$ . Then, for every  $a \in P$ , either  $P(\leq_P a)$  or  $P(\geq_P a)$  has a scattered linear extension. Partition  $P$  into

$$X = \{a \in P \mid P(\leq_P a) \text{ has a scattered linear extension}\}$$

---

<sup>3</sup>At the end of the proof, we show how this proof can be modified to give Becker's stronger result that  $\Pi_1^1\text{-CA}_0$  is sufficient to prove the extendibility of  $\eta$ .

and  $P \setminus X$ . We use this decomposition to build a scattered linear extension of  $\mathcal{P}$ .

First, we build a scattered linear extension of  $(X, \leq_P)$  by arithmetical recursion over  $\mathcal{N}$ . Let  $x_0, x_1, \dots$  be a list of the elements of  $X$  in  $<_{\mathcal{N}}$ -increasing order. By  $\Sigma_2^1$ -choice (which is provable in  $\Pi_2^1\text{-CA}_0$ , see [Si99, Section VII.6]), we can fix a scattered linear extension for each cone  $P(\leq_P x_s)$ ,  $s \in \mathcal{N}$ . Let  $X_s = \{p \in P \mid p \leq_P x_0 \text{ or } \dots \text{ or } p \leq_P x_s\} \subseteq X$ . We define  $L_0$  to be the fixed scattered linear extension of  $X_0$ . Assume that we have a scattered linear extension  $L_s$  of  $X_s$ . If  $x_{s+1} \in X_s$ , then let  $L_{s+1} = L_s$ . Otherwise, let  $Z = P(\leq_P x_{s+1}) \setminus X_s$ , and fix a scattered linear extension  $L_Z$  of  $Z$  by restricting the scattered extension of  $P(\leq_P x_{s+1})$ . Let  $L_{s+1}$  be the linear extension of  $X_{s+1}$  which agrees with  $L_s$  on  $X_s$ , agrees with  $L_Z$  on  $Z$ , and places elements in  $Z$  above everything from  $X_s$ .  $L_{s+1}$  is a scattered linear extension of  $X_{s+1}$  which is an end-extension of  $L_s$ . Combining these orders,  $\cup L_s$  is a scattered linear extension of  $X$ .

It remains to handle  $P \setminus X$ . We list these elements as  $y_0, y_1, \dots$  and let  $Y_s = \{p \in P \mid y_0 \leq_p p \text{ or } \dots \text{ or } y_s \leq_p p\}$ . We use a similar construction to the one above to build linear extensions  $L'_s$  of  $Y_s$  such that  $L'_{s+1}$  extends  $L'_s$  downwards. We combine  $\cup L_s$  with  $\cup L'_s$  by placing all elements of  $P \setminus X$  above all elements of  $X$ . This gives a scattered linear extension of  $P$ , and finishes the claim that  $\mathcal{P}$  has a scattered linear extension if and only if  $K(\mathcal{P}) = \emptyset$ .

We can now prove the extendibility of  $\eta$  from  $\Pi_2^1\text{-CA}_0$  as follows: Fix a partial ordering  $\mathcal{P}$  and assume that it does not have a scattered linear extension. Add to  $\mathcal{P}$  a new least and a new greatest element,  $x_0$  and  $x_1$ , respectively. Let  $D = \{\frac{m}{2^n} \mid 0 \leq m \leq 2^n \text{ and } n \in \omega\}$  be the set of dyadic rationals in  $[0, 1]$ . By induction, we will now construct a subset  $X = \{x_d \mid d \in D \cap (0, 1)\}$  of  $P$  such that  $x_d <_P x_e$  iff  $d < e$ ; so  $X$  is a dense subset of  $\mathcal{P}$  as desired. Suppose  $x_d$  has been defined for all indices with denominator  $< 2^n$ , and fix  $d = \frac{m}{2^n} \in D$  with  $m$  odd. By induction, the  $\mathcal{P}$ -interval  $[x_{\frac{m-1}{2^n}}, x_{\frac{m+1}{2^n}}]$  has no scattered linear extension, so we can choose an element  $x_d$  in its  $\eta$ -kernel  $K([x_{\frac{m-1}{2^n}}, x_{\frac{m+1}{2^n}}])$ . By the definition of  $\eta$ -kernel, the intervals  $[x_{\frac{m-1}{2^n}}, x_d]$  and  $[x_d, x_{\frac{m+1}{2^n}}]$  have no scattered linear extension, and so the induction can continue.<sup>4</sup>

---

<sup>4</sup>Becker's proof modifies ours as follows: Note that our proof has two parts: (i) Any partial ordering  $\mathcal{P}$  has a scattered linear extension if and only if the  $\eta$ -kernel of  $\mathcal{P}$  is nonempty. (ii) If the kernel is nonempty then  $\mathcal{P}$  is not scattered. Now (i) can actually be shown within  $\text{ATR}_0$  in the following version: (i') Any partial ordering  $\mathcal{P}$  has a scattered linear extension which is hyperarithmetical in  $\mathcal{P}$  iff there is no  $a \in P$  which is neither  $<_P$ -good nor  $>_P$ -good, where now an element  $a$  is  $<_P$ -good (or  $>_P$ -good, respectively) if  $\langle P(\leq_P a), <_P \rangle$  (or  $\langle P(\geq_P a), <_P \rangle$ ) has a scattered linear extension *hyperarithmetical* in  $\mathcal{P}$ . Then the sets of  $<_P$ -good (or  $>_P$ -good, respectively) elements are  $\Pi_1^1$ -definable in  $\mathcal{P}$ , and our set  $X$  above can be chosen  $\Delta_1^1$  in  $\mathcal{P}$  by  $\Sigma_1^1$ -Separation (or more precisely, by  $\Pi_1^1$ -Reduction, see Simpson [Si99, V.5.1]) and such that  $X$  is downward closed and contains only  $<_P$ -good elements, whereas  $P - X$  contains only  $>_P$ -good elements. By the Kreisel Selection Theorem (see [Si99, VIII.4.7]), there are now functions  $f : X \rightarrow N$  and  $g : P - X \rightarrow N$  (which are  $\Delta_1^1$  in  $\mathcal{P}$ ) such that for all  $a \in X$ ,  $f(a)$  is the index of a scattered linear extension of  $\langle P(\leq_P a), <_P \rangle$  hyperarithmetical in  $\mathcal{P}$ , and for all  $a \in P - X$ ,  $g(a)$  is the index of a scattered linear extension of  $\langle P(\geq_P a), <_P \rangle$  hyperarithmetical in  $\mathcal{P}$ . Now proceed as in our proof above to obtain the linear extension hyperarithmetical in  $\mathcal{P}$  using  $X$ ,  $f$  and  $g$ . Part (ii) can be shown within  $\Pi_1^1\text{-CA}_0$  as in our original proof.

To prove part (2) of Theorem 2 as well as Theorem 2A, we will show that, given a sequence  $X_0 \leq_T X_1 \leq_T \dots$  of uniformly low, uniformly  $\Delta_2^0$ -sets, there is a computable partial ordering  $\mathcal{P} = \langle S, <_P \rangle$  which is classically scattered (i. e., there is no densely ordered  $<_P$ -subchain of *any* complexity) such that for any  $i$ , any  $X_i$ -computable linear extension  $<_L$  of  $<_P$  contains a densely ordered  $<_L$ -subchain Turing computable in  $<_L$ . Again, since, by Jockusch and Soare [JS72] and Simpson [Si99, VIII.2], there is a model of  $WKL_0$  whose second-order part consists of all sets in the Turing ideal generated by a sequence  $X_0 \leq_T X_1 \leq_T \dots$  of uniformly low, uniformly  $\Delta_2^0$ -sets, this implies that  $WKL_0$  does not imply the extendibility of  $\eta$ .

The construction of  $\mathcal{P}$  is again a finite-injury priority argument. We construct  $\mathcal{P}$  as the disjoint union of sub-partial orderings  $\mathcal{P}_{i,e}$  (for  $e, i \in \omega$ ) such that each  $\mathcal{P}_{i,e}$  is a connected component of  $\mathcal{P}$  (when viewed as a directed graph). Each  $\mathcal{P}_{i,e}$  will be devoted to showing that if the  $e$ th binary  $X_i$ -computable relation  $\mathcal{L}_{i,e} = L_e^{X_i}$  is a linear extension  $<_L$  of  $<_P$  then it has a densely ordered  $<_L$ -subchain inside  $P_{i,e}$  which is computable in  $\mathcal{L}_{i,e}$ . At the same time, we have to show that there is no densely ordered  $<_P$ -subchain inside  $P_{i,e}$ . (Note here that any  $<_P$ -subchain in  $\mathcal{P}$  must be completely contained in a single  $P_{i,e}$ .)

We can thus fix indices  $i$  and  $e$  and concentrate on the construction of the subordering  $\mathcal{P}_{i,e}$ . (The constructions for the various  $i$  and  $e$  can be fit together using a computable partition of  $\omega$ . Since if  $\mathcal{L}_{i,e}$  is not a linear extension of  $<_P$  on  $P_{i,e}$ ,  $\mathcal{P}_{i,e}$  may turn out to be finite, this computable partition cannot be fixed beforehand but must be constructed simultaneously with the components  $\mathcal{P}_{i,e}$ .) The Remark in the proof of part (3) of Theorem 1 also applies here verbatim.

We will now construct a partial ordering  $\mathcal{P}$  such that

- (1) if  $\mathcal{L}_{i,e} \upharpoonright P_{i,e}$  is a linear extension  $<_L$  of  $<_P$  on  $P_{i,e}$ , then we have an  $\mathcal{L}_{i,e}$ -computable densely ordered  $<_L$ -subchain  $C$  in  $P_{i,e}$ , and
- (2) for any element  $x$  of  $P_{i,e}$ , there are either only finitely many elements  $>_P x$ , or only finitely many elements  $<_P x$ .

We first illustrate our construction by showing how to perform a single step: We start with three  $<_P$ -incomparable elements  $a_0, a_1$  and  $a_2$ , calling them the *0-critical elements*, and wait for  $\mathcal{L}_{i,e}$  to decide the  $<_L$ -ordering on these three elements. Possibly relabeling them, we will assume  $a_0 <_L a_1 <_L a_2$ . We then place  $a_1$  into  $C$  and call the interval  $(-\infty, a_0)$  and the interval  $(a_2, \infty)$  the *0-active intervals*. (If  $\mathcal{L}_{i,e}$  does not converge on the ordering of  $a_0, a_1$ , and  $a_2$  then  $P_{i,e}$  will be finite.) Note that from  $<_L$  we can compute the first element of  $C$ . And among the elements  $a_0, a_1$ , and  $a_2$ , at most one is  $<_P$ -comparable to any element of the 0-active intervals.

The full construction simply nests the above: The previous paragraph describes the definition of the 0-active intervals. Given an  $n$ -active  $<_P$ -interval  $(b, c)$ , we create three new pairwise incomparable elements  $b_0, b_1$ , and  $b_2$  in  $(b, c)$ , calling them the  $(n+1)$ -critical elements, and wait for  $\mathcal{L}_{e,i}$  to decide the  $<_L$ -ordering on these three elements. Possibly relabeling them, we will assume  $b_0 <_L b_1 <_L b_2$ . We then place  $b_1$  into  $C$  and call the interval  $(b, b_0)$  and the interval  $(b_2, c)$   $(n+1)$ -active intervals. (If  $\mathcal{L}_{i,e}$  does not converge on the ordering of  $b_0, b_1$ , and  $b_2$  then  $P_{i,e}$  will be finite.) Note that from  $<_L$  we can compute the first element of  $C$  in the interval

$(b, c)$ . And among the elements  $b_0$ ,  $b_1$ , and  $b_2$ , at most one is  $<_P$ -comparable to any element of the  $(n + 1)$ -active intervals.

As mentioned in the Remark above, the above nested construction for each  $n$  is controlled by our computable approximation of how  $<_L$  orders the  $n$ -critical elements. If  $<_L$  does not decide the ordering of the  $n$ -critical elements, then  $P_{i,e}$  will be finite; otherwise,  $P_{i,e}$  will be infinite, and for each  $n$ , we will eventually settle on the correct  $n$ -active intervals (while making finitely many mistakes before then). In the latter case, it is not hard to check that we are building a  $<_L$ -computable densely  $<_L$ -ordered chain; and that for any element  $x$  of  $P_{i,e}$ , there are either only finitely many elements  $>_P x$ , or only finitely many elements  $<_P x$ .

This concludes the proof of part (2) of Theorem 2.

#### 4. THE PROOF OF THEOREM 3

To prove that  $ATR_0$  implies the extendibility of  $\zeta$ , observe that the proof of Jullien [Ju69] (see also [BP82, p. 141], note a typo there:  $\omega + \omega^*$  should be  $\omega^* + \omega$ , i. e.,  $\zeta$ ) can be adapted here: Fix a partial ordering  $\mathcal{P} = \langle S, <_P \rangle$  without any subchain (in our model  $\mathcal{N}$  of  $ATR_0$ ) of order type  $\zeta$ . Call an element  $a$  of  $P$   $<_P$ -good if  $\langle P(\leq_P a), <_P \rangle$  contains no subchain of order type  $\omega^*$ ; and  $>_P$ -good if  $\langle P(\geq_P a), <_P \rangle$  contains no subchain of order type  $\omega$ . By our assumption on  $\mathcal{P}$ , any element of  $P$  is either  $<_P$ -good or  $>_P$ -good. Since  $<_P$ -goodness and  $>_P$ -goodness are both  $\Pi_1^1$ -definable, our model  $\mathcal{N}$  of  $ATR_0$  contains a set  $S$  by  $\Sigma_1^1$ -separation such that any element of  $S$  is  $<_P$ -good, and any element of  $P - S$  is  $>_P$ -good. (We use here that  $ATR_0$  is equivalent to  $\Sigma_1^1$ -separation (see Simpson [Si99, Theorem V.5.1]).)

Since the set of  $<_P$ -good elements is downward closed in  $\mathcal{P}$ , we may assume that  $S$  is downward closed in  $\mathcal{P}$ . Now, by part (1) of Theorem 1,  $ACA_0$  (and thus *a fortiori*  $ATR_0$ ) proves the extendibility of  $\omega$  and  $\omega^*$ ; so we can fix linear extensions  $<_{L_1}$  and  $<_{L_2}$  of  $S$  and  $P - S$ , respectively, which have no subchains of order type  $\omega^*$  and  $\omega$ , respectively. We can now patch  $<_{L_1}$  and  $<_{L_2}$  together (by placing the elements of  $S$  left of the elements of  $P - S$ ) to obtain a linear extension  $<_L$  of  $\mathcal{P}$  without a subchain of order type  $\zeta$ .

We now establish the other direction of the proof of Theorem 3 by two claims:

**Claim 1.**  $RCA_0$  and the extendibility of  $\zeta$  imply  $ACA_0$ .

**Claim 2.**  $ACA_0$  and the extendibility of  $\zeta$  imply  $ATR_0$ .

*Proof of Claim 1.* Fix a model  $\mathcal{N}$  of  $RCA_0 + \text{“}\zeta \text{ is extendible”}$  and an injective function  $f : N \rightarrow N$  in this model. In order to establish  $ACA_0$ , we must show that the range of  $f$  is in this model (see Simpson [Si99, Lemma III.1.3]). Construct a partial ordering  $\mathcal{P}$  in  $\mathcal{N}$  with distinguished elements  $a_n$  and  $b_n$  (for  $n \in N$ ) as follows: If  $n$  is not in the range of  $f$  then  $a_n$  is the first element of an  $\omega$ -chain, and  $b_n$  is the last element of an  $\omega^*$ -chain in  $\mathcal{P}$ . If  $n = f(m)$  then  $a_n$  is the  $m$ th element from the end of an  $\omega^*$ -chain, and  $b_n$  is the  $m$ th element of an  $\omega$ -chain in  $\mathcal{P}$ . Since all  $a_n$  and  $b_n$  are in different connected components of  $\mathcal{P}$  (when viewed as a directed graph),  $\mathcal{P}$  contains no  $\zeta$ -chain, so we can fix a linear extension  $<_L$

without a  $\zeta$ -chain. But then  $n$  is in the range of  $f$  iff  $b_n <_L a_n$ , the range of  $f$  is in the model  $\mathcal{N}$  as desired.

*Proof of Claim 2.* We use again that  $ATR_0$  is equivalent to  $\Sigma_1^1$ -separation. Fix a model  $\mathcal{N}$  of  $ACA_0 + \text{“}\zeta \text{ is extendible”}$  and two disjoint sets  $S_0, S_1 \subseteq N$  which are  $\Sigma_1^1$ -definable over this model. We will separate  $S_0$  and  $S_1$  as follows: Fix two sequences of trees  $\mathcal{L}_n^0$  and  $\mathcal{L}_n^1$  such that for all  $i < 2$  and  $n \in N$ ,  $n \in S_i$  iff  $\mathcal{L}_n^i$  is not well-founded. (Such sequences exist within  $\mathcal{N}$ ; see Simpson [Si99, Theorem V.1.8].) Construct a partial ordering  $\mathcal{P}$  in  $\mathcal{N}$  with distinguished elements  $a_n$  and  $b_n$  (for  $n \in N$ ) as follows:

- (1)  $\mathcal{P}(<_P a_n)$  is isomorphic to  $\mathcal{L}_n^0$ ,
- (2)  $\mathcal{P}(>_P a_n)$  is isomorphic to  $(\mathcal{L}_n^1)^*$ ,
- (3)  $\mathcal{P}(<_P b_n)$  is isomorphic to  $\mathcal{L}_n^1$ , and
- (4)  $\mathcal{P}(>_P b_n)$  is isomorphic to  $(\mathcal{L}_n^0)^*$ ,

(where  $\mathcal{L}^*$  denotes  $\mathcal{L}$  under the reverse ordering). Since all  $a_n$  and  $b_n$  are in different connected components of  $\mathcal{P}$  (when viewed as a directed graph) and since for each  $n$ , at most one of  $\mathcal{L}_n^0$  and  $\mathcal{L}_n^1$  is not well-founded, we have that  $\mathcal{P}$  contains no  $\zeta$ -chain; so we can fix a linear extension  $<_L$  without a  $\zeta$ -chain. But then  $\{n \in N \mid b_n <_L a_n\}$  is in the model  $\mathcal{N}$  and separates  $S_0$  and  $S_1$  as desired.

## REFERENCES

- [Bo69] Robert Bonnet, *Stratifications et extension des genres de chaînes dénombrables*, C. R. Acad. Sci. Paris Sér. A-B **269** (1969), A880–A882.
- [BP69] Robert Bonnet and Maurice Pouzet, *Extension et stratification d’ensembles dispersés*, C. R. Acad. Sci. Paris Sér. A-B **268** (1969), A1512–A1515.
- [BP82] Robert Bonnet and Maurice Pouzet, *Linear extensions of ordered sets*, Ordered Sets (Proceedings of a NATO Advanced Study Institute held in Banff, Alta., August 28–September 12, 1981) (Ivan Rival, ed.), D. Reidel Publishing Co., Dordrecht-Boston, Mass., 1982, pp. 125–170.
- [De12] Max Dehn, *Über unendliche diskontinuierliche Gruppen*, Math. Ann. **71** (1912), 116–144.
- [Do98] Rodney G. Downey, *Computability theory and linear orderings*, Handbook of Recursive Mathematics, Vol. 2, North-Holland, Amsterdam, 1998, pp. 823–976.
- [EGNR98] Yurii L. Ershov, Sergey S. Goncharov, Anil Nerode, and Jeffrey Remmel (eds.), *Handbook of Recursive Mathematics, Vols. 1 and 2*, North-Holland, Amsterdam, 1998.
- [Fr74] Harvey Friedman, *Some systems of second order arithmetic and their uses*, Proceedings of the International Congress of Mathematicians, Vancouver, 1974, Canadian Mathematical Congress, Montreal, Que., 1975, pp. 235–242.
- [FRS87] Harvey Friedman, Neil Robertson, and Paul Seymour, *The metamathematics of the graph minor theorem*, Logic and Combinatorics (Stephen G. Simpson, ed.), American Mathematical Society, Providence, R. I., 1987, pp. 229–261.
- [FSS83] Harvey Friedman, Stephen G. Simpson, and Rick L. Smith, *Countable algebra and set existence axioms*, Ann. Pure Appl. Logic **25** (1983), 141–181.
- [FS56] Albrecht Fröhlich and John C. Shepherdson, *Effective procedures in field theory*, Trans. London Roy. Soc. Ser. A **248** (1956), 407–432.
- [Jo72] Carl G. Jockusch, Jr., *Ramsey’s theorem and recursion theory*, J. Symbolic Logic **37** (1972), 268–280.
- [JS72] Carl G. Jockusch, Jr. and Robert I. Soare,  $\Pi_1^0$  classes and degrees of theories, Trans. Amer. Math. Soc. **173** (1972), 33–56.

- [Ju69] Pierre Jullien, *Contribution à l'étude des types d'ordre dispersés*, Thèse, Marseille, 1969.
- [MN79] George Metakides and Anil Nerode, *Effective content of field theory*, Ann. Math. Logic **17** (1979), 289–320.
- [PH77] Jeff Paris and Leo Harrington, *A mathematical incompleteness in Peano arithmetic*, Handbook of Mathematical Logic (Jon Barwise, ed.), North-Holland Publishing Co., Amsterdam, 1977, pp. 1133–1142.
- [Ra60] Michael O. Rabin, *Computable algebra, general theory and theory of computable fields*, Trans. Amer. Math. Soc. **95** (1960), 341–360.
- [Ro67] Hartley G. Rogers, Jr., *Theory of Recursive Functions and Effective Computability*, McGraw-Hill, New York, 1967.
- [Ro82] Joseph G. Rosenstein, *Linear orderings*, Academic Press, New York-London, 1982.
- [Ro84] Joseph G. Rosenstein, *Recursive linear orderings*, Orders: Description and Roles (Proceedings of the fourth conference on ordered sets and their applications held in L'Arbresle, July 5–11, 1982) (Maurice Pouzet and Denis Richard, eds.), North-Holland, Amsterdam-New York, 1984, pp. 465–475.
- [Si99] Stephen G. Simpson, *Subsystems of Second Order Arithmetic*, Springer-Verlag, Berlin, 1999.
- [SW98] Theodore A. Slaman and W. Hugh Woodin, *Extending partial orders to dense linear orders*, Ann. Pure Appl. Logic **94** (1998), 253–261.
- [So87] Robert I. Soare, *Recursively Enumerable Sets and Degrees*, Springer-Verlag, Berlin-New York, 1987.
- [Sol99] Reed Solomon, *Ordered groups: a case study in reverse mathematics*, Bull. Symbolic Logic **5** (1999), 45–58.
- [Sz30] Edward Szpilrajn, *Sur l'extension de l'ordre partiel*, Fund. Math. **16** (1930), 386–389.

SCHOOL OF MATHEMATICAL AND COMPUTING SCIENCES, VICTORIA UNIVERSITY, POST OFFICE BOX 600, WELLINGTON, NEW ZEALAND

*E-mail address:* rod.downey@mcs.vuw.ac.nz

SCHOOL OF MATHEMATICAL AND COMPUTING SCIENCES, VICTORIA UNIVERSITY, POST OFFICE BOX 600, WELLINGTON, NEW ZEALAND

*Current address:* Department of Mathematics, University of Chicago, Chicago, IL 60637-1514, USA

*E-mail address:* drh@math.uchicago.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON, WI 53706-1388, USA

*E-mail address:* lempp@math.wisc.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON, WI 53706-1388, USA

*Current address:* Department of Mathematics, University of Connecticut, Storrs, CT 06269-3009, USA

*E-mail address:* david.solomon@uconn.edu