

Algebra & Number Theory

Volume 10

2016

No. 4

Local bounds for L^p norms of Maass forms in the
level aspect

Simon Marshall



Local bounds for L^p norms of Maass forms in the level aspect

Simon Marshall

We apply techniques from harmonic analysis to study the L^p norms of Maass forms of varying level on a quaternion division algebra. Our first result gives a candidate for the local bound for the sup norm in terms of the level, which is new when the level is not squarefree. The second result is a bound for L^p norms in the level aspect that is analogous to Sogge's theorem on L^p norms of Laplace eigenfunctions.

1. Introduction

Let ϕ be a cuspidal newform of level $\Gamma_0(N)$ on GL_2/\mathbb{Q} or a quaternion division algebra over \mathbb{Q} , which we shall assume is L^2 -normalised with respect to the measure that gives $\Gamma_0(N)\backslash\mathbb{H}^2$ mass 1. There has recently been interest in bounding the sup norm $\|\phi\|_\infty$ in terms of N and the infinite component of ϕ , see [Blomer and Holowinsky 2010; Harcos and Templier 2013; 2012; Saha 2015b; Templier 2010; 2014; 2015]. The “trivial” bound in the level aspect (with the infinite component remaining bounded) is generally considered to be $\|\phi\|_\infty \ll_\epsilon N^{1/2+\epsilon}$, provided N is squarefree; see [Abbes and Ullmo 1995] or any of the previously cited papers. Our first result is a candidate for the generalisation of this to arbitrary N .

Theorem 1. *Let D/\mathbb{Q} be a quaternion division algebra that is split at infinity. Let ϕ be an L^2 -normalised newform of level $K_0(N)$ on $\mathrm{PGL}_1(D)$, where N is odd and coprime to the primes that ramify in D . Assume that ϕ is spherical at infinity with spectral parameter t , which is related to the Laplace eigenvalue by the equation $(\Delta + 1/4 + t^2)\phi = 0$. Let $N_0 \geq 1$ be the smallest number with $N|N_0^2$. We have*

$$\|\phi\|_\infty \ll (1 + |t|)^{1/2} N_0^{1/2} \prod_{p|N} (1 + 1/p)^{1/2}. \quad (1)$$

Supported by NSF grants DMS-1509331 and DMS-1501230.

MSC2010: primary 35P20; secondary 11F41, 11F25.

Keywords: L^p norms, sup norms, Maass forms, automorphic forms.

Notation is standard, and specified below. When t is bounded, [Theorem 1](#) gives a bound of $N^{1/2+\epsilon}$ for N squarefree, but roughly $N^{1/4+\epsilon}$ for powerful N . While the theorem is restricted to compact quotients, [Section 3.1](#) gives a weaker result in the case of $\mathrm{PGL}_2/\mathbb{Q}$. This has been strengthened by Saha [\[2015a\]](#), who proves a bound on $\mathrm{PGL}_2/\mathbb{Q}$ that combines [\(1\)](#) in the level aspect with the $t^{5/12+\epsilon}$ bound of Iwaniec and Sarnak [\[1995\]](#) in the eigenvalue aspect. He does this by combining the methods of Iwaniec and Sarnak with bounds for Whittaker functions and an amplification argument.

Our second result is the analogue in the level aspect of a classical theorem of Sogge [\[1988\]](#), which we now recall. Let M be a compact Riemannian surface with Laplacian Δ , and let ψ be a function on M satisfying $(\Delta + \lambda^2)\psi = 0$ and $\|\psi\|_2 = 1$. Define $\delta : [2, \infty) \rightarrow \mathbb{R}$ by

$$\delta(p) = \begin{cases} \frac{1}{2} - \frac{2}{p}, & 0 \leq \frac{1}{p} \leq \frac{1}{6}, \\ \frac{1}{4} - \frac{1}{2p}, & \frac{1}{6} \leq \frac{1}{p} \leq \frac{1}{2}. \end{cases} \tag{2}$$

Sogge’s theorem states that

$$\|\psi\|_p \ll \lambda^{\delta(p)} \quad \text{for } 2 \leq p \leq \infty. \tag{3}$$

In particular, this is stronger than the bound obtained by interpolating between bounds for the L^2 and L^∞ norms. Our next theorem demonstrates that something similar is possible in the level aspect.

Theorem 2. *Let D/\mathbb{Q} be a quaternion division algebra that is split at infinity. Let ϕ be an L^2 -normalised newform of level $K_0(q^2)$ on $\mathrm{PGL}_1(D)$, where q is an odd prime that does not ramify in D . Assume that ϕ is principal series at q , that ϕ is spherical at infinity with spectral parameter t , and that $|t| \leq A$ for some $A > 0$. We have*

$$\|\phi\|_p \ll_A q^{\delta(p)}.$$

It should be possible to give some extension of [Theorem 2](#) to general ϕ , although in some cases the method may not give any improvement over the bound given by interpolating between L^2 and L^∞ norms. In particular, this seems to occur when ϕ is special at q . We have chosen to work in the simplest case where the method gives a nontrivial result.

[Theorems 1](#) and [2](#) are an attempt to prove the correct local bounds for L^p norms of eigenfunctions in the level aspect, in the same way that [\(3\)](#) is the local bound in the eigenvalue aspect. The term “local bound” means the best bound that may be proved by only considering the behaviour of ϕ in one small open set at a time, without taking the global structure of the space into account.

The bound [\(3\)](#) is sharp on the round sphere. In the same way, one may obtain limited evidence that [Theorem 1](#), and [Theorem 2](#) for $p \geq 6$, are the sharp local

bounds by comparing them with what may be proved on the “compact form” of the arithmetic quotient being considered. In the case of [Theorem 2](#), this means taking an L^2 -normalised function ψ on $\mathrm{PGL}(2, \mathbb{Z}_q)$ of the same type as ϕ' defined below — in other words, invariant under the group $K(q, q)$ defined in [Section 2.1](#), and generating an irreducible representation of the same type as ϕ' under right translation — and proving bounds for $\|\psi\|_p$. We may prove that $\|\psi\|_p \ll q^{\delta(p)}$ in the same way as [Theorem 2](#), after which [Equation \(4\)](#) and [Lemma 11](#) imply that this is sharp for $p \geq 6$. An analogous statement may be proved for [Theorem 1](#) when N is a growing power of a fixed prime. However, we do not yet know if [Theorem 2](#) is sharp in this sense for $2 \leq p \leq 6$. We expect the bound of [Theorem 1](#) to have a natural expression as the square root of the Plancherel density around the representation of ϕ .

Because the proofs do not make use of the global structure of the arithmetic quotient, it should be possible to improve the exponents by using arithmetic amplification.

2. Notation

2.1. Adelic groups. Let \mathbb{A} and \mathbb{A}_f be the adèles and finite adèles of \mathbb{Q} . Let D/\mathbb{Q} be a quaternion division algebra that is split at infinity. Let S be the set containing 2 and all primes that ramify in D , and let $S_\infty = S \cup \{\infty\}$. Let $G = \mathrm{PGL}_1(D)$. If v is a place of \mathbb{Q} , let $G_v = G(\mathbb{Q}_v)$. Let $X = G(\mathbb{Q}) \backslash G(\mathbb{A})$. Let $\mathcal{O} \subset D$ be a maximal order. Let $K = \bigotimes_p K_p \subset G(\mathbb{A}_f)$ be a compact subgroup with the properties that K_p is open in G_p for $p \in S$, and K_p is isomorphic to the image of \mathcal{O}_p^\times in G_p when $p \notin S$. This allows us to choose isomorphisms $K_p \simeq \mathrm{PGL}(2, \mathbb{Z}_p)$ when $p \notin S$. When $M, N \geq 1$ are prime to S , we shall use these isomorphisms to define the upper triangular congruence subgroup $K_0(N)$, principal congruence subgroup $K(N)$, and

$$K(M, N) = \left\{ k \in K : k \equiv \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} (M), k \equiv \begin{pmatrix} * & 0 \\ * & * \end{pmatrix} (N) \right\}$$

in the natural way. We choose a maximal compact subgroup $K_\infty \subset G_\infty$.

We fix a Haar measure on $G(\mathbb{A})$ by taking the product of the measures on G_p assigning mass 1 to K_p , and any Haar measure on G_∞ . We use this measure to define convolution of functions on $G(\mathbb{A})$, which we denote by $*$, and if $f \in C_0^\infty(G(\mathbb{A}))$ we use it to define the operator $R(f)$ by which f acts on $L^2(X)$. If H is a group and f is a function on H , we define the function f^\vee by $f^\vee(h) = \bar{f}(h^{-1})$. If $f \in C_0^\infty(G(\mathbb{A}))$, the operators $R(f)$ and $R(f^\vee)$ are adjoints.

2.2. Newforms. Let $N \geq 1$ be prime to S . We shall say that $\phi \in L^2(X)$ is a newform of level $K_0(N)$ if ϕ lies in an automorphic representation $\pi = \bigotimes_v \pi_v$ of G , ϕ is invariant under $K_0(N)$, and we have a factorisation $\phi = \bigotimes_v \phi_v$ where ϕ_v is a local

newvector of level N for $v \notin S_\infty$. We shall say that ϕ is spherical with spectral parameter $t \in \mathbb{C}$ if π_∞ satisfies these conditions, and ϕ is invariant under K_∞ . Note that our normalisation of t is such that the tempered principal series corresponds to $t \in \mathbb{R}$.

2.3. The Harish-Chandra transform. Given $k \in C_0^\infty(G_\infty)$, we define its Harish-Chandra transform by

$$\hat{k}(t) = \int_{G_\infty} k(g)\varphi_t(g) dg$$

for $t \in \mathbb{C}$, where φ_t is the standard spherical function with spectral parameter t . We will use the following standard result on the existence of a K_∞ -biinvariant function with concentrated spectral support.

Lemma 3. *There is a compact set $B \subset G_\infty$ such that for any $t \in \mathbb{R} \cup [0, i/2]$, there is a K_∞ -biinvariant function $k \in C_0^\infty(G_\infty)$ with the following properties:*

- (a) *The function k is supported in B , and $\|k\|_\infty \ll 1 + |t|$.*
- (b) *The Harish-Chandra transform \hat{k} is nonnegative on $\mathbb{R} \cup [0, i/2]$, and satisfies $\hat{k}(t) \geq 1$.*

Proof. When $t \in \mathbb{R}$ and $|t| \geq 1$, this is, e.g., Lemma 2.1 of [Templier 2015]. When $|t| \leq 1$, one may fix a K_∞ -biinvariant real bump function k_0 supported near the identity and define $k = k_0 * k_0$. □

Note that condition (b) implies that $k = k^\vee$.

2.4. Inner products of matrix coefficients. The following lemma is known as Schur orthogonality, see Theorem 2.4 and Proposition 2.11 of [Bump 2013] for the proof.

Lemma 4. *Let H be a finite group, and let dh be the Haar measure of mass 1 on H . Let (ρ, V) be an irreducible representation of H , and let $\langle \cdot, \cdot \rangle$ be a positive definite H -invariant Hermitian form on V . If $v_i \in V$ for $1 \leq i \leq 4$, we have*

$$\int_H \langle \rho(h)v_1, v_2 \rangle \overline{\langle \rho(h)v_3, v_4 \rangle} dh = \frac{\langle v_1, v_3 \rangle \overline{\langle v_2, v_4 \rangle}}{\dim V}. \tag{4}$$

3. Proof of Theorem 1

Choose $h \in G(\mathbb{A})$ by setting $h_v = 1$ for $v \in S_\infty$, and

$$h_v = \begin{pmatrix} N & \\ & N_0 \end{pmatrix}$$

when $v \notin S_\infty$. We define $\phi' = R(h)\phi$, so that ϕ' is invariant under $K(N_0, N/N_0)$. Let $V = \otimes V_v \subset \pi$ be the space generated by ϕ' under the action of K .

Lemma 5. V is an irreducible representation of K , and

$$\dim V \leq N_0 \prod_{p|N} (1 + 1/p).$$

Proof. It suffices to prove the analogous statement for the tensor factors V_p , and we may assume that $p \notin S$. If we could write $V_p = V^1 + V^2$, where V^i are nontrivial K_p -invariant subspaces, then the projection of ϕ'_p to each subspace would be invariant under $K_p(N_0, N/N_0)$. However, this contradicts the uniqueness of the newvector.

As V_p is irreducible and factors through $K_p/K_p(N_0)$, the lemma now follows from the results of Silberger [1970, §3.4], in particular the remarks on pages 96–97. Note that we use our assumption that $2 \in S$ at this point. \square

We define $k_f \in C_0^\infty(G(\mathbb{A}_f))$ to be $\overline{\langle R(g)\phi', \phi' \rangle}$ for $g \in K$ and 0 otherwise. Choose a function $k_\infty \in C_0^\infty(G_\infty)$ as in Lemma 3, and define $k = k_\infty k_f$. It may be seen that $k = k^\vee$, which implies that $R(k)$ is self-adjoint. Lemma 5 and Equation (4) imply that $k_f = \dim V k_f * k_f$, and combining this with Lemma 3(b) gives that $R(k)$ is nonnegative. Lemmas 3 and 5 and Equation (4) imply that $R(k)\phi' = \lambda\phi'$, where $\lambda > 0$ and

$$\lambda^{-1} \leq \dim V \leq N_0 \prod_{p|N} (1 + 1/p). \quad (5)$$

Extend ϕ' to an orthonormal basis $\{\phi_i\}$ of eigenfunctions for $R(k)$ with eigenvalues $\lambda_i \geq 0$. The pretrace formula associated to k is

$$\sum_i \lambda_i |\phi_i(x)|^2 = \sum_{\gamma \in G(\mathbb{Q})} k(x^{-1}\gamma x)$$

and dropping all terms from the left hand side but ϕ' gives

$$\lambda |\phi'(x)|^2 \leq \sum_{\gamma \in G(\mathbb{Q})} k(x^{-1}\gamma x). \quad (6)$$

The compactness of X and uniformly bounded support of k implies that the number of nonzero terms on the right hand side is bounded independently of x , and combining (5) and Lemma 3(a) completes the proof.

3.1. A result in the noncompact case. If we set $G = \mathrm{PGL}_2/\mathbb{Q}$, it may be seen that we have the following analogue of Theorem 1.

Proposition 6. Let $\Omega \subset G(\mathbb{Q}) \backslash G(\mathbb{A})$ be compact. Let ϕ be an L^2 -normalised newform of level $K_0(N)$ on G , where N is odd. Assume that ϕ is spherical at infinity with spectral parameter t . Let $N_0 \geq 1$ be the smallest number with $N|N_0^2$. If ϕ' is related to ϕ as above, we have $\|\phi'\|_\Omega \ll (1 + |t|)^{1/2} N_0^{1/2} \prod_{p|N} (1 + 1/p)^{1/2}$.

See [Saha 2015a] for a strengthening of this result.

4. Proof of Theorem 2

We maintain the notation ϕ' , V , and k_f from Section 3. Our assumption $2 \in S$ implies that $q \geq 3$.

Lemma 7. *We have $\dim V = q$ or $q + 1$.*

Proof. We have assumed that π_q is isomorphic to an irreducible principal series representation $\mathcal{I}(\chi, \chi^{-1})$, for some character χ of \mathbb{Q}_q^\times with conductor q . By considering the compact model of $\mathcal{I}(\chi, \chi^{-1})$, and applying the fact that V factors through $K/K(q) \simeq \text{PGL}(2, \mathbb{F}_q)$, we see that V must be a subrepresentation of a principal series representation of $\text{PGL}(2, \mathbb{F}_q)$. (Here \mathbb{F}_q denotes the field with q elements.) It follows that $\dim V$ must be either 1, q , or $q + 1$. The possibility $\dim V = 1$ is ruled out because any one-dimensional representation of K that is trivial on $K(q, q)$ must be trivial, and this contradicts our assumption that ϕ is new at $K_0(q^2)$. □

Let $k_\infty^0 \in C_0^\infty(G_\infty)$ be a real-valued K_∞ -biinvariant function, so that $k_\infty^0 = (k_\infty^0)^\vee$. If we choose k_∞^0 to be a nonnegative bump function with sufficiently small support, we may assume that its Harish-Chandra transform satisfies $\hat{k}_\infty^0(t) \geq 1$ for $t \in [0, A] \cup [0, i/2]$. We define $k_\infty = k_\infty^0 * k_\infty^0$. Let $k_0 = k_\infty^0 k_f$, and $k = k_\infty k_f$. Let $T_0 = R(k_0)$ and $T = R(k)$. We see that T_0 is self-adjoint, and Equation (4) implies that $T = \dim V T_0^2$. Let $W \subset K_q$ be the subgroup $\{1, w\}$, where

$$w = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

Let $k_{1,f} \in C_0^\infty(G(\mathbb{A}_f))$ be k_f times the characteristic function of $WK(q, q)$, and let $k_{2,f} = k_f - k_{1,f}$. Let $k_i = k_\infty k_{i,f}$, and $T_i = R(k_i)$. The proof of Theorem 2 works by combining the decomposition $T = T_1 + T_2$ with interpolation between the following bounds.

Lemma 8. *We have*

$$\|T_1 f\|_\infty \ll \|f\|_1, \quad \|T_2 f\|_\infty \ll q^{-1/2} \|f\|_1,$$

for any $f \in C^\infty(X)$.

Proof. The integral kernels of T_i are given by

$$\sum_{\gamma \in G(\mathbb{Q})} k_i(x^{-1}\gamma y).$$

The result now follows from the compactness of $G(\mathbb{Q}) \backslash G(\mathbb{A})$, the bound $\|k_1\|_\infty \ll 1$, and the bound $\|k_2\|_\infty \ll q^{-1/2}$ which follows from Lemma 11 below. □

Lemma 9. *We have*

$$\|T_1 f\|_2 \ll q^{-2} \|f\|_2, \quad \|T_2 f\|_2 \ll q^{-1} \|f\|_2,$$

for any $f \in C^\infty(X)$.

Proof. The choice of k_∞ and the identity $k_f = \dim V k_f * k_f$ imply that the $L^2 \rightarrow L^2$ norm of T is $\ll (\dim V)^{-1} \leq q^{-1}$. [Lemma 11](#) implies that $k_{1,f} = k_{1,f}^\vee$ and

$$k_{1,f} = [K : WK(q, q)] k_{1,f} * k_{1,f} = (q(q+1)/2) k_{1,f} * k_{1,f},$$

and this implies that the $L^2 \rightarrow L^2$ norm of T_1 is $\ll q^{-2}$. The bound for T_2 follows from the triangle inequality. \square

Interpolating between these bounds gives the following.

Lemma 10. *We have $\|Tf\|_p \ll q^{2\delta(p)-1} \|f\|_{p'}$ for any $2 \leq p \leq \infty$, where p' is the dual exponent to p .*

Proof. Applying the Riesz–Thorin interpolation theorem [[Folland 1999](#), Theorem 6.27] to the bounds

$$\|T_1 f\|_\infty \ll \|f\|_1, \quad \|T_1 f\|_2 \ll q^{-2} \|f\|_2$$

gives $\|T_1 f\|_p \ll q^{-4/p} \|f\|_{p'}$ for $2 \leq p \leq \infty$, and applying it to

$$\|T_2 f\|_\infty \ll q^{-1/2} \|f\|_1, \quad \|T_2 f\|_2 \ll q^{-1} \|f\|_2$$

gives $\|T_2 f\|_p \ll q^{-1/2-1/p} \|f\|_{p'}$ for $2 \leq p \leq \infty$. Applying Minkowski's inequality then gives

$$\|Tf\|_p \leq \|T_1 f\|_p + \|T_2 f\|_p \ll (q^{-4/p} + q^{-1/2-1/p}) \|f\|_{p'},$$

and the observation $2\delta(p) - 1 = \max(-4/p, -1/2 - 1/p)$ completes the proof. \square

We now combine [Lemma 10](#) with the usual adjoint-square argument: we have

$$\begin{aligned} \langle \dim V T_0^2 f, f \rangle &= \langle Tf, f \rangle \\ &\ll q^{2\delta(p)-1} \|f\|_{p'}^2, \\ \langle T_0 f, T_0 f \rangle &\ll q^{2\delta(p)-2} \|f\|_{p'}^2, \\ \|T_0 f\|_2 &\ll q^{\delta(p)-1} \|f\|_{p'}. \end{aligned}$$

Taking adjoints gives $\|T_0 f\|_p \ll q^{\delta(p)-1} \|f\|_2$. Applying this with $f = \phi'$ and estimating the eigenvalue of T_0 on ϕ' as in [Section 3](#) completes the proof.

Lemma 11. *Let π_q be isomorphic to an irreducible principal series representation $\mathcal{I}(\chi, \chi^{-1})$, for some character χ of \mathbb{Q}_q^\times with conductor q . When $g \in K_q$, the matrix coefficient $\langle \pi_q(g)\phi'_q, \phi'_q \rangle$ satisfies*

$$\langle \pi_q(g)\phi'_q, \phi'_q \rangle = 1, \quad g \in K_q(q, q), \quad (7)$$

$$\langle \pi_q(g)\phi'_q, \phi'_q \rangle = \chi(-1), \quad g \in wK_q(q, q), \quad (8)$$

$$\langle \pi_q(g)\phi'_q, \phi'_q \rangle \ll q^{-1/2}, \quad g \notin WK_q(q, q), \quad (9)$$

where the implied constant is absolute.

Proof. We may reduce the problem to one for the group $\mathrm{PGL}(2, \mathbb{F}_q)$ as in Lemma 7. We let T and B be the usual diagonal and upper triangular subgroups of $\mathrm{PGL}(2, \mathbb{F}_q)$. We now think of χ as a nontrivial character of \mathbb{F}_q^\times , and let (ρ, H) denote the corresponding induced representation of $\mathrm{PGL}(2, \mathbb{F}_q)$. We realise H as the space of functions $f : \mathrm{PGL}(2, \mathbb{F}_q) \rightarrow \mathbb{C}$ satisfying

$$f\left(\begin{pmatrix} a & b \\ 0 & d \end{pmatrix}g\right) = \chi(a/d)f(g)$$

with the invariant Hermitian form

$$\langle f_1, f_2 \rangle = \sum_{g \in B \backslash \mathrm{PGL}(2, \mathbb{F}_q)} f_1(g)\overline{f_2(g)}. \quad (10)$$

It may be seen that there is a unique function $f_0 \in H$ that is invariant under T , up to scaling, and we may choose it to be

$$f_0 : \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto \begin{cases} \chi(\det/cd)/\sqrt{q-1}, & cd \neq 0, \\ 0, & cd = 0. \end{cases} \quad (11)$$

It follows that $\|f_0\| = 1$, and so $\langle \pi_q(g)\phi'_q, \phi'_q \rangle = \langle \rho(g)f_0, f_0 \rangle$. Equation (7) is immediate, and (8) follows from $\rho(w)f_0 = \chi(-1)f_0$. To prove (9), we assume that

$$g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \notin WT.$$

We may write $\langle \rho(g)f_0, f_0 \rangle$ as a sum over \mathbb{F}_q as follows.

Lemma 12. *We have*

$$\langle \rho(g)f_0, f_0 \rangle = (q-1)^{-1}\chi(\det(g))\sum_n \chi^{-1}((c+an)(d+bn))\chi(n). \quad (12)$$

Proof. We choose a set of coset representatives for $B \backslash \mathrm{PGL}(2, \mathbb{F}_q)$ consisting of

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 1 & 0 \\ n & 1 \end{pmatrix}, \quad n \in \mathbb{F}_q.$$

Applying (11) for these representatives gives

$$f_0\left(\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}\right) = 0 \quad \text{and} \quad f_0\left(\begin{pmatrix} 1 & 0 \\ n & 1 \end{pmatrix}\right) = \chi^{-1}(n)/\sqrt{q-1}.$$

The first coset representative therefore makes no contribution to $\langle \rho(g)f_0, f_0 \rangle$. For the others, we calculate

$$\begin{aligned} [\rho(g)f_0]\left(\begin{pmatrix} 1 & 0 \\ n & 1 \end{pmatrix}\right) &= f_0\left(\begin{pmatrix} 1 & 0 \\ n & 1 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix}\right) \\ &= f_0\left(\begin{pmatrix} a & b \\ c+an & d+bn \end{pmatrix}\right) \\ &= \chi(\det(g))\chi^{-1}((c+an)(d+bn))/\sqrt{q-1}. \end{aligned}$$

Substituting these into (10) completes the proof. \square

We bound the sum (12) by rewriting it as

$$\sum_n \chi^{-1}((c+an)(d+bn)n^{q-2})$$

and applying [Schmidt 2004, Chapter 2, Theorem 2.4] (see also [Iwaniec and Kowalski 2004, Theorem 11.23]). We must first check that $(c+an)(d+bn)n^{q-2}$ is not a proper power. The assumption $g \notin WT$ implies that one or both of $a+cn$ and $b+dn$ have a root distinct from 0. If they both have the same root distinct from 0, this contradicts the invertibility of g . Therefore $(c+an)(d+bn)n^{q-2}$ must have at least one root of multiplicity 1, so it cannot be a power. As $(c+an)(d+bn)n^{q-2}$ has at most 3 distinct roots, [Schmidt 2004] or [Iwaniec and Kowalski 2004] therefore give

$$\left| \sum_n \chi((c+an)(d+bn)n^{q-2}) \right| \leq 2\sqrt{q},$$

which completes the proof of (9). \square

Acknowledgements

We would like to thank Valentin Blomer, Abhishek Saha, and Nicolas Templier for helpful discussions, and the referee for a careful reading of the paper.

References

- [Abbes and Ullmo 1995] A. Abbes and E. Ullmo, “Comparaison des métriques d’Arakelov et de Poincaré sur $X_0(N)$ ”, *Duke Math. J.* **80**:2 (1995), 295–307. MR 1369394 Zbl 0895.14007
- [Blomer and Holowinsky 2010] V. Blomer and R. Holowinsky, “Bounding sup-norms of cusp forms of large level”, *Invent. Math.* **179**:3 (2010), 645–681. MR 2587342 Zbl 1243.11059

- [Bump 2013] D. Bump, *Lie groups*, 2nd ed., Graduate Texts in Mathematics **225**, Springer, New York, 2013. [MR 3136522](#) [Zbl 1279.22001](#)
- [Folland 1999] G. B. Folland, *Real analysis*, 2nd ed., Wiley, New York, 1999. [MR 1681462](#) [Zbl 0924.28001](#)
- [Harcos and Templier 2012] G. Harcos and N. Templier, “On the sup-norm of Maass cusp forms of large level: II”, *Int. Math. Res. Not.* **2012**:20 (2012), 4764–4774. [MR 2989618](#) [Zbl 1332.11048](#)
- [Harcos and Templier 2013] G. Harcos and N. Templier, “On the sup-norm of Maass cusp forms of large level, III”, *Math. Ann.* **356**:1 (2013), 209–216. [MR 3038127](#) [Zbl 1332.11049](#)
- [Iwaniec and Kowalski 2004] H. Iwaniec and E. Kowalski, *Analytic number theory*, American Mathematical Society Colloquium Publications **53**, American Mathematical Society, Providence, RI, 2004. [MR 2061214](#) [Zbl 1059.11001](#)
- [Iwaniec and Sarnak 1995] H. Iwaniec and P. Sarnak, “ L^∞ norms of eigenfunctions of arithmetic surfaces”, *Ann. of Math. (2)* **141**:2 (1995), 301–320. [MR 1324136](#) [Zbl 0833.11019](#)
- [Saha 2015a] A. Saha, “Hybrid sup-norm bounds for Maass newforms of powerful level”, preprint, 2015. [arXiv 1509.07489](#)
- [Saha 2015b] A. Saha, “On sup-norms of cusp forms of powerful level”, preprint, 2015. [arXiv 1404.3179](#)
- [Schmidt 2004] W. Schmidt, *Equations over finite fields: an elementary approach*, 2nd ed., Kendrick Press, Heber City, UT, 2004. [MR 2121285](#) [Zbl 1128.11032](#)
- [Silberger 1970] A. J. Silberger, PGL_2 over the p -adics: its representations, spherical functions, and Fourier analysis, Lecture Notes in Mathematics **166**, Springer, Berlin, 1970. [MR 0285673](#) [Zbl 0204.44102](#)
- [Sogge 1988] C. D. Sogge, “Concerning the L^p norm of spectral clusters for second-order elliptic operators on compact manifolds”, *J. Funct. Anal.* **77**:1 (1988), 123–138. [MR 930395](#) [Zbl 0641.46011](#)
- [Templier 2010] N. Templier, “On the sup-norm of Maass cusp forms of large level”, *Selecta Math. (N.S.)* **16**:3 (2010), 501–531. [MR 2734340](#) [Zbl 1260.11032](#)
- [Templier 2014] N. Templier, “Large values of modular forms”, *Camb. J. Math.* **2**:1 (2014), 91–116. [MR 3272013](#) [Zbl 1307.11062](#)
- [Templier 2015] N. Templier, “Hybrid sup-norm bounds for Hecke–Maass cusp forms”, *J. Eur. Math. Soc. (JEMS)* **17**:8 (2015), 2069–2082. [MR 3372076](#) [Zbl 06481085](#)

Communicated by Philippe Michel

Received 2015-03-09

Revised 2015-12-31

Accepted 2016-04-23

marshall@math.wisc.edu

Department of Mathematics, University of Wisconsin–Madison,
480 Lincoln Drive, Madison, WI 53706, United States

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Bjorn Poonen
Massachusetts Institute of Technology
Cambridge, USA

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Georgia Benkart	University of Wisconsin, Madison, USA	Susan Montgomery	University of Southern California, USA
Dave Benson	University of Aberdeen, Scotland	Shigefumi Mori	RIMS, Kyoto University, Japan
Richard E. Borcherds	University of California, Berkeley, USA	Raman Parimala	Emory University, USA
John H. Coates	University of Cambridge, UK	Jonathan Pila	University of Oxford, UK
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Anand Pillay	University of Notre Dame, USA
Brian D. Conrad	Stanford University, USA	Victor Reiner	University of Minnesota, USA
Hélène Esnault	Freie Universität Berlin, Germany	Peter Sarnak	Princeton University, USA
Hubert Flenner	Ruhr-Universität, Germany	Joseph H. Silverman	Brown University, USA
Sergey Fomin	University of Michigan, USA	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Andrew Granville	Université de Montréal, Canada	J. Toby Stafford	University of Michigan, USA
Joseph Gubeladze	San Francisco State University, USA	Ravi Vakil	Stanford University, USA
Roger Heath-Brown	Oxford University, UK	Michel van den Bergh	Hasselt University, Belgium
Craig Huneke	University of Virginia, USA	Marie-France Vignéras	Université Paris VII, France
Kiran S. Kedlaya	Univ. of California, San Diego, USA	Kei-Ichi Watanabe	Nihon University, Japan
János Kollár	Princeton University, USA	Efim Zelmanov	University of California, San Diego, USA
Yuri Manin	Northwestern University, USA	Shou-Wu Zhang	Princeton University, USA
Philippe Michel	École Polytechnique Fédérale de Lausanne		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor


See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2016 is US \$290/year for the electronic version, and \$485/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

ANT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 10 No. 4 2016

Moduli of morphisms of logarithmic schemes	695
JONATHAN WISE	
Residual intersections and the annihilator of Koszul homologies	737
SEYED HAMID HASSANZADEH and JOSE NAÉLITON	
The Prym map of degree-7 cyclic coverings	771
HERBERT LANGE and ANGELA ORTEGA	
Local bounds for L^p norms of Maass forms in the level aspect	803
SIMON MARSHALL	
Hasse principle for Kummer varieties	813
YONATAN HARPAZ and ALEXEI N. SKOROBOGATOV	
Analytic continuation on Shimura varieties with μ-ordinary locus	843
STÉPHANE BIJAKOWSKI	
A note on secondary K-theory	887
GONÇALO TABUADA	
Nef cones of Hilbert schemes of points on surfaces	907
BARBARA BOLOGNESE, JACK HUIZENGA, YINBANG LIN, ERIC RIEDL, BENJAMIN SCHMIDT, MATTHEW WOOLF and XIAOLEI ZHAO	
Interpolation for restricted tangent bundles of general curves	931
ERIC LARSON	