

The Limitations of Cupping in the Local Structure of the Enumeration Degrees

Mariya I. Soskova

Department of Pure Mathematics
University of Leeds
mariya@maths.leeds.ac.uk

17.06.08

The cupping property

Let $\langle \mathcal{A}, \leq, \vee, \mathbf{1} \rangle$ be an upper semi-lattice with least and greatest element.

Definition

An element $\mathbf{a} \in \mathcal{A}$ is *cuppable* if there exists an element $\mathbf{b} \in \mathcal{A}$, $\mathbf{b} \neq \mathbf{1}$ such that $\mathbf{a} \vee \mathbf{b} = \mathbf{1}$.

The element \mathbf{b} is called a *cupping partner* for \mathbf{a} .

Results in the Turing degrees

- ▶ Posner, Robinson: Every nonzero degree in $\mathcal{D}_T(\leq 0')$ is cuppable.
- ▶ Cooper, Yates: There exists a nonzero c.e. Turing degree which cannot be cupped by any incomplete c.e. Turing degree.

Enumeration degrees

Definition


A set A is *enumeration reducible* (\leq_e) to a set B if there is a c.e. set Φ such that:

$$n \in A \Leftrightarrow \exists u (\langle n, u \rangle \in \Phi \wedge D_u \subseteq B),$$


where D_u denotes the finite set with code u under the standard coding of finite sets.

- ▶ $A \equiv_e B \Leftrightarrow A \leq_e B \wedge B \leq_e A$
- ▶ $\langle \mathcal{D}_e, 0_e, \leq, \cup, ' \rangle$.
- ▶ $\iota : \mathcal{D}_T \rightarrow \mathcal{D}_e$.

The local structure of the enumeration degrees $\mathcal{D}_e(\leq 0'_e)$



Transferring results from the Turing degrees



Cupping results in the enumeration degrees

- ▶ Negative Results:

(Cooper, Sorbi, Yi): There exists a nonzero Σ_2^0 enumeration degree that is not cuppable.

- ▶ Positive Results:

(Cooper, Sorbi and Yi): Every nonzero Δ_2^0 e-degree is cuppable by a total incomplete Δ_2^0 e-degree.


(S, Wu): Every nonzero Δ_2^0 e-degree is cuppable by a partial and low Δ_2^0 e-degree.

Cupping partners

Question

How much further can we limit the search for cupping partners.

Δ_2^0 e-degrees



Reaching the first limit

Theorem

Let $\{\mathbf{a}_i\}_{i < \omega}$ be a Δ_2^0 -computably enumerable sequence of enumeration degrees. There exists a nonzero Δ_2^0 enumeration degree \mathbf{b} such that for every $i < \omega$ if \mathbf{a}_i is incomplete then $\mathbf{a}_i \vee \mathbf{b} \neq 0'_e$.


Here a class $\{\mathbf{a}_i\}_{i < \omega}$ of Δ_2^0 enumeration degrees is Δ_2^0 -computably enumerable if there is a computable sequence of Δ_2^0 approximations $\{A_i[s]\}_{i, s < \omega}$ to representatives A_i of every degree \mathbf{a}_i in the class.

The Difference Hierarchy

Definition (Ershov)

1. A set A is n -c.e. if there is a computable function f such that for each x , $f(x, 0) = 0$,
 $|\{s + 1 \mid f(x, s) \neq f(x, s + 1)\}| \leq n$ and $A(x) = \lim_s f(x, s)$.
2. A is ω -c.e. if there are two computable functions $f(x, s), g(x)$ such that for all x , $f(x, 0) = 0$,
 $|\{s + 1 \mid f(x, s) \neq f(x, s + 1)\}| \leq g(x)$ and $\lim_s f(x, s) = A(x)$.
3. A degree \mathbf{a} is n -c.e. (ω -c.e.) if it contains a n -c.e. (ω -c.e.) set.

A finer partition of the Δ_2^0 enumeration degrees




Consequences

Corollary

There exists a nonzero Δ_2^0 enumeration degree that cannot be cupped by any incomplete ω -c.e. degree.

Wu, S: For every nonzero ω -c.e. enumeration degree **a** there exists an incomplete 3-c.e. enumeration degree **b** that cups **a**.

Cupping classes of enumeration degrees



(Cooper, Seetapun and Li): There exists a single incomplete Δ_2^0 Turing degree that cups every nonzero c.e. Turing degree.

The second limitation

For any larger subclass, which contains the nonzero 3-c.e enumeration degrees this cannot be done as:

Theorem

Let \mathbf{a} be an incomplete Σ_2^0 enumeration degree. There exists a nonzero 3-c.e. enumeration degree \mathbf{b} such that $\mathbf{a} \vee \mathbf{b} \neq \mathbf{0}'_e$.

Proof(ideas)

Let A be a representative of the given Σ_2^0 e-degree. We construct two 3-c.e. sets X and Y so that:

- ▶ For every natural number e :


$$\mathcal{N}_e : W_e \neq X \wedge W_e \neq Y.$$

- ▶ If for some i the requirement \mathcal{P}_i^0 is not satisfied then for every i the requirement \mathcal{P}_i^1 is satisfied, where:




$$\mathcal{P}_i^0 : \Theta_i^{A,X} \neq \bar{K};$$

$$\mathcal{P}_i^1 : \Psi_i^{A,Y} \neq \bar{K}.$$





The tree



Bibliography

-  S.B. Cooper: On a theorem of C.E.M. Yates. Handwritten notes (1973)
-  A. E. M. Lewis: Finite cupping sets. Arch. Math. Logic 43, 845–858 (2004)
-  D. Posner, R. Robinson: Degrees joining to $0'$. J. Symbolic Logic 46, 714–722 (1981)

Bibliography

-  S. B. Cooper: Enumeration reducibility, nondeterministic computations and relative computability of partial functions. In: K. Ambos-Spies, G. H. Muller, G. E. Sacks (eds.) Recursion Theory Week, Proceedings Oberwolfach 1989, pp 57–110. LNM 1432 (1990)
-  S. B. Cooper, A. Sorbi, X. Yi: Cupping and noncupping in the enumeration degrees of Σ_2^0 sets. Ann. Pure Appl. Logic 82, 317–342 (1996)
-  A. H. Lachlan, R. A. Shore: The n -re enumeration degrees are dense. Arch. Math. Logic 31, 277–285 (1992)
-  M. I. Soskova, G. Wu: Cupping Δ_2^0 enumeration degrees to $0'$. In: S. Cooper, B. Löwe, A. Sorbi (eds.) Computation and Logic in the Real World, pp 727–738. LNCS 4497 (2007)