

EXERCISES 2.3

2.3.1. For the following partial differential equations, what ordinary differential equations are implied by the method of separation of variables?

$$* (a) \quad \frac{\partial u}{\partial t} = \frac{k}{r} \frac{\partial}{\partial r} \left(r \frac{\partial u}{\partial r} \right)$$

$$(b) \quad \frac{\partial u}{\partial t} = k \frac{\partial^2 u}{\partial x^2} - v_0 \frac{\partial u}{\partial x}$$

$$* (c) \quad \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$$

$$(d) \quad \frac{\partial u}{\partial t} = \frac{k}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial u}{\partial r} \right)$$

$$* (e) \quad \frac{\partial u}{\partial t} = k \frac{\partial^4 u}{\partial x^4}$$

$$* (f) \quad \frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2}$$

2.3.2. Consider the differential equation

$$\frac{d^2 \phi}{dx^2} + \lambda \phi = 0.$$

Determine the eigenvalues λ (and corresponding eigenfunctions) if ϕ satisfies the following boundary conditions. Analyze three cases ($\lambda > 0$, $\lambda = 0$, $\lambda < 0$). You may assume that the eigenvalues are real.

$$(a) \quad \phi(0) = 0 \text{ and } \phi(\pi) = 0$$

$$* (b) \quad \phi(0) = 0 \text{ and } \phi(1) = 0$$

$$(c) \quad \frac{d\phi}{dx}(0) = 0 \text{ and } \frac{d\phi}{dx}(L) = 0 \text{ (If necessary, see Sec. 2.4.1.)}$$

$$* (d) \quad \phi(0) = 0 \text{ and } \frac{d\phi}{dx}(L) = 0$$

$$(e) \quad \frac{d\phi}{dx}(0) = 0 \text{ and } \phi(L) = 0$$

$$* (f) \quad \phi(a) = 0 \text{ and } \phi(b) = 0 \text{ (You may assume that } \lambda > 0 \text{.)}$$

$$(g) \quad \phi(0) = 0 \text{ and } \frac{d\phi}{dx}(L) + \phi(L) = 0 \text{ (If necessary, see Sec. 5.8.)}$$

2.3.3. Consider the heat equation

$$\frac{\partial u}{\partial t} = k \frac{\partial^2 u}{\partial x^2},$$

subject to the boundary conditions

$$u(0, t) = 0 \quad \text{and} \quad u(L, t) = 0.$$

Solve the initial value problem if the temperature is initially

$$(a) \quad u(x, 0) = 6 \sin \frac{9\pi x}{L}$$

$$(b) \quad u(x, 0) = 3 \sin \frac{\pi x}{L} - \sin \frac{3\pi x}{L}$$

$$* (c) \quad u(x, 0) = 2 \cos \frac{3\pi x}{L}$$

$$(d) \quad u(x, 0) = \begin{cases} 1 & 0 < x \leq L/2 \\ 2 & L/2 < x < L \end{cases}$$

[Your answer in part (c) may involve certain integrals that do not need to be evaluated.]

2.3.4. Consider

$$\frac{\partial u}{\partial t} = k \frac{\partial^2 u}{\partial x^2},$$

subject to $u(0, t) = 0$, $u(L, t) = 0$, and $u(x, 0) = f(x)$.

*(a) What is the total heat energy in the rod as a function of time?

(b) What is the flow of heat energy out of the rod at $x = 0$? at $x = L$?

*(c) What relationship should exist between parts (a) and (b)?

2.3.5. Evaluate (be careful if $n = m$)

$$\int_0^L \sin \frac{n\pi x}{L} \sin \frac{m\pi x}{L} dx \quad \text{for } n > 0, m > 0.$$

Use the trigonometric identity

$$\sin a \sin b = \frac{1}{2} [\cos(a - b) - \cos(a + b)].$$

*2.3.6. Evaluate

$$\int_0^L \cos \frac{n\pi x}{L} \cos \frac{m\pi x}{L} dx \quad \text{for } n \geq 0, m \geq 0.$$

Use the trigonometric identity

$$\cos a \cos b = \frac{1}{2} [\cos(a + b) + \cos(a - b)].$$

(Be careful if $a - b = 0$ or $a + b = 0$.)

2.3.7. Consider the following boundary value problem (if necessary, see Sec. 2.4.1):

$$\frac{\partial u}{\partial t} = k \frac{\partial^2 u}{\partial x^2} \quad \text{with} \quad \frac{\partial u}{\partial x}(0, t) = 0, \frac{\partial u}{\partial x}(L, t) = 0, \quad \text{and} \quad u(x, 0) = f(x).$$

(a) Give a one-sentence physical interpretation of this problem.

(b) Solve by the method of separation of variables. First show that there are no separated solutions which exponentially grow in time. [Hint: The answer is

$$u(x, t) = A_0 + \sum_{n=1}^{\infty} A_n e^{-\lambda_n kt} \cos \frac{n\pi x}{L}]$$

What is λ_n ?

(c) Show that the initial condition, $u(x, 0) = f(x)$, is satisfied if

$$f(x) = A_0 + \sum_{n=1}^{\infty} A_n \cos \frac{n\pi x}{L}.$$

(d) Using Exercise 2.3.6, solve for A_0 and A_n ($n \geq 1$).

(e) What happens to the temperature distribution as $t \rightarrow \infty$? Show that it approaches the steady-state temperature distribution (see Sec. 1.4).

*2.3.8. Consider

$$\frac{\partial u}{\partial t} = k \frac{\partial^2 u}{\partial x^2} - \alpha u.$$

This corresponds to a one-dimensional rod either with heat loss through the lateral sides with outside temperature 0° ($\alpha > 0$, see Exercise 1.2.4) or with insulated lateral sides with a heat sink proportional to the temperature. Suppose that the boundary conditions are

$$u(0, t) = 0 \quad \text{and} \quad u(L, t) = 0.$$

(a) What are the possible equilibrium temperature distributions if $\alpha > 0$?

(b) Solve the time-dependent problem [$u(x, 0) = f(x)$] if $\alpha > 0$. Analyze the temperature for large time ($t \rightarrow \infty$) and compare to part (a).

*2.3.9. Redo Exercise 2.3.8 if $\alpha < 0$. [Be especially careful if $-\alpha/k = (n\pi/L)^2$.]

2.3.10. For two- and three-dimensional vectors, the fundamental property of dot products, $\mathbf{A} \cdot \mathbf{B} = |\mathbf{A}||\mathbf{B}| \cos \theta$, implies that

$$|\mathbf{A} \cdot \mathbf{B}| \leq |\mathbf{A}||\mathbf{B}|. \quad (2.3.44)$$

In this exercise we generalize this to n -dimensional vectors and functions, in which case (2.3.44) is known as **Schwarz's inequality**. [The names of Cauchy and Buniakovsky are also associated with (2.3.44).]

(a) Show that $|\mathbf{A} - \gamma\mathbf{B}|^2 > 0$ implies (2.3.44), where $\gamma = \mathbf{A} \cdot \mathbf{B} / \mathbf{B} \cdot \mathbf{B}$.

(b) Express the inequality using both

$$\mathbf{A} \cdot \mathbf{B} = \sum_{n=1}^{\infty} a_n b_n = \sum_{n=1}^{\infty} a_n c_n \frac{b_n}{c_n}.$$

* (c) Generalize (2.3.44) to functions. [Hint: Let $\mathbf{A} \cdot \mathbf{B}$ mean the integral $\int_0^L A(x)B(x) dx$.]

2.3.11. Solve Laplace's equation inside a rectangle:

$$\nabla^2 u = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$$

subject to the boundary conditions

$$\begin{aligned} u(0, y) &= g(y) & u(x, 0) &= 0 \\ u(L, y) &= 0 & u(x, H) &= 0. \end{aligned}$$

(Hint: If necessary, see Sec. 2.5.1.)