

heat exchange and exit times

Jean-Luc Thiffeault

Department of Mathematics
University of Wisconsin – Madison

with Florence Marcotte, William R. Young, Charles R. Doering

Mathematics of Turbulence Reunion, Lake Arrowhead, CA
5 June 2017

Supported by NSF grant CMMI-1233935



WISCONSIN
UNIVERSITY OF WISCONSIN-MADISON


advection–diffusion equation in a bounded region



Advection and diffusion of heat in a **bounded region** Ω , with Dirichlet boundary conditions:

$$\partial_t \theta + \mathbf{u} \cdot \nabla \theta = D \Delta \theta, \quad \mathbf{u} \cdot \hat{\mathbf{n}}|_{\partial \Omega} = 0, \quad \theta|_{\partial \Omega} = 0,$$

with $\nabla \cdot \mathbf{u} = 0$ and $\theta(\mathbf{x}, t) \geq 0$.




This is the **heat exchanger** configuration: given an initial distribution of heat, it is fluxed away through the cooled boundaries.

This happens through **diffusion** (conduction) alone, but is greatly aided by **stirring**.

heat exchangers



Our domain will be a 2D cross-section of a traditional coil.



Write $\langle \cdot \rangle$ for an integral over Ω .

$$\langle \cdot \rangle := \int_{\Omega} \cdot \, dV$$

The **rate of heat loss is equal to the flux** through the boundary $\partial\Omega$:

$$\partial_t \langle \theta \rangle = D \int_{\partial\Omega} \nabla \theta \cdot \hat{\mathbf{n}} \, dS =: -F[\theta] \leq 0. \quad *$$

Goal: find velocity fields \mathbf{u} that maximize the heat flux.

Note that $*$ is not so good for this, since velocity does not appear.

The role of \mathbf{u} is to **increase gradients** near the boundary. What it does internally is not directly relevant. **This is in contrast to the traditional Neumann IVP** (chaotic mixing, etc).



Take **steady velocity** $\mathbf{u}(\mathbf{x})$. The **mean exit time** $\tau(\mathbf{x})$ of a Brownian particle initially at \mathbf{x} satisfies

$$-\mathbf{u} \cdot \nabla \tau = D \Delta \tau + 1, \quad \tau|_{\partial\Omega} = 0,$$

This is a steady advection–diffusion equation with velocity $-\mathbf{u}$ and source 1.

Intuitively, a **small integrated mean exit time** $\langle \tau \rangle = \|\tau\|_1$ implies that the velocity is efficient at taking heat out of the system.

The mean exit time equation is much nicer than the equation for the concentration: it is **steady**, and it applies for any **initial concentration** $\theta_0(\mathbf{x})$.

relationship between exit time and mean temperature

Recall that $\langle \cdot \rangle$ is an integral over space, and take $\langle \theta_0 \rangle = 1$. The quantity

$$\int_0^\infty \langle \theta \rangle dt$$

is a **cooling time**. **Smaller is better** for good heat exchange.

We have the rigorous bounds

$$\int_0^\infty \langle \theta \rangle dt \leq \|\tau\|_\infty \quad \int_0^\infty \langle \theta \rangle dt \leq \|\tau\|_1 \|\theta_0\|_\infty.$$

Thus, decreasing a norm like $\|\tau\|_1$ or $\|\tau\|_\infty$ will typically decrease the cooling time, as expected.

does stirring always help?



[Iyer, G., Novikov, A., Ryzhik, L., & Zlatoš, A. (2010). *SIAM J. Math. Anal.* **42** (6), 2484–2498]

Theorem (Iyer *et al.* 2010)

$\Omega \in \mathbb{R}^n$ bounded, $\partial\Omega \in C^1$. Then

$$\|\tau\|_{L^p(\Omega)} \leq \|\tau_0\|_{L^p(\mathcal{B})}, \quad 1 \leq p \leq \infty,$$

where $\mathcal{B} \in \mathbb{R}^n$ is a ball of the same volume as Ω , and τ_0 is the 'purely diffusive' solution, $0 = D\Delta\tau_0 + 1$ on \mathcal{B} .

That is, measured in any norm, the exit time is maximized for a disk with no stirring. So **for a disk stirring always helps**, or at least isn't harmful.

They also prove that, surprisingly, if Ω is not a disk, then it's **always** possible to make $\|\tau\|_{L^\infty(\Omega)}$ **increase** by stirring. (Related to unmixing flows? [IMA 2010 gang; see review Thiffeault (2012)])



Let's formulate an optimization problem to find the best incompressible \mathbf{u} .

Advection–diffusion operator and its **adjoint**:

$$\mathcal{L} := \mathbf{u} \cdot \nabla - D\Delta, \quad \mathcal{L}^\dagger = -\mathbf{u} \cdot \nabla - D\Delta.$$

Minimize $\langle \tau \rangle$ over steady $\mathbf{u}(\mathbf{x})$ with fixed total kinetic energy $E = \frac{1}{2} \|\mathbf{u}\|_2^2$.

The functional to optimize:

$$\mathcal{F}[\tau, \mathbf{u}, \vartheta, \mu, p] = \langle \tau \rangle - \langle \vartheta (\mathcal{L}^\dagger \tau - 1) \rangle + \frac{1}{2} \mu (\|\mathbf{u}\|_2^2 - 2E) - \langle p \nabla \cdot \mathbf{u} \rangle$$

Here ϑ , μ , p are **Lagrange multipliers**.

Introduce streamfunction Ψ to satisfy $\nabla \cdot \mathbf{u} = 0$:

$$u_x = -\partial_y \psi, \quad u_y = \partial_x \psi.$$

The variational problem gives the [Euler–Lagrange equations](#)

$$\begin{aligned} \mathcal{L}^\dagger \tau &= 1, & \tau|_{\partial\Omega} &= 0; \\ \mathcal{L} \vartheta &= 1, & \vartheta|_{\partial\Omega} &= 0; \\ \mu \Delta \psi &= J(\tau, \vartheta), & \psi|_{\partial\Omega} &= 0; \\ \langle |\nabla \psi|^2 \rangle &= 2E, \end{aligned}$$

with the Jacobian

$$J(\tau, \vartheta) := (\nabla \tau \times \nabla \vartheta) \cdot \hat{\mathbf{z}}.$$



Transform to new functions η, ξ

$$\tau = \tau_0 + \frac{1}{2}(\eta + \xi), \quad \vartheta = \tau_0 + \frac{1}{2}(\eta - \xi)$$

where recall that τ_0 is the solution without flow (purely diffusive).

Then by using the Euler–Lagrange equations we can eventually show

$$\langle \tau \rangle = \langle \tau_0 \rangle - \frac{1}{4} \langle |\nabla \xi|^2 \rangle - \frac{1}{4} \langle |\nabla \eta|^2 \rangle.$$

Hence, solutions to E–L equations cannot make $\langle \tau \rangle$ increase. So stirring is always better than not stirring.

For a disk the purely diffusive solution is $\tau_0 = \frac{1}{4}(1 - r^2)$. We then make the *ansatz*

$$\xi = \sqrt{2\mu} B(r) \cos m\theta, \quad \eta = B(r) \sin m\theta, \quad \psi = \xi / \sqrt{2\mu},$$

and look for solutions of that form.

Inserting this into the full system gives solutions provided the radial functions $B(r)$ satisfy the **nonlinear eigenvalue problem**

$$r^2 B'' + rB' + (r^2 \lambda - m^2)B = \frac{1}{2} m^2 B^3, \quad \lambda = m / \sqrt{2\mu}.$$



The left-hand side is Bessel's equation.

Note that it is rather unusual for such a linear-type ansatz to give nonlinear solutions. We also have no guarantee that this is the **true optimal solution**.


For **small energy** E , exact solution in terms of Bessel functions $J_m(\rho_{mn}r)$, where ρ_{mn} are zeros:

$$\langle \tau \rangle / \langle \tau_0 \rangle = 1 - (4m^2 / \pi \rho_{mn}^4) E + O(E^2).$$


Pick the solution with the smallest $\langle \tau \rangle$: $m = 2, n = 1$ for all $E \ll 1$:



Numerical solution with Matlab's **bvp5c**, using a continuation method:




Larger m worse at small E , then better, then maybe worse again?



Three regions:

- Stagnation zone (SZ)
- Bulk
- Peripheral boundary layer (PBL)

structure of the radial solution $B(r)$ for large E





Rescaled variables $B = E^\alpha \tilde{B}$ and $\lambda = E^\beta \tilde{\lambda}$:

$$r^2 \tilde{B}'' E^\alpha + r \tilde{B}' E^\alpha + r^2 \tilde{\lambda} \tilde{B} E^{\alpha+\beta} - m^2 \tilde{B} E^\alpha = \frac{1}{2} m^2 \tilde{B}^3 E^{3\alpha}.$$

Outside the boundary layer, the large- E balance must occur between the terms $r^2 \tilde{\lambda} \tilde{B} E^{\alpha+\beta}$ and $\frac{1}{2} m^2 \tilde{B}^3 E^{3\alpha}$, so $\beta = 2\alpha$.

This gives the outer solution

$$B_{\text{outer}} = E^\alpha \tilde{B} = \sqrt{2/m^3 \tilde{\lambda}} E^\alpha r.$$

(This does not include the stagnation zone in the center. Neglect for now.)

Cannot satisfy $B_{\text{outer}}(1) = 0$: need **boundary layer**.



Inner variable $r = 1 - \epsilon\rho$:

$$\begin{aligned} \frac{(1 - \epsilon\rho)^2}{\epsilon^2} \bar{B}'' E^\alpha + \frac{(1 - \epsilon\rho)}{\epsilon} \bar{B}' E^\alpha + (1 - \epsilon\rho)^2 \tilde{\lambda} \bar{B} E^{3\alpha} - m^2 \bar{B} E^\alpha \\ = \frac{1}{2} m^2 \bar{B}^3 E^{3\alpha}. \end{aligned}$$

Dominant balance: highest derivative with $E^\alpha = \epsilon^{-1}$:

$$\bar{B}'' + \tilde{\lambda} \bar{B} = \frac{1}{2} m^2 \bar{B}^3.$$

This has an exact **tanh** solution, which after matching with the outer solution as $\rho \rightarrow \infty$ gives

$$B_{\text{inner}} = \sqrt{2\tilde{\lambda}/m^2} E^\alpha \tanh\left(\sqrt{\lambda/2}\rho\right)$$




Finally we apply the energy constraint, which reads

$$\begin{aligned} \frac{2E}{\pi} &= \int_0^1 \left\{ rB'^2 + \frac{m^2}{r} B^2 \right\} dr \\ &= \int_0^{1-\delta} \left\{ rB_{\text{outer}}'^2 + \frac{m^2}{r} B_{\text{outer}}^2 \right\} dr + \int_{1-\delta}^1 \left\{ B_{\text{inner}}'^2 + m^2 B_{\text{inner}}^2 \right\} dr. \end{aligned}$$

We skip the details, but dominant balance requires $\alpha = 1/3$, and so $\beta = 2\alpha = 2/3$.


The optimal integrated exit time thus scales as $m^{-2/3} E^{-1/3}$.

large- E case: asymptotics at fixed m



Fixed- E asymptotic optimal $\langle \tau \rangle$ seems to decrease to zero as $m^{-2/3}$. This implies no optimal flow, since arbitrarily efficient at large m . Not so!

large- E , large- m case



To truly capture the optimal solution, have to let $m \sim E^{1/4}$.
This is the **dashed line** (envelope).

- Transport in heat exchangers has a very different character than ‘freely-decaying’ problem.
- Using the probabilistic **mean exit time** formulation simplifies the problem. (Idea came from Iyer et al. 2010.)
- Optimal solutions for \mathbf{u} are reminiscent of **Dean flow**.
- At small energy optimal solution has $m = 2$, $n = 1$.
- At larger energy there is a boundary layer, which enhances the heat transfer or decreases exit time: $\langle \tau \rangle \sim m^{-2/3} E^{-1/3}$.
- This asymptotic solution breaks down when m gets too large. The **stagnation zone** becomes larger and penalizes large m .
- A distinguished limit in m gives $\langle \tau \rangle \sim E^{-1/2}$.
- Generalizations: use different norms, spatial weight. . .



- Iyer, G., Novikov, A., Ryzhik, L., & Zlatoš, A. (2010). *SIAM J. Math. Anal.* **42** (6), 2484–2498.
- Thiffeault, J.-L. (2012). *Nonlinearity*, **25** (2), R1–R44.