
Mixing with Ghost Rods

Jean-Luc Thiffeault

Matthew Finn

Emmanuelle Guillard

<http://www.ma.imperial.ac.uk/~jeanluc>

Department of Mathematics

Imperial College London


Experiment of Boyland, Aref, & Stremler




[P. L. Boyland, H. Aref, and M. A. Stremler, *J. Fluid Mech.* **403**, 277 (2000)]

[P. L. Boyland, M. A. Stremler, and H. Aref, *Physica D* **175**, 69 (2003)]

The Connection with Braiding



Generators of the n -Braid Group



A generator

$$\sigma_i, \quad i = 1, \dots, n-1$$

is the clockwise interchange of the i th and $(i+1)$ th rod.

The generators obey the **presentation**


$$\sigma_{i+1} \sigma_i \sigma_{i+1} = \sigma_i \sigma_{i+1} \sigma_i$$

$$\sigma_i \sigma_j = \sigma_j \sigma_i, \quad |i - j| > 1$$


These generators are used to characterise the motion of the rods.

The Two BAS Stirring Protocols

$\sigma_1\sigma_2$ protocol



$\sigma_1^{-1}\sigma_2$ protocol




[P. L. Boyland, H. Aref, and M. A. Stremler, *J. Fluid Mech.* **403**, 277 (2000)]

Topological Entropy of a Braid

Practically speaking, the topological entropy of a braid is a **lower bound** on the **line-stretching exponent** of the flow!

This is reasonable:



In practice, we compute the topological entropy of a braid using a **train-track** algorithm due to Bestvina & Handel. The end result is a **transition matrix** showing the how each edge is mapped under the action of the braid.

[M. Bestvina and M. Handel, *Topology* **34**, 109 (1995)]

The Difference between BAS's Two Protocols

- The matrices associated the generators have eigenvalues on the unit-circle (but their product doesn't necessarily).
- The first (**bad**) stirring protocol has eigenvalues on the unit circle
- The second (**good**) protocol has largest eigenvalue $(3 + \sqrt{5})/2 = 2.6180$.
- So for the second protocol the length of a line joining the rods grows exponentially!
- That is, material lines have to stretch by at least a factor of 2.6180 each time we execute the protocol $\sigma_1^{-1}\sigma_2$.
- This is guaranteed to hold in some neighbourhood of the rods (**Thurston–Nielsen theorem**).

One Rod Mixer: The Kenwood Chef



More than a Mixer
at CHRISTMAS

and all through the year...

The Kenwood^{RESO}
Chef
Your Servant Madam!

The Kenwood 'Chef' not only
★ MIXES, KNEADS and WHISKS
but, with its wide range of optional attachments, it
★ LIQUIDISES, BLENDS, PUREES,
EXTRACTS JUICES
★ SLICES, MINCES, GRINDS, SIEVES
★ PEELS POTATOES and
★ Even OPENS CANS

Yes, you need the Kenwood 'Chef' in your home every day of the year. It makes every meal a festive occasion, with tempting dishes and drinks having the true professional touch.

What a wonderful Christmas gift . . . especially if it arrives in good time to prepare the Christmas spread!

Standard Pack: £28 7 6
complete with Mixing Bowl, K-Beater, Whisk, Dough Hook, Rubber Spatula and Plastic Dust Cover.
Optional attachments extra.

H.P. TERMS AVAILABLE


For illustrated literature of all Kenwood products and name of nearest dealer, write to

Other delightful Kenwood Gifts


<p>The ROTO-BLEND Gives you Wealth — Why? Because it gives you health and vitamin in seconds. This versatile machine, retains all the natural goodness of fruit and vegetables, giving you juices unobtainable without it. Versatile because you can add to the Roto-Blend a multi-purpose Blender attachment and high-speed Slicer and Shredder attachment. Standard Unit: £14 11 2</p>	<p>The KENMIX '55' The Continental Wonder! Does the work of hours in a few seconds, retaining vitamins and full flavours. So simple to use; blends fruits and vegetables, drinks cocktails and eggs, pulverises nuts and dry cereals, mixes drinks and confections. £14 12 7</p>	<p>The MINOR A really efficient Portable Food Mixer for only £9 18 0. It mixes, whips, beats and blends—in fact does everything a Food Mixer should, saving hours of wrist-aching work. Weighs only 3½ lbs., light enough to be used anywhere in the kitchen—at the stove, over the sink or in the table. £9 18 0</p>
---	---	--

For illustrated literature of all Kenwood products and name of nearest dealer, write to
KENWOOD ELECTRICS LTD., 26 North Audley Street, London, W.1

Poincaré Section




Stretching of Lines



Particle Orbits are Topological Obstacles

Choose any fluid particle orbit (green dot).



Material lines must bend around the orbit: **it acts just like a “rod”!**

[J-LT, *Phys. Rev. Lett.* **94**, 084502 (2005)]


Today: focus on **periodic orbits**.

How do they braid around each other?

Motion of Islands

Make a braid from the motion of the rod and the **periodic islands**.

Most (74%) of the topological entropy is accounted for by this braid.




Motion of Islands and Unstable Periodic Orbits

Now we also include **unstable** periodic orbits as well as the stable ones (islands).

Almost all (99%) of the topological entropy is accounted for by this braid.

But are the periodic orbits really “ghost rods”?


That is, do material lines really get out of their way?



Periodic Orbits as Ghost Rods

[movie 3: sf_periodic.avi]

Blowup of the Tip



So Which Orbits Make Good Rods?

A good ghost rod should “look” like a real rod: material lines wrap around it.

Look at linearisation of the period-1 map:

Two eigenvalues Λ and Λ^{-1} (Floquet multipliers),
with $|\Lambda| \geq |\Lambda|^{-1}$.

Λ complex	Elliptic	Good rod (obvious).
$ \Lambda > 1$	Hyperbolic	Crap rod.
$\Lambda = 1$	Parabolic	Good rod?

The parabolic points are the most interesting: they are associated with the crucial property of **folding**.

Normal Form near a Parabolic Point

$$X' = X + \text{quadratic terms}$$

$$Y' = Y + \text{quadratic terms}$$

or

$$X' = X + Y + X^2$$


$$Y' = Y + X^2$$

The first of these does not lead to folding.

The coefficient of X^2 determines the “size” of the rod, which is a function of time.

[[movie 4: folditer.avi](#)]

Curvature of the Tip at the Periodic Orbit



Conclusions

- Topological chaos involves moving **obstacles** in a 2D flow, which create nontrivial braids.
- Periodic orbits make perfect obstacles (in periodic flows).
- This is a good way to “explain” the chaos in a flow — accounts for stretching of material lines.
- Islands (elliptic orbits) look just like rods, and parabolic orbits can look a lot like rods.
- No need for infinitesimal separation of trajectories or derivatives of the velocity field — this is an inherently **global** description.
- The size of the rods is important — for islands this is obvious but for parabolic points the apparent size is a function of time.