

Topological optimization of rod-stirring devices

Jean-Luc Thiffeault

Department of Mathematics
University of Wisconsin – Madison

SIAM–UW Seminar, 13 April 2011

Collaborators:

Matthew Finn	University of Adelaide
Phil Boyland	University of Florida

The taffy puller




[Photo and movie by M. D. Finn.]

[movie 1]

The mixograph

Model experiment for kneading bread dough:



[Department of Food Science, University of Wisconsin. Photos by J-LT.]

Planetary mixers

In food processing, **rods** are often used for stirring.




[movie 2] ©BLT Inc.

Experiment of Boyland, Aref & Stremler




[movie 3] [movie 4]




[P. L. Boyland, H. Aref, and M. A. Stremler, *J. Fluid Mech.* **403**, 277 (2000)]

Braid description of taffy puller



The three rods of the taffy puller in a space-time diagram. Defines a braid on $n = 3$ strands, $\sigma_1^2 \sigma_2^{-2}$ (three periods shown).

Braid description of mixograph



$$\sigma_3 \sigma_2 \sigma_3 \sigma_5 \sigma_6^{-1} \sigma_2 \sigma_3 \sigma_4 \sigma_3 \sigma_1^{-1} \sigma_2^{-1} \sigma_5$$

braid on B_7 , the braid group on 7 strands.

Topological entropy of a braid

Burau representation for 3-braids:

$$[\sigma_1] = \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}, \quad [\sigma_2] = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix},$$

$$[\sigma_1^{-1} \sigma_2] = [\sigma_1^{-1}] \cdot [\sigma_2] = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \cdot \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}.$$

This matrix has **spectral radius** $(3 + \sqrt{5})/2$ (**Golden Ratio²**), and hence the topological entropy is $\log[(3 + \sqrt{5})/2]$.

This is the growth rate of a 'rubber band' caught on the rods.

This matrix trick only works for 3-braids, unfortunately.

Optimizing over generators

- Entropy can grow without bound as the length of a braid increases;
- A proper definition of optimal entropy requires a **cost** associated with the braid.
- Divide the entropy by the **smallest number of generators** required to write the braid word.
- For example, the braid $\sigma_1^{-1} \sigma_2$ has entropy $\log[(3 + \sqrt{5})/2]$ and consists of two generators.
- Its **Topological Entropy Per Generator (TEPG)** is thus $\frac{1}{2} \log[(3 + \sqrt{5})/2] = \log[\text{Golden Ratio}]$.
- Assume all the generators are used (**stronger: irreducible**).

Optimal braid

- In B_3 and B_4 , the optimal TEPG is $\log[\text{Golden Ratio}]$.
- Realized by $\sigma_1^{-1}\sigma_2$ and $\sigma_1^{-1}\sigma_2\sigma_3^{-1}\sigma_2$, respectively.
- In B_n , $n > 4$, the optimal TEPG is $< \log[\text{Golden Ratio}]$.

Why? Recall Burau representation:

$$[\sigma_1] = \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}, \quad [\sigma_2] = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix},$$


Its spectral radius provides a lower bound on entropy. However,

$$|[\sigma_1]| = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, \quad |[\sigma_2]| = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix},$$

provides an upper bound! Need to find **Joint Spectral Radius**.

Periodic array of rods


- Consider periodic lattice of rods.
- Move all the rods such that they execute $\sigma_1 \sigma_2^{-1}$ with their neighbor (Boyland et al., 2000).





- The entropy per 'switch' is $\log(1 + \sqrt{2})$, the **Silver Ratio**!
- This is **optimal** for a periodic lattice of two rods (follows from D'Alessandro et al. (1999)).
- Also optimal if we assign cost by **simultaneous operation**.

Silver mixers

- The designs with entropy given by the Silver Ratio can be realized with simple gears.
- All the rods move at once: very efficient.




Build it!




[movie 6] [movie 7]

Experiment: Silver mixer with four rods



Silver mixer with six rods



[movie 8]

Conclusions

- Having rods undergo ‘braiding’ motion guarantees a minimal amount of entropy ([stretching of material lines](#)).
- Can optimize to find the best rod motions, but depends on choice of ‘cost function.’
- For two natural cost functions, the **Golden Ratio** and **Silver Ratio** pop up!

References

- Boyland, P. L., Aref, H. & Stremler, M. A. 2000 Topological fluid mechanics of stirring. *J. Fluid Mech.* **403**, 277–304.
- D'Alessandro, D., Dahleh, M. & Mezić, I. 1999 Control of mixing in fluid flow: A maximum entropy approach. *IEEE Transactions on Automatic Control* **44**, 1852–1863.
- Finn, M. D., & Thiffeault, J.-L. 2010 Topological optimisation of rod-stirring devices. *SIAM Review*, in press.
- Gouillart, E., Finn, M. D. & Thiffeault, J.-L. 2006 Topological Mixing with Ghost Rods. *Phys. Rev. E* **73**, 036311.
- Moussafir, J.-O. 2006 On the Entropy of Braids. *Func. Anal. and Other Math.* **1**, 43–54.
- Thiffeault, J.-L. 2005 Measuring Topological Chaos. *Phys. Rev. Lett.* **94**, 084502.
- Thiffeault, J.-L. & Finn, M. D. 2006 Topology, Braids, and Mixing in Fluids. *Phil. Trans. R. Soc. Lond. A* **364**, 3251–3266.
- Thiffeault, J.-L., Finn, M. D., Gouillart, E. & Hall, T. 2008 Topology of Chaotic Mixing Patterns. *Chaos* **18**, 033123.
- Thurston, W. P. 1988 On the geometry and dynamics of diffeomorphisms of surfaces. *Bull. Am. Math. Soc.* **19**, 417–431.