

Topological optimization

Jean-Luc Thiffeault


Department of Mathematics
University of Wisconsin – Madison

WIDDOW Seminar, Wisconsin Institute for Discovery
13 February 2012

Collaborators:

Matthew Finn	University of Adelaide
Erwan Lanneau	CPT Marseille
Phil Boyland	University of Florida

The taffy puller




[Photo and movie by M. D. Finn.]

[movie 1]

The mixograph

Model experiment for kneading bread dough:



[Department of Food Science, University of Wisconsin. Photos by J-LT.]


Planetary mixers

In food processing, **rods** are often used for stirring.



[movie 2] ©BLT Inc.


Experiment of Boyland, Aref & Stremler



[movie 3] [movie 4]


[P. L. Boyland, H. Aref, and M. A. Stremler, *J. Fluid Mech.* **403**, 277 (2000)]

Braid description of taffy puller



The three rods of the taffy puller in a space-time diagram. Defines a braid on $n = 3$ strands, $\sigma_1^2 \sigma_2^{-2}$ (three periods shown).

Braid description of mixograph



$$\sigma_3 \sigma_2 \sigma_3 \sigma_5 \sigma_6^{-1} \sigma_2 \sigma_3 \sigma_4 \sigma_3 \sigma_1^{-1} \sigma_2^{-1} \sigma_5$$

braid on B_7 , the braid group on 7 strands.

Topological entropy of a braid

Burau representation for 3-braids:

$$[\sigma_1] = \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}, \quad [\sigma_2] = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix},$$

$$[\sigma_1^{-1} \sigma_2] = [\sigma_1^{-1}] \cdot [\sigma_2] = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \cdot \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}.$$

This matrix has **spectral radius** $(3 + \sqrt{5})/2$ (**Golden Ratio²**), and hence the topological entropy is $\log[(3 + \sqrt{5})/2]$.

This is the growth rate of a 'rubber band' caught on the rods.

This matrix trick only works for 3-braids, unfortunately.

Optimizing over generators

- Entropy can grow without bound as the length of a braid increases;
- A proper definition of optimal entropy requires a **cost** associated with the braid.
- Divide the entropy by the **smallest number of generators** required to write the braid word.
- For example, the braid $\sigma_1^{-1} \sigma_2$ has entropy $\log[(3 + \sqrt{5})/2]$ and consists of two generators.
- Its **Topological Entropy Per Generator (TEPG)** is thus $\frac{1}{2} \log[(3 + \sqrt{5})/2] = \log[\text{Golden Ratio}]$.
- Assume all the generators are used (**stronger: irreducible**).

Optimal braid

- In B_3 and B_4 , the optimal TEPG is $\log[\text{Golden Ratio}]$.
- Realized by $\sigma_1^{-1}\sigma_2$ and $\sigma_1^{-1}\sigma_2\sigma_3^{-1}\sigma_2$, respectively.
- In B_n , $n > 4$, the optimal TEPG is $< \log[\text{Golden Ratio}]$.

Why? Recall Burau representation:

$$[\sigma_1] = \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}, \quad [\sigma_2] = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix},$$


Its spectral radius provides a lower bound on entropy. However,

$$|[\sigma_1]| = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, \quad |[\sigma_2]| = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix},$$

provides an upper bound! Need to find **Joint Spectral Radius**.

Periodic array of rods


- Consider periodic lattice of rods.
- Move all the rods such that they execute $\sigma_1 \sigma_2^{-1}$ with their neighbor (Boyland et al., 2000).





- The entropy per 'switch' is $\log(1 + \sqrt{2})$, the **Silver Ratio**!
- This is **optimal** for a periodic lattice of two rods (follows from D'Alessandro et al. (1999)).
- Also optimal if we assign cost by **simultaneous operation**.

Silver mixers

- The designs with entropy given by the Silver Ratio can be realized with simple gears.
- All the rods move at once: very efficient.




Build it!




[movie 6] [movie 7]

[M. D. Finn and J-LT, *SIAM Review* **53**, 723 (2011)]

Experiment: Silver mixer with four rods



Silver mixer with six rods



[movie 8]

The Minimizer problem

- On a given surface of genus g , which pA has the least λ ?
- If the foliation is orientable (vector field), then things are much simpler;
- Action of the pA on first homology captures dilatation λ ;
- Polynomials of degree $2g$;
- Procedure:
 - We have a guess for the minimizer;
 - Find all integer-coefficient, reciprocal polynomials that could have smaller largest root;
 - Show that they can't correspond to pA s;
 - For the smallest one that can, construct pA .

Newton's formulas

We need an efficient way to bound the number of polynomials with largest root smaller than λ . Given a reciprocal polynomial

$$P(x) = x^{2g} + a_1 x^{2g-1} + \dots + a_2 x^2 + a_1 x + 1$$

we have Newton's formulas for the traces,

$$\mathrm{Tr}(\phi_*^k) = - \sum_{m=1}^{k-1} a_m \mathrm{Tr}(\phi_*^{k-m}) - ka_k,$$

where

- ϕ is a (hypothetical) pA associated with $P(x)$;
- ϕ_* is the matrix giving the action of the pA ϕ on first homology;
- $\mathrm{Tr}(\phi_*)$ is its trace.

Bounding the traces

The trace satisfies

$$|\mathrm{Tr}(\phi_*^k)| = \left| \sum_{m=1}^g (\lambda_m^k + \lambda_m^{-k}) \right| \leq g(r^k + r^{-k})$$

where λ_m are the roots of ϕ_* , and $r = \max_m(|\lambda_m|)$.

- Bound $\mathrm{Tr}(\phi_*^k)$ with $r < \lambda$, $k = 1, \dots, g$;
- Use these g traces and Newton's formulas to construct candidate $P(x)$;
- Overwhelming majority have fractional coeffs \rightarrow discard!
- Carefully check the remaining polynomials:
 - Is their largest root real?
 - Is it strictly greater than all the other roots?
 - Is it really less than λ ?
- Largest tractable case: $g = 8$ (10^{12} polynomials).

Lefschetz's fixed point theorem

This procedure still leaves a fair number of polynomials — though not enormous (10's to 100's, even for $g = 8$.)

The next step involves using [Lefschetz's fixed point theorem](#) to eliminate more polynomials:

$$L(\phi) = 2 - \text{Tr}(\phi_*) = \sum_{p \in \text{Fix}(\phi)} \text{Ind}(\phi, p)$$

where

- $L(\phi)$ is the Lefschetz number;
- $\text{Fix}(\phi)$ is set of fixed points of ϕ ;
- $\text{Ind}(\phi, p)$ is index of ϕ at p .

We can easily compute $L(\phi^k)$ for every iterate using Newton's formula.

Eliminating polynomials

Outline of procedure: for a surface of genus g ,

- Use the Euler–Poincaré formula to list possible singularity data for the foliations;
- For each singularity data, compute possible contributions to the index (depending on how the singularities and their separatrices are permuted);
- Check if index is consistent with Lefschetz's theorem.

With this, we can reduce the number of polynomials to one or two!

Minimizers for orientable foliations

g	polynomial	minimizer
2	$X^4 - X^3 - X^2 - X + 1$	$\simeq 1.72208$ †
3	$X^6 - X^4 - X^3 - X^2 + 1$	$\simeq 1.40127$
4	$X^8 - X^5 - X^4 - X^3 + 1$	$\simeq 1.28064$
5	$X^{10} + X^9 - X^7 - X^6 - X^5 - X^4 - X^3 + X + 1$	$\simeq 1.17628$ *
6	$X^{12} - X^7 - X^6 - X^5 + 1$	$\gtrsim 1.17628$
7	$X^{14} + X^{13} - X^9 - X^8 - X^7 - X^6 - X^5 + X + 1$	$\simeq 1.11548$
8	$X^{16} - X^9 - X^8 - X^7 + 1$	$\simeq 1.12876$

† Zhironv (1995)'s result; also for nonorientable [Lanneau–T];

* Lehmer's number; realized by Leininger (2004)'s pA;

- For genus 6 we have not explicitly constructed the pA;

- Genus 6 is the first **nondecreasing** case.

- Genus 7 and 8: pA's found by Aaber & Dunfield (2010) and Kin & Takasawa (2010b) [$g = 7$]; Hironaka (2009) [$g = 8$].

Conclusions

- Having rods undergo ‘braiding’ motion guarantees a minimal amount of entropy ([stretching of material lines](#)).
- Can optimize to find the best rod motions, but depends on choice of ‘cost function.’
- For two natural cost functions, the **Golden Ratio** and **Silver Ratio** pop up!
- Found orientable minimizer on surfaces of genus $g \leq 8$; only known nonorientable case is for genus 2.

References

- Aaber, J. & Dunfield, N.. 2010 Closed surface bundles of least volume. Preprint.
- Boylard, P. L., Aref, H. & Stremler, M. A. 2000 Topological fluid mechanics of stirring. *J. Fluid Mech.* **403**, 277–304.
- D'Alessandro, D., Dahleh, M. & Mezić, I. 1999 Control of mixing in fluid flow: A maximum entropy approach. *IEEE Transactions on Automatic Control* **44**, 1852–1863.
- Finn, M. D., & Thiffeault, J.-L. 2011 Topological optimization of rod-stirring devices. *SIAM Review* **53**, 623–743.
- Gouillart, E., Finn, M. D. & Thiffeault, J.-L. 2006 Topological Mixing with Ghost Rods. *Phys. Rev. E* **73**, 036311.
- Ham, J.-Y. & Song, W. T. 2007 The minimum dilatation of pseudo-Anosov 5-braids. *Experiment. Math.* **16**, 167–179.
- Hironaka, E. 2009 Small dilatation pseudo-Anosov mapping classes coming from the simplest hyperbolic braid. Preprint.
- Kin, E. & Takasawa, M. 2010a Pseudo-Anosovs on closed surfaces having small entropy and the Whitehead sister link exterior. Preprint.
- Kin, E. & Takasawa, M. 2010b Pseudo-Anosov braids with small entropy and the magic 3-manifold. Preprint.
- Lanneau, E. & Thiffeault, J.-L. 2011a On the minimum dilatation of pseudo-Anosov diffeomorphisms on surfaces of small genus. *Annales de l'Institut Fourier* **61**, 105-144.
- Lanneau, E. & Thiffeault, J.-L. 2011b On the minimum dilatation of braids on the punctured disc. *Geometriae Dedicata* **152**, 165–182.
- Leininger, C. J. 2004 On groups generated by two positive multi-twists: Teichmüller curves and Lehmer's number. *Geom. Topol.* **8**, 1301–1359.
- Moussafir, J.-O. 2006 On the Entropy of Braids. *Func. Anal. and Other Math.* **1**, 43–54.
- Song, W. T., Ko, K. H. & Los, J. E. 2002 Entropies of braids. *J. Knot Th. Ramifications* **11**, 647–666.
- Thiffeault, J.-L. & Finn, M. D. 2006 Topology, Braids, and Mixing in Fluids. *Phil. Trans. R. Soc. Lond. A* **364**, 3251–3266.
- Thiffeault, J.-L., Finn, M. D., Gouillart, E. & Hall, T. 2008 Topology of Chaotic Mixing Patterns. *Chaos* **18**, 033123.
- Thurston, W. P. 1988 On the geometry and dynamics of diffeomorphisms of surfaces. *Bull. Am. Math. Soc.* **19**, 417–431.
- Zhirov, A. Y. 1995 On the minimum dilation of pseudo-Anosov diffeomorphisms of a double torus. *Russ. Math. Surv.* **50**, 223–224.