

# The topology of fluid mixing

Jean-Luc Thiffeault

Department of Mathematics  
University of Wisconsin – Madison

Zhejiang University, Hangzhou, China

7 June 2013

Supported by NSF grants DMS-0806821 and CMMI-1233935



# the taffy puller



Taffy is a type of candy.

Needs to be **pulled**: this aerates it and makes it lighter and chewier.

We can assign a **growth**: length multiplier per period.

(Here  $(1 + \sqrt{2})^2$ ... more later.)

[movie by M. D. Finn]

play movie



# making candy cane



[*Wired*: This Is How You Craft 16,000 Candy Canes in a Day]

# four-pronged taffy puller




play movie


<http://www.youtube.com/watch?v=Y7t1LHdsquVM>

[studied in detail by Halbert & Yorke (2013)]

Experimental device for kneading bread dough:



play movie



[Department of Food Science, University of Wisconsin. Photos by J-LT.]

# the mixograph as a braid



Encode the topological information as a sequence of **generators of the Artin braid group  $B_n$** .

Equivalent to the 7-braid


$$\sigma_3 \sigma_2 \sigma_3 \sigma_5 \sigma_6^{-1} \sigma_2 \sigma_3 \sigma_4 \sigma_3 \sigma_1^{-1} \sigma_2^{-1} \sigma_5$$

The **growth** is the largest root of

$$x^8 - 4x^7 - x^6 + 4x^4 - x^2 - 4x + 1$$


$$\simeq 4.186$$

Compare to taffy pullers:  $\boxed{5.828}$





play movie



play movie

[Boyland, P. L., Aref, H., & Stremler, M. A. (2000). *J. Fluid Mech.* **403**, 277–304; Simulations by M. D. Finn, S. E. Tumas, and J-LT.]



Periodic stirring protocols in two dimensions can be described by a **homeomorphism**  $\varphi : \mathcal{S} \rightarrow \mathcal{S}$ , where  $\mathcal{S}$  is a surface.

For instance, in a closed circular container,

- $\varphi$  describes the mapping of fluid elements after one full period of stirring, obtained by solving the Stokes equation;
- $\mathcal{S}$  is the **disc** with holes in it, corresponding to the stirring rods.

Goal: **Topological characterization of  $\varphi$ .**


[The theory extends to handlebodies, but not as relevant for applications. . . ]



$\varphi$  and  $\psi$  are **isotopic** if  $\psi$  can be continuously 'reached' from  $\varphi$  without moving the rods. Write  $\varphi \simeq \psi$ .

(Defines **isotopy classes**.)

Convenient to think of isotopy in terms of material loops. Isotopic maps act the same way on loops (up to continuous deformation).



(Loops will always mean **essential** loops.)

## Theorem

$\varphi$  is isotopic to a homeomorphism  $\psi$ , where  $\psi$  is in one of the following three categories:

**finite-order** for some integer  $k > 0$ ,  $\psi^k \simeq \text{identity}$ ;

**reducible**  $\psi$  leaves invariant a disjoint union of essential simple closed curves, called *reducing curves*;

**pseudo-Anosov**  $\psi$  leaves invariant a pair of transverse measured **singular foliations**,  $\mathcal{F}^u$  and  $\mathcal{F}^s$ , such that  $\psi(\mathcal{F}^u, \mu^u) = (\mathcal{F}^u, \lambda \mu^u)$  and  $\psi(\mathcal{F}^s, \mu^s) = (\mathcal{F}^s, \lambda^{-1} \mu^s)$ , for **dilatation**  $\lambda > 1$ .

The three categories characterize the **isotopy class** of  $\varphi$ .

We want **pseudo-Anosov** for good mixing.




- Consider a **motion of stirring elements**, such as rods.
- Determine if the motion is **isotopic to a pseudo-Anosov mapping**.
- **Compute** topological quantities, such as foliation, entropy, etc.
- **Analyze** and **optimize**.

# train tracks: computing entropy and foliations



play movie


'Figure-8' motion:  $\sigma_2^{-2}\sigma_1^2$



[Guillart *et al.* (2007)]

Thurston introduced **train tracks** as a way of characterizing the measured foliation. The name stems from the 'cusps' that look like train switches.

# train track map for figure-eight



$$a \mapsto a\bar{2}\bar{a}\bar{1}ab\bar{3}\bar{b}\bar{a}1a, \quad b \mapsto \bar{2}\bar{a}\bar{1}ab$$

Easy to show that this map is **efficient**: under repeated iteration, cancellations of the type  $a\bar{a}$  or  $b\bar{b}$  never occur.

[There are algorithms, such as Bestvina & Handel (1995), to find efficient train tracks. (Toby Hall has an implementation in C++.)]



As the TT map is iterated, the number of symbols grows exponentially, at a rate given by the **topological entropy**,  $\log \lambda$ . This is a lower bound on the **minimal length of a material line** caught on the rods.


Find from the TT map by **Abelianizing**: count the number of occurrences of  $a$  and  $b$ , and write as matrix:

$$\begin{pmatrix} a \\ b \end{pmatrix} \mapsto \begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix}$$

The largest eigenvalue of the matrix is  $\lambda = (1 + \sqrt{2})^2 \simeq 5.83$ . Hence, asymptotically, the length of the 'blob' is multiplied by 5.83 for each full stirring period.


[This is the growth for the 3 and 4-pronged taffy pullers.]

- Consider periodic lattice of rods.
- Move all the rods such that they execute the Boyland *et al.* (2000) rod motion (J-LT & Finn, 2006; Finn & J-LT, 2011).



- The dilatation per period is  $\chi^2$ , where  $\chi = 1 + \sqrt{2}$  is the **Silver Ratio!**
- This is **optimal** for a periodic lattice of two rods (Follows from D'Alessandro *et al.* (1999)).


- The designs with dilatation given by the silver ratio can be realized with simple gears.
- All the rods move at once: very efficient.







play movie

[play movie](#)

[See Finn, M. D. & J-LT (2011). *SIAM Rev.* **53** (4), 723–743 for proofs, heavily influenced by work on  $\pi_1$ -stirrers of Boyland, P. L. & Harrington, J. (2011). *Algeb. Geom. Topology*, **11** (4), 2265–2296.]

# ghost rods ('tiges fantômes')




Topological analysis can be done on other objects than rods – for instance, **islands** or **unstable periodic orbits**.

We simply follow the islands and examine the braid they form, which gives us bounds on **topological entropy**.


In this framework we call the islands **ghost rods**.

[Gouillart, E., Finn, M. D., & J-LT (2006). *Phys. Rev. E*, **73**, 036311]

[implemented by Stremmer & Chen (2007); J-LT *et al.* (2009); Binder (2010); Stremmer *et al.* (2011)]



One of the best examples of ghost rods is from Stremler *et al.* (2011):




The islands are made to follow the  $\sigma_2\sigma_1^{-1}$  stirring protocol by clever wall motions! (viscous Stokes flow)

[Stremler, M. A., Ross, S. D., Grover, P., & Kumar, P. (2011). *Phys. Rev. Lett.* **106**, 114101]

play movie

# oceanic float trajectories





What can we measure?

- single-particle dispersion (not a good use of all data)
- correlation functions (what do they mean?)
- Lyapunov exponents (some luck needed!)

Another possibility:

Compute the **braid group generators**  $\sigma_i$  for the float trajectories (convert to a sequence of symbols), then look at how loops grow. Obtain a **topological entropy** for the motion (similar to Lyapunov exponent).



It is well-known that the entropy can be obtained by applying the motion of the punctures to a closed curve (loop) repeatedly, and measuring the growth of the length of the loop (Bowen, 1978).

The problem is twofold:


- 1 Need to keep track of the loop, since its length is growing exponentially;
- 2 Need a simple way of transforming the loop according to the motion of the punctures.

However, simple closed curves are easy objects to manipulate in 2D. Since they cannot self-intersect, we can describe them **topologically** with very few numbers.

# solution to problem 1: loop coordinates




What saves us is that a closed loop can be uniquely reconstructed from the number of intersections with a set of curves. For instance, the [Dybnikov coordinates](#) involve intersections with vertical lines:







Moving the punctures according to a braid generator changes some crossing numbers:



There is an explicit formula for the change in the coordinates! [Dyannikov (2002); Moussafir (2006); Hall & Yurttas (2009); J-LT (2010)]

For a specific rod motion, say as given by the braid  $\sigma_3^{-1}\sigma_2^{-1}\sigma_3^{-1}\sigma_2\sigma_1$ , we can easily see the exponential growth of  $L$  and thus measure the entropy:






$m$  is the number of times the braid acted on the initial loop.


[Moussafir, J.-O. (2006). *Func. Anal. and Other Math.* 1 (1), 37–46]

10 floats from Davis' Labrador sea data:




Floats have an entanglement time of about 50 days — timescale for horizontal stirring.

Source: WOCE subsurface float data assembly center (2004)



- There is a lot more information in the braid than just entropy;
- For instance: imagine there is an **isolated region** in the flow that does not interact with the rest, bounded by **Lagrangian coherent structures (LCS)**;
- Identify LCS and invariant regions from particle trajectory data by searching for curves that grow slowly or not at all.
- [see Haller, G. & Beron-Vera, F. J. (2012). *Physica D*, **241** (20), 1680–1702.]

# double-gyre coherent structures



play movie

[Allshouse, M. R. & J-LT (2012). *Physica D*, 241 (2), 95–105]



- The **nature of the isotopy** between the pA and real system.
- **Sharpness** of the entropy bound (Tumasz & J-LT, 2013).
- **Computational methods** for isotopy class (random entanglements of trajectories – LCS method, see Allshouse & J-LT (2012), ongoing work with Tom Peacock).
- **'Designing'** for topological chaos (see Stremler & Chen (2007)).
- Combine with **other measures**, e.g., **mix-norms** (Mathew *et al.*, 2005; Lin *et al.*, 2011; J-LT, 2012).
- We're developing a Matlab toolbox — **braidlab**.
- **3D?!** (lots of missing theory)



- Allshouse, M. R. & J-LT (2012). *Physica D*, **241** (2), 95–105.
- Bestvina, M. & Handel, M. (1995). *Topology*, **34** (1), 109–140.
- Binder, B. J. (2010). *Phys. Lett. A*, **374**, 3483–3486.
- Binder, B. J. & Cox, S. M. (2008). *Fluid Dyn. Res.* **40**, 34–44.
- Bowen, R. (1978). In: *Structure of Attractors* volume 668 of *Lecture Notes in Math.* pp. 21–29, New York: Springer.
- Boyland, P. L., Aref, H., & Stremler, M. A. (2000). *J. Fluid Mech.* **403**, 277–304.
- Boyland, P. L. & Harrington, J. (2011). *Algeb. Geom. Topology*, **11** (4), 2265–2296.
- Boyland, P. L., Stremler, M. A., & Aref, H. (2003). *Physica D*, **175**, 69–95.
- D'Alessandro, D., Dahleh, M., & Mezić, I. (1999). *IEEE Transactions on Automatic Control*, **44** (10), 1852–1863.
- Dynnikov, I. A. (2002). *Russian Math. Surveys*, **57** (3), 592–594.
- Finn, M. D. & J-LT (2011). *SIAM Rev.* **53** (4), 723–743.
- Gouillart, E., Finn, M. D., & J-LT (2006). *Phys. Rev. E*, **73**, 036311.
- Gouillart, E., Kuncio, N., Dauchot, O., Dubrulle, B., Roux, S., & J-LT, J.-L. (2007). *Phys. Rev. Lett.* **99**, 114501.
- Halbert, J. T. & Yorke, J. A. (2013). *Topology Proceedings*, . in press.



- Hall, T. & Yurttas, S. Ö. (2009). *Topology Appl.* **156** (8), 1554–1564.
- Haller, G. & Beron-Vera, F. J. (2012). *Physica D*, **241** (20), 1680–1702.
- Handel, M. (1985). *Ergod. Th. Dynam. Sys.* **8**, 373–377.
- Kobayashi, T. & Umeda, S. (2007). In: *Proceedings of the International Workshop on Knot Theory for Scientific Objects, Osaka, Japan* pp. 97–109, Osaka, Japan: Osaka Municipal Universities Press.
- Lin, Z., Doering, C. R., & J-LT (2011). *J. Fluid Mech.* **675**, 465–476.
- Mathew, G., Mezić, I., & Petzold, L. (2005). *Physica D*, **211** (1-2), 23–46.
- Moussafir, J.-O. (2006). *Func. Anal. and Other Math.* **1** (1), 37–46.
- Stremmer, M. A. & Chen, J. (2007). *Phys. Fluids*, **19**, 103602.
- Stremmer, M. A., Ross, S. D., Grover, P., & Kumar, P. (2011). *Phys. Rev. Lett.* **106**, 114101.
- J-LT (2005). *Phys. Rev. Lett.* **94** (8), 084502.
- J-LT (2010). *Chaos*, **20**, 017516.
- J-LT (2012). *Nonlinearity*, **25** (2), R1–R44.
- J-LT & Finn, M. D. (2006). *Phil. Trans. R. Soc. Lond. A*, **364**, 3251–3266.
- J-LT, Finn, M. D., Gouillart, E., & Hall, T. (2008). *Chaos*, **18**, 033123.
- J-LT, Gouillart, E., & Finn, M. D. (2009). In: *Analysis and Control of Mixing with Applications to Micro and Macro Flow Processes*, (Cortelezzi, L. & Mezić, I., eds) volume 510 of *CISM International Centre for Mechanical Sciences* pp. 339–350, Vienna: Springer.
- Thurston, W. P. (1988). *Bull. Am. Math. Soc.* **19**, 417–431.
- Tumasz, S. E. & J-LT (2013). *J. Nonlinear Sci.* .